



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 13, 2017 – 04:11 pm GMT

PDB ID : 3KMZ  
Title : Crystal structure of RARalpha ligand binding domain in complex with the inverse agonist BMS493 and a corepressor fragment  
Authors : Bourguet, W.; le Maire, A.  
Deposited on : 2009-11-11  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	266	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 8%, green 75%, grey 17%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>75%</span> <span>8%</span> <span>17%</span> </div> </div>
1	B	266	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 12%, green 70%, grey 17%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>70%</span> <span>12%</span> <span>17%</span> </div> </div>
2	C	19	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 89%, yellow 11%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span></span> <span>89%</span> <span>11%</span> </div> </div>
2	D	19	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 95%, yellow 5%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span></span> <span>95%</span> <span>5%</span> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	3	-	-	-	X
4	GOL	A	422	-	-	X	X
4	GOL	A	6	-	-	-	X
4	GOL	A	7	-	-	-	X
4	GOL	A	8	-	-	-	X
4	GOL	B	422	-	-	-	X
4	GOL	B	5	-	-	-	X
4	GOL	B	9	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4262 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Retinoic acid receptor alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	221	Total	C	N	O	S	0	4	0
			1761	1115	297	334	15			
1	A	221	Total	C	N	O	S	0	7	0
			1775	1124	298	338	15			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	156	MET	-	EXPRESSION TAG	UNP P10276
B	157	GLY	-	EXPRESSION TAG	UNP P10276
B	158	SER	-	EXPRESSION TAG	UNP P10276
B	159	SER	-	EXPRESSION TAG	UNP P10276
B	160	HIS	-	EXPRESSION TAG	UNP P10276
B	161	HIS	-	EXPRESSION TAG	UNP P10276
B	162	HIS	-	EXPRESSION TAG	UNP P10276
B	163	HIS	-	EXPRESSION TAG	UNP P10276
B	164	HIS	-	EXPRESSION TAG	UNP P10276
B	165	HIS	-	EXPRESSION TAG	UNP P10276
B	166	SER	-	EXPRESSION TAG	UNP P10276
B	167	SER	-	EXPRESSION TAG	UNP P10276
B	168	GLY	-	EXPRESSION TAG	UNP P10276
B	169	LEU	-	EXPRESSION TAG	UNP P10276
B	170	VAL	-	EXPRESSION TAG	UNP P10276
B	171	PRO	-	EXPRESSION TAG	UNP P10276
B	172	ARG	-	EXPRESSION TAG	UNP P10276
B	173	GLY	-	EXPRESSION TAG	UNP P10276
B	174	SER	-	EXPRESSION TAG	UNP P10276
B	175	HIS	-	EXPRESSION TAG	UNP P10276
A	156	MET	-	EXPRESSION TAG	UNP P10276
A	157	GLY	-	EXPRESSION TAG	UNP P10276
A	158	SER	-	EXPRESSION TAG	UNP P10276
A	159	SER	-	EXPRESSION TAG	UNP P10276
A	160	HIS	-	EXPRESSION TAG	UNP P10276

*Continued on next page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	161	HIS	-	EXPRESSION TAG	UNP P10276
A	162	HIS	-	EXPRESSION TAG	UNP P10276
A	163	HIS	-	EXPRESSION TAG	UNP P10276
A	164	HIS	-	EXPRESSION TAG	UNP P10276
A	165	HIS	-	EXPRESSION TAG	UNP P10276
A	166	SER	-	EXPRESSION TAG	UNP P10276
A	167	SER	-	EXPRESSION TAG	UNP P10276
A	168	GLY	-	EXPRESSION TAG	UNP P10276
A	169	LEU	-	EXPRESSION TAG	UNP P10276
A	170	VAL	-	EXPRESSION TAG	UNP P10276
A	171	PRO	-	EXPRESSION TAG	UNP P10276
A	172	ARG	-	EXPRESSION TAG	UNP P10276
A	173	GLY	-	EXPRESSION TAG	UNP P10276
A	174	SER	-	EXPRESSION TAG	UNP P10276
A	175	HIS	-	EXPRESSION TAG	UNP P10276

- | Mol | Chain | Residues | Atoms        |         |         |         |        | ZeroOcc | AltConf | Trace |
|-----|-------|----------|--------------|---------|---------|---------|--------|---------|---------|-------|
| 2   | C     | 19       | Total<br>156 | C<br>98 | N<br>29 | O<br>28 | S<br>1 | 0       | 0       | 0     |
| 2   | D     | 19       | Total<br>156 | C<br>98 | N<br>29 | O<br>28 | S<br>1 | 0       | 0       | 0     |

- # EQO
- 
- The chemical structure of EQO (Equisetol) is shown, featuring a complex polycyclic system. The structure includes a central benzene ring substituted with a carboxylic acid group (OAC, OH, OAD) and a side chain. The side chain consists of a propenyl group (CAG, CAH) attached to a cyclohexadiene ring (CAQ, CBA, CAU, CBE, CAV, CAI). This cyclohexadiene ring is further substituted with a CAA and CAB group, and a CCB group. The CCB group is attached to a benzene ring (CAR, CAO, CAZ, CAP, CAS). The CCB group is also attached to a side chain (CAE, CAF) which is further substituted with a CAY group. The CAY group is attached to a benzene ring (CAN, CAM, CAJ, CAL).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			31	29	2		
3	A	1	Total	C	O	0	0
			31	29	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	135	Total 135	O 135	0	0
5	C	15	Total 15	O 15	0	0
5	A	136	Total 136	O 136	0	0
5	D	12	Total 12	O 12	0	0





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.83Å 105.62Å 53.38Å 90.00° 89.92° 90.00°	Depositor
Resolution (Å)	43.40 – 2.10 43.39 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.6 (43.40-2.10) 96.2 (43.39-2.10)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.35 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.4.0062	Depositor
R, $R_{free}$	0.172 , 0.230 0.171 , 0.228	Depositor DCC
$R_{free}$ test set	1666 reflections (5.36%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.6	Xtriage
Anisotropy	0.183	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 31.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.027 for k,h,-l 0.026 for -k,-h,-l 0.467 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4262	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CSO, EQO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.48	0/1825	0.58	0/2466
1	B	0.48	0/1802	0.58	0/2435
2	C	0.46	0/149	0.72	1/199 (0.5%)
2	D	0.47	0/149	0.57	0/199
All	All	0.48	0/3925	0.59	1/5299 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2065	ARG	NE-CZ-NH2	-5.16	117.72	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1775	0	1828	16	0
1	B	1761	0	1811	27	0
2	C	156	0	160	3	0
2	D	156	0	160	1	0
3	A	31	0	23	0	0
3	B	31	0	23	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	30	0	40	7	0
4	B	24	0	32	9	0
5	A	136	0	0	0	0
5	B	135	0	0	0	0
5	C	15	0	0	0	0
5	D	12	0	0	0	0
All	All	4262	0	4077	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 43 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:367:ARG:HG3	4:B:9:GOL:H12	1.63	0.80
1:A:319:LEU:O	4:A:422:GOL:H31	1.88	0.73
1:A:347:ARG:HG2	1:A:347:ARG:HH11	1.54	0.72
1:B:186:GLU:O	1:B:190:LYS:HG3	1.99	0.62
1:B:218:VAL:H	1:B:299:ASN:HD21	1.47	0.60

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/266 (85%)	224 (99%)	2 (1%)	0	100	100
1	B	223/266 (84%)	220 (99%)	3 (1%)	0	100	100
2	C	16/19 (84%)	16 (100%)	0	0	100	100
2	D	16/19 (84%)	16 (100%)	0	0	100	100
All	All	481/570 (84%)	476 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/238 (86%)	200 (98%)	4 (2%)	60	66
1	B	201/238 (84%)	196 (98%)	5 (2%)	53	57
2	C	16/16 (100%)	16 (100%)	0	100	100
2	D	16/16 (100%)	16 (100%)	0	100	100
All	All	437/508 (86%)	428 (98%)	9 (2%)	62	64

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	394	ARG
1	A	385	ARG
1	A	254	ILE
1	B	265[B]	CYS
1	A	205	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	340	GLN
1	A	299	ASN
1	A	204	GLN
1	B	296	GLN
1	A	296	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	CSO	C	2056	2	4,6,7	1.77	1 (25%)	1,6,8	1.64	0
2	CSO	D	2056	2	4,6,7	1.69	1 (25%)	1,6,8	1.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSO	C	2056	2	-	0/1/5/7	0/0/0/0
2	CSO	D	2056	2	-	0/1/5/7	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2056	CSO	CA-C	3.21	1.54	1.50
2	C	2056	CSO	CA-C	3.43	1.54	1.50

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EQO	A	2	-	29,34,34	2.87	5 (17%)	42,48,48	1.99	8 (19%)
4	GOL	A	3	-	5,5,5	0.42	0	5,5,5	0.35	0
4	GOL	A	422	-	5,5,5	0.35	0	5,5,5	0.42	0
4	GOL	A	6	-	5,5,5	0.33	0	5,5,5	0.25	0
4	GOL	A	7	-	5,5,5	0.41	0	5,5,5	0.54	0
4	GOL	A	8	-	5,5,5	0.37	0	5,5,5	0.15	0
3	EQO	B	1	-	29,34,34	2.83	5 (17%)	42,48,48	1.94	6 (14%)
4	GOL	B	4	-	5,5,5	0.40	0	5,5,5	0.07	0
4	GOL	B	422	-	5,5,5	0.34	0	5,5,5	0.26	0
4	GOL	B	5	-	5,5,5	0.31	0	5,5,5	0.61	0
4	GOL	B	9	-	5,5,5	0.33	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EQO	A	2	-	-	0/8/30/30	0/4/4/4
4	GOL	A	3	-	-	0/4/4/4	0/0/0/0
4	GOL	A	422	-	-	0/4/4/4	0/0/0/0
4	GOL	A	6	-	-	0/4/4/4	0/0/0/0
4	GOL	A	7	-	-	0/4/4/4	0/0/0/0
4	GOL	A	8	-	-	0/4/4/4	0/0/0/0
3	EQO	B	1	-	-	0/8/30/30	0/4/4/4
4	GOL	B	4	-	-	0/4/4/4	0/0/0/0
4	GOL	B	422	-	-	0/4/4/4	0/0/0/0
4	GOL	B	5	-	-	0/4/4/4	0/0/0/0
4	GOL	B	9	-	-	0/4/4/4	0/0/0/0

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2	EQO	CAE-CAX	-11.23	1.29	1.44
3	B	1	EQO	CAE-CAX	-11.05	1.29	1.44
3	B	1	EQO	CAY-CAF	-6.04	1.29	1.44
3	A	2	EQO	CAY-CAF	-5.92	1.30	1.44
3	A	2	EQO	CAV-CBE	-5.60	1.46	1.54

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2	EQO	CAA-CBE-CAV	-5.50	100.01	109.15
3	B	1	EQO	CAA-CBE-CAV	-5.44	100.11	109.15
3	A	2	EQO	CAF-CAE-CAX	-3.05	168.52	176.07
3	A	2	EQO	CAS-CBB-CAW	-2.24	117.44	120.45
3	B	1	EQO	CBC-CAX-CAE	2.10	123.06	119.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	422	GOL	5	0
4	A	7	GOL	2	0
4	B	4	GOL	1	0
4	B	5	GOL	3	0
4	B	9	GOL	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	221/266 (83%)	-0.15	3 (1%) 75 79	19, 27, 52, 65	0
1	B	221/266 (83%)	-0.15	2 (0%) 84 86	18, 28, 53, 64	0
2	C	18/19 (94%)	-0.23	0 100 100	22, 27, 41, 45	0
2	D	18/19 (94%)	-0.12	0 100 100	22, 26, 42, 44	0
All	All	478/570 (83%)	-0.15	5 (1%) 82 85	18, 27, 53, 65	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	370	ARG	3.6
1	A	400	MET	3.3
1	A	181	THR	3.0
1	B	400	MET	2.4
1	B	371	PRO	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	CSO	C	2056	7/8	0.92	0.10	-	25,27,34,35	0
2	CSO	D	2056	7/8	0.96	0.09	-	23,25,31,32	0



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	B	422	6/6	0.87	0.19	6.36	61,62,64,66	0
4	GOL	A	8	6/6	0.89	0.25	6.19	56,57,57,57	0
4	GOL	A	6	6/6	0.88	0.24	4.55	66,67,70,71	0
4	GOL	A	7	6/6	0.84	0.30	4.04	54,55,57,58	1
4	GOL	A	422	6/6	0.88	0.43	3.56	68,69,70,70	0
4	GOL	A	3	6/6	0.94	0.16	2.79	37,41,42,43	0
4	GOL	B	5	6/6	0.92	0.17	2.37	43,46,47,51	0
3	EQO	A	2	31/31	0.94	0.13	1.87	18,24,34,36	0
4	GOL	B	4	6/6	0.81	0.18	1.67	61,63,64,65	0
4	GOL	B	9	6/6	0.95	0.24	1.46	62,63,65,65	0
3	EQO	B	1	31/31	0.95	0.11	0.27	19,23,31,33	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.