



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2017 – 06:06 AM EST

PDB ID : 4KMH  
Title : Crystal structure of Suppressor of Fused d20  
Authors : Zhang, Y.; Qi, X.; Zhang, Z.; Wu, G.  
Deposited on : unknown  
Resolution : 3.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2176 (3.08-3.00)
Clashscore	112137	2542 (3.08-3.00)
Ramachandran outliers	110173	2458 (3.08-3.00)
Sidechain outliers	110143	2461 (3.08-3.00)
RSRZ outliers	101464	2202 (3.08-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	484	
1	B	484	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6075 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Suppressor of fused homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	383	Total	C	N	O	S	0	0	0
			3012	1922	509	568	13			
1	B	389	Total	C	N	O	S	0	0	0
			3051	1948	515	575	13			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP Q9UMX1
A	-18	GLY	-	EXPRESSION TAG	UNP Q9UMX1
A	-17	SER	-	EXPRESSION TAG	UNP Q9UMX1
A	-16	SER	-	EXPRESSION TAG	UNP Q9UMX1
A	-15	HIS	-	EXPRESSION TAG	UNP Q9UMX1
A	-14	HIS	-	EXPRESSION TAG	UNP Q9UMX1
A	-13	HIS	-	EXPRESSION TAG	UNP Q9UMX1
A	-12	HIS	-	EXPRESSION TAG	UNP Q9UMX1
A	-11	HIS	-	EXPRESSION TAG	UNP Q9UMX1
A	-10	HIS	-	EXPRESSION TAG	UNP Q9UMX1
A	-9	SER	-	EXPRESSION TAG	UNP Q9UMX1
A	-8	SER	-	EXPRESSION TAG	UNP Q9UMX1
A	-7	GLY	-	EXPRESSION TAG	UNP Q9UMX1
A	-6	LEU	-	EXPRESSION TAG	UNP Q9UMX1
A	-5	VAL	-	EXPRESSION TAG	UNP Q9UMX1
A	-4	PRO	-	EXPRESSION TAG	UNP Q9UMX1
A	-3	ARG	-	EXPRESSION TAG	UNP Q9UMX1
A	-2	GLY	-	EXPRESSION TAG	UNP Q9UMX1
A	-1	SER	-	EXPRESSION TAG	UNP Q9UMX1
A	0	HIS	-	EXPRESSION TAG	UNP Q9UMX1
B	-19	MET	-	EXPRESSION TAG	UNP Q9UMX1
B	-18	GLY	-	EXPRESSION TAG	UNP Q9UMX1
B	-17	SER	-	EXPRESSION TAG	UNP Q9UMX1
B	-16	SER	-	EXPRESSION TAG	UNP Q9UMX1
B	-15	HIS	-	EXPRESSION TAG	UNP Q9UMX1

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	EXPRESSION TAG	UNP Q9UMX1
B	-13	HIS	-	EXPRESSION TAG	UNP Q9UMX1
B	-12	HIS	-	EXPRESSION TAG	UNP Q9UMX1
B	-11	HIS	-	EXPRESSION TAG	UNP Q9UMX1
B	-10	HIS	-	EXPRESSION TAG	UNP Q9UMX1
B	-9	SER	-	EXPRESSION TAG	UNP Q9UMX1
B	-8	SER	-	EXPRESSION TAG	UNP Q9UMX1
B	-7	GLY	-	EXPRESSION TAG	UNP Q9UMX1
B	-6	LEU	-	EXPRESSION TAG	UNP Q9UMX1
B	-5	VAL	-	EXPRESSION TAG	UNP Q9UMX1
B	-4	PRO	-	EXPRESSION TAG	UNP Q9UMX1
B	-3	ARG	-	EXPRESSION TAG	UNP Q9UMX1
B	-2	GLY	-	EXPRESSION TAG	UNP Q9UMX1
B	-1	SER	-	EXPRESSION TAG	UNP Q9UMX1
B	0	HIS	-	EXPRESSION TAG	UNP Q9UMX1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	6	Total O 6 6	0	0
2	B	6	Total O 6 6	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Suppressor of fused homolog



#### • Molecule 1: Suppressor of fused homolog



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.63Å 122.33Å 118.75Å 90.00° 90.41° 90.00°	Depositor
Resolution (Å)	50.00 – 3.04 42.60 – 3.04	Depositor EDS
% Data completeness (in resolution range)	97.5 (50.00-3.04) 97.5 (42.60-3.04)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.79 (at 3.06Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.224 , 0.277 0.215 , 0.242	Depositor DCC
$R_{free}$ test set	1010 reflections (5.36%)	DCC
Wilson B-factor (Å <sup>2</sup> )	83.4	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for -1/2*h+1/2*k,3/2*h+1/2*k,-l 0.021 for -1/2*h-1/2*k,-3/2*h+1/2*k,-l 0.019 for 1/2*h+1/2*k,3/2*h-1/2*k,-l 0.018 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.406 for -h,-k,l	Xtriage
Reported twinning fraction	0.632 for H,K,L 0.368 for -h,-k,l	Depositor
Outliers	0 of 19784 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6075	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/3098	0.52	1/4217 (0.0%)
1	B	0.33	0/3138	0.52	1/4274 (0.0%)
All	All	0.33	0/6236	0.52	2/8491 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	20	PRO	N-CA-CB	5.98	110.48	103.30
1	A	20	PRO	N-CA-CB	5.91	110.39	103.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3012	0	2876	41	0
1	B	3051	0	2914	40	0
2	A	6	0	0	0	0
2	B	6	0	0	0	0
All	All	6075	0	5790	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:278:LEU:HB2	1:A:368:SER:OG	1.45	1.14
1:B:278:LEU:HB2	1:B:368:SER:OG	1.59	1.02
1:A:371:LEU:HD11	1:A:433:ILE:CD1	1.90	1.01
1:B:449:ASP:HA	1:B:450:LEU:C	1.94	0.84
1:A:103:ARG:HG3	1:A:104:VAL:HG23	1.63	0.81
1:A:371:LEU:HD11	1:A:433:ILE:HD11	1.67	0.75
1:A:76:ASN:HD22	1:A:78:GLY:H	1.34	0.74
1:A:278:LEU:HB2	1:A:368:SER:HG	1.53	0.73
1:B:76:ASN:HD22	1:B:78:GLY:H	1.36	0.72
1:A:362:ARG:HG2	1:A:363:THR:HG23	1.73	0.71
1:A:449:ASP:HA	1:A:450:LEU:C	2.14	0.68
1:B:103:ARG:HG3	1:B:104:VAL:HG23	1.78	0.64
1:B:163:TRP:H	1:B:176:HIS:HD2	1.45	0.63
1:B:171:GLU:OE2	1:B:171:GLU:HA	1.96	0.63
1:A:163:TRP:H	1:A:176:HIS:HD2	1.46	0.62
1:B:215:ASN:HD22	1:B:218:GLY:H	1.50	0.59
1:A:180:THR:HG21	1:A:234:LEU:HD12	1.85	0.58
1:A:371:LEU:HD21	1:A:433:ILE:HD11	1.86	0.57
1:B:431:LEU:HD13	1:B:433:ILE:HG22	1.88	0.56
1:B:215:ASN:ND2	1:B:218:GLY:H	2.05	0.55
1:B:279:SER:O	1:B:280:ARG:CB	2.54	0.55
1:B:433:ILE:HD11	1:B:435:LEU:HD21	1.88	0.55
1:A:386:ARG:HA	1:A:390:LEU:HD12	1.89	0.53
1:B:163:TRP:HB3	1:B:175:GLN:O	2.09	0.53
1:B:136:TRP:CD2	1:B:168:ASP:HB3	2.45	0.52
1:A:374:ASN:ND2	1:A:377:SER:H	2.08	0.52
1:A:252:GLN:O	1:A:255:VAL:HG22	2.10	0.51
1:B:386:ARG:HA	1:B:390:LEU:HD12	1.92	0.50
1:B:168:ASP:CG	1:B:170:SER:OG	2.50	0.50
1:A:431:LEU:HD13	1:A:433:ILE:HG13	1.93	0.50
1:A:362:ARG:HG2	1:A:363:THR:N	2.26	0.49
1:B:278:LEU:CB	1:B:368:SER:OG	2.48	0.49
1:B:409:VAL:HB	1:B:413:VAL:HG21	1.95	0.48
1:B:164:HIS:H	1:B:176:HIS:CD2	2.32	0.48
1:A:215:ASN:HD22	1:A:218:GLY:H	1.62	0.47
1:A:221:GLU:HA	1:A:224:ARG:HH11	1.80	0.47
1:B:30:PHE:CD2	1:B:30:PHE:N	2.83	0.47
1:A:76:ASN:ND2	1:A:78:GLY:H	2.09	0.46
1:B:277:ASP:OD1	1:B:279:SER:HB2	2.14	0.46
1:A:136:TRP:CE2	1:A:168:ASP:HB3	2.51	0.46
1:B:389:LEU:HD11	1:B:408:PHE:HB3	1.97	0.46

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:431:LEU:CD1	1:A:433:ILE:HG13	2.46	0.46
1:A:371:LEU:HD11	1:A:433:ILE:HD13	1.89	0.46
1:A:164:HIS:H	1:A:176:HIS:CD2	2.34	0.45
1:A:271:ALA:HB1	1:A:377:SER:HB3	1.97	0.45
1:B:180:THR:HG21	1:B:234:LEU:HD13	1.97	0.45
1:B:229:ALA:HB3	1:B:230:GLY:HA3	1.98	0.45
1:A:90:TYR:O	1:A:121:THR:HA	2.16	0.45
1:A:222:LEU:HD22	1:A:251:LEU:HD22	1.99	0.45
1:B:403:ASP:HB3	1:B:429:PRO:HG3	1.99	0.45
1:B:278:LEU:HB2	1:B:368:SER:HG	1.78	0.44
1:B:384:CYS:O	1:B:388:ARG:HB2	2.17	0.44
1:B:102:ASN:OD1	1:B:106:GLU:HA	2.18	0.44
1:A:136:TRP:CZ2	1:A:168:ASP:HB3	2.53	0.43
1:A:166:PRO:HB2	1:A:170:SER:O	2.18	0.43
1:A:382:PRO:O	1:A:386:ARG:HB2	2.19	0.43
1:A:275:TRP:CE2	1:A:431:LEU:HB2	2.54	0.43
1:A:362:ARG:HE	1:A:363:THR:H	1.67	0.43
1:A:122:PHE:CE2	1:A:124:LEU:HB2	2.54	0.42
1:A:36:ALA:HB3	1:A:190:THR:HG22	2.01	0.42
1:A:77:VAL:HA	1:A:86:GLU:HG3	2.01	0.42
1:A:229:ALA:N	1:A:230:GLY:HA3	2.34	0.42
1:B:166:PRO:CB	1:B:170:SER:O	2.67	0.42
1:B:439:PHE:CZ	1:B:467:LYS:HG3	2.55	0.42
1:B:368:SER:HB2	1:B:468:LYS:HB3	2.01	0.42
1:B:30:PHE:HD2	1:B:30:PHE:N	2.18	0.41
1:A:31:PRO:HA	1:A:32:PRO:HD3	1.90	0.41
1:B:374:ASN:ND2	1:B:377:SER:H	2.19	0.41
1:A:383:LEU:HD12	1:A:383:LEU:C	2.41	0.41
1:A:380:LEU:HD22	1:A:383:LEU:HD23	2.02	0.41
1:B:226:VAL:O	1:B:230:GLY:HA3	2.20	0.41
1:B:22:ALA:O	1:B:42:ARG:NH2	2.45	0.41
1:B:143:GLY:HA2	1:B:146:ARG:HH11	1.85	0.41
1:B:31:PRO:HA	1:B:32:PRO:HD3	1.88	0.41
1:B:271:ALA:HB1	1:B:377:SER:HB3	2.02	0.41
1:B:448:GLU:O	1:B:449:ASP:C	2.59	0.41
1:A:447:LEU:HB3	1:A:450:LEU:HD22	2.03	0.41
1:A:374:ASN:HB2	1:A:476:ASP:OD1	2.21	0.41
1:B:22:ALA:HA	1:B:23:PRO:HD3	1.96	0.40
1:B:275:TRP:CE2	1:B:431:LEU:HB2	2.56	0.40
1:A:389:LEU:HD11	1:A:408:PHE:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	375/484 (78%)	364 (97%)	11 (3%)	0	100	100
1	B	381/484 (79%)	363 (95%)	15 (4%)	3 (1%)	22	61
All	All	756/968 (78%)	727 (96%)	26 (3%)	3 (0%)	38	75

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	172	SER
1	B	21	THR
1	B	174	ILE

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	325/418 (78%)	307 (94%)	18 (6%)	25	61
1	B	329/418 (79%)	313 (95%)	16 (5%)	29	66
All	All	654/836 (78%)	620 (95%)	34 (5%)	27	64

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	THR
1	A	76	ASN
1	A	125	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	128	THR
1	A	144	LEU
1	A	209	HIS
1	A	215	ASN
1	A	256	ASP
1	A	266	LEU
1	A	278	LEU
1	A	279	SER
1	A	362	ARG
1	A	364	ARG
1	A	368	SER
1	A	374	ASN
1	A	383	LEU
1	A	447	LEU
1	A	469	LEU
1	B	4	LEU
1	B	5	ARG
1	B	55	THR
1	B	76	ASN
1	B	125	LYS
1	B	215	ASN
1	B	266	LEU
1	B	278	LEU
1	B	363	THR
1	B	374	ASN
1	B	384	CYS
1	B	386	ARG
1	B	433	ILE
1	B	434	LEU
1	B	469	LEU
1	B	476	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	ASN
1	A	164	HIS
1	A	175	GLN
1	A	176	HIS
1	A	215	ASN
1	A	370	HIS
1	A	374	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	76	ASN
1	B	175	GLN
1	B	176	HIS
1	B	215	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	383/484 (79%)	0.14	2 (0%) 90 74	63, 82, 108, 149	15 (3%)
1	B	389/484 (80%)	0.22	3 (0%) 86 64	63, 82, 113, 202	15 (3%)
All	All	772/968 (79%)	0.18	5 (0%) 89 71	63, 82, 111, 202	30 (3%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	450	LEU	4.9
1	B	403	ASP	3.5
1	A	450	LEU	3.4
1	B	-1	SER	3.0
1	A	279	SER	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.