



wwPDB NMR Structure Validation Summary Report ⓘ

Feb 12, 2017 – 11:10 pm GMT

PDB ID : 2KN7
Title : Structure of the XPF-single strand DNA complex
Authors : Das, D.; Folkers, G.E.; van Dijk, M.; Jaspers, N.G.J.; Hoeijmakers, J.H.J.;
Kaptein, R.; Boelens, R.
Deposited on : 2009-08-16

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk28760
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 63%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	67	 66% 25% 7%
1	D	67	 63% 22% 12%
2	B	10	 40% 50% 10%
2	C	10	 40% 50% 10%

2 Ensemble composition and analysis

This entry contains 20 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:12-A:73, D:115-D:173 (121)	0.17	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters. No single-model clusters were found.

Cluster number	Models
1	3, 5, 6, 8, 12, 15, 16, 17
2	1, 4, 13, 20
3	9, 10, 14, 18
4	7, 19
5	2, 11

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2720 atoms, of which 1268 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called DNA repair endonuclease XPF.

Mol	Chain	Residues	Atoms						Trace
1	A	67	Total	C	H	N	O	S	0
			1041	329	520	90	99	3	
1	D	67	Total	C	H	N	O	S	0
			1041	329	520	90	99	3	

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*AP*GP*TP*GP*GP*CP*TP*GP*A)-3').


Mol	Chain	Residues	Atoms						Trace
2	B	10	Total	C	H	N	O	P	0
			319	98	114	40	58	9	
2	C	10	Total	C	H	N	O	P	0
			319	98	114	40	58	9	

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: DNA repair endonuclease XPF




- Molecule 1: DNA repair endonuclease XPF



- Molecule 2: DNA (5'-D(*CP*AP*GP*TP*GP*GP*CP*TP*GP*A)-3')



- Molecule 2: DNA (5'-D(*CP*AP*GP*TP*GP*GP*CP*TP*GP*A)-3')



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 5. Colouring as in section 4.1 above.


- Molecule 1: DNA repair endonuclease XPF

Chain A:  70% 19% • 7%





- Molecule 1: DNA repair endonuclease XPF

Chain D:  58% 27% •• 12%




- Molecule 2: DNA (5'-D(*CP*AP*GP*TP*GP*GP*CP*TP*GP*A)-3')

Chain B:  80% 20%



- Molecule 2: DNA (5'-D(*CP*AP*GP*TP*GP*GP*CP*TP*GP*A)-3')

Chain C:  10% 80% 10%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing, torsion angle dynamics*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
CNS	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 16449
Number of chemical shift lists	1
Total number of shifts	1414
Number of shifts mapped to atoms	1408
Number of unparsed shifts	0
Number of shifts with mapping errors	6
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	63%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.29±0.01	0±0/492 (0.0±0.0%)	0.38±0.01	0±0/667 (0.0±0.0%)
1	D	0.29±0.01	0±0/463 (0.0±0.0%)	0.37±0.01	0±0/626 (0.0±0.0%)
2	B	0.73±0.01	0±0/230 (0.0±0.0%)	0.98±0.01	2±0/354 (0.6±0.0%)
2	C	0.73±0.01	0±0/230 (0.0±0.0%)	0.98±0.01	2±0/354 (0.6±0.0%)
All	All	0.48	0/28300 (0.0%)	0.66	80/40020 (0.2%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	9	DG	N3-C2-N2	-6.39	115.42	119.90	12	20
2	C	9	DG	N3-C2-N2	-6.33	115.47	119.90	11	20
2	C	3	DG	N3-C2-N2	-5.77	115.86	119.90	19	20
2	B	3	DG	N3-C2-N2	-5.74	115.88	119.90	1	20

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	481	479	476	10±3
1	D	454	457	454	11±2
2	B	205	114	114	7±3

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes
2	C	205	114	114	7±2
All	All	26900	23280	23160	591

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

5 of 126 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:3:DG:H2''	2:B:4:DT:O5'	0.94	1.62	10	18
2:C:3:DG:H2''	2:C:4:DT:O5'	0.91	1.63	18	20
2:B:5:DG:N3	2:B:5:DG:H5''	0.90	1.81	11	4
2:C:5:DG:N3	2:C:5:DG:H5''	0.89	1.82	11	6
2:B:5:DG:H5''	2:B:5:DG:N3	0.75	1.97	12	4

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	62/67 (93%)	58±1 (94±1%)	3±1 (4±1%)	1±0 (2±0%)	15	58
1	D	59/67 (88%)	57±1 (96±1%)	1±1 (2±1%)	1±0 (2±0%)	15	58
All	All	2420/2680 (90%)	2295 (95%)	84 (3%)	41 (2%)	15	58

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	D	140	ASN	20
1	A	40	ASN	20
1	A	12	TYR	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	52/56 (93%)	43±2 (83±4%)	9±2 (17±4%)	5	41
1	D	49/56 (88%)	40±1 (82±3%)	9±1 (18±3%)	5	39
All	All	2020/2240 (90%)	1663 (82%)	357 (18%)	5	40

5 of 43 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	17	GLN	20
1	A	52	LEU	20
1	A	53	THR	20
1	D	117	GLN	20
1	A	27	ASN	20

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 63% for the well-defined parts and 62% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 16449

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1414
Number of shifts mapped to atoms	1408
Number of unparsed shifts	0
Number of shifts with mapping errors	6
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	6

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Residue not found in structure. First 5 (of 6) occurrences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
D	75	LYS	H	8.143	0.0	1
D	75	LYS	CB	35.026	0.0	1
A	75	LYS	CB	35.026	0.0	1
A	75	LYS	N	128.74	0.0	1
D	75	LYS	N	128.74	0.0	1

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	124	-1.81 ± 0.14	Should be applied
$^{13}\text{C}_\beta$	126	-1.14 ± 0.12	Should be applied
$^{13}\text{C}'$	2	—	None (insufficient data)

Continued on next page...

Continued from previous page...

Nucleus	# values	Correction \pm precision, ppm	Suggested action
^{15}N	122	1.00 \pm 0.40	Should be applied

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 63%, i.e. 1180 atoms were assigned a chemical shift out of a possible 1868. 16 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	470/595 (79%)	236/237 (100%)	119/242 (49%)	115/116 (99%)
Sidechain	642/747 (86%)	394/438 (90%)	231/278 (83%)	17/31 (55%)
Aromatic	68/126 (54%)	46/66 (70%)	16/48 (33%)	6/12 (50%)
Overall	1180/1868 (63%)	676/981 (69%)	366/700 (52%)	138/187 (74%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
2	D	160	ALA	C	56.63	188.57 – 166.97	-56.1
1	A	60	ALA	C	56.63	188.57 – 166.97	-56.1
2	D	112	TYR	CE1	134.40	124.14 – 111.74	13.3
1	A	12	TYR	CE1	134.40	124.14 – 111.74	13.3
1	A	12	TYR	CE2	134.40	124.68 – 111.18	12.2
2	D	112	TYR	CE2	134.40	124.68 – 111.18	12.2

7.1.5 Random Coil Index (RCI) plots [i](#)

The images below report *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain D:

