



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2017 – 02:44 PM EST

PDB ID : 4KNZ  
Title : Thymidylate synthase ternary complex with dUMP AND CB3717  
Authors : Newby, Z.; Lee, T.T.; Finer-Moore, J.; Stroud, R.M.  
Deposited on : unknown  
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1131 (1.32-1.28)
Clashscore	112137	1185 (1.32-1.28)
Ramachandran outliers	110173	1138 (1.32-1.28)
Sidechain outliers	110143	1138 (1.32-1.28)
RSRZ outliers	101464	1133 (1.32-1.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	<div> <div>2%</div> <div>90%</div> <div>9%</div> </div>
1	B	264	<div> <div>11%</div> <div>94%</div> <div>6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NA	A	903	-	-	-	X

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9681 atoms, of which 4336 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thymidylate synthase.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	264	Total	C	H	N	O	S	0	9	0
			4385	1418	2165	385	404	13			
1	B	264	Total	C	H	N	O	S	0	6	0
			4306	1395	2129	371	398	13			

- Molecule 2 is 2'-deoxy-5'-uridylic acid (three-letter code: UMC) (formula:  $C_9H_{15}N_2O_8P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	B	1	Total	C	N	O	P	0	0
			20	9	2	8	1		

- Molecule 3 is 10-PROPARGYL-5,8-DIDEAZAFOLIC ACID (three-letter code: CB3) (formula:  $C_{24}H_{23}N_5O_6$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			56	24	21	5	6		
3	B	1	Total	C	H	N	O	0	0
			56	24	21	5	6		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	466	Total	O	0	1
			466	466		
5	B	371	Total	O	0	0
			371	371		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Thymidylate synthase



#### • Molecule 1: Thymidylate synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	125.41Å 125.41Å 66.58Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.60 – 1.30 45.65 – 1.21	Depositor EDS
% Data completeness (in resolution range)	96.5 (45.60-1.30) 92.2 (45.65-1.21)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.97 (at 1.21Å)	Xtriage
Refinement program	ELVES, PHENIX (PHENIX.REFINE: 1.7.2_869)	Depositor
R, $R_{free}$	0.131 , 0.154 0.140 , 0.161	Depositor DCC
$R_{free}$ test set	14036 reflections (11.04%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.4	Xtriage
Anisotropy	0.175	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 46.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.025 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	9681	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CB3, CXM, UMC, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.44	0/2281	0.68	0/3097
1	B	0.40	0/2248	0.63	0/3052
All	All	0.42	0/4529	0.65	0/6149

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2220	2165	2151	24	0
1	B	2177	2129	2114	16	0
2	A	20	0	12	0	0
2	B	20	0	12	0	0
3	A	35	21	21	2	0
3	B	35	21	21	4	0
4	A	1	0	0	0	0
5	A	466	0	0	15	1
5	B	371	0	0	4	1
All	All	5345	4336	4331	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 5.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:CYS:SG	5:A:1323:HOH:O	1.97	0.98
1:A:64:GLN:NE2	5:A:1338:HOH:O	2.07	0.87
1:A:229:LYS:HD2	5:A:1429:HOH:O	1.82	0.79
1:A:52:LEU:HD22	1:A:249:ILE:HG13	1.63	0.79
1:B:108:HIS:O	5:B:646:HOH:O	2.08	0.71
1:B:116:LEU:HD22	1:B:188[A]:MET:SD	2.31	0.70
1:A:103[A]:THR:O	1:A:105[A]:ASP:N	2.25	0.68
1:B:116:LEU:HD22	1:B:188[A]:MET:HE1	1.74	0.68
1:B:116:LEU:HD22	1:B:188[A]:MET:CE	2.24	0.68
1:B:116:LEU:CD2	1:B:188[A]:MET:SD	2.84	0.65
1:B:116:LEU:CD2	1:B:188[A]:MET:CE	2.79	0.60
1:B:139[B]:ASP:OD2	5:B:749:HOH:O	2.16	0.60
1:A:52:LEU:CD2	1:A:249:ILE:HG13	2.31	0.59
1:A:102:PRO:O	5:A:1324:HOH:O	2.18	0.55
1:A:53[B]:ARG:HG3	1:A:244:PHE:CZ	2.43	0.53
1:A:190:GLN:CG	5:A:1318:HOH:O	2.56	0.53
1:A:190:GLN:NE2	5:A:1318:HOH:O	2.43	0.51
1:B:116:LEU:CD2	1:B:188[A]:MET:HE1	2.40	0.51
1:B:183:LEU:HD11	1:B:247:PHE:CE2	2.47	0.50
1:B:17:GLN:NE2	5:B:674:HOH:O	2.46	0.49
3:B:302:CB3:C6	3:B:302:CB3:H13	2.44	0.48
1:A:104[A]:PRO:CD	5:A:1229:HOH:O	2.62	0.47
1:A:237:GLU:HG2	1:A:241:ASP:OD2	2.15	0.47
1:A:190:GLN:HG3	5:A:1318:HOH:O	2.15	0.46
1:B:103:THR:OG1	1:B:105:ASP:OD2	2.18	0.46
1:A:129:ILE:HD12	1:B:133:TRP:CH2	2.51	0.45
1:B:107:ARG:HB3	5:B:646:HOH:O	2.16	0.45
5:A:1324:HOH:O	1:B:104:PRO:HG3	2.16	0.45
1:A:183:LEU:HD11	1:A:247:PHE:CE2	2.52	0.45
1:A:190:GLN:CD	5:A:1318:HOH:O	2.54	0.44
1:B:116:LEU:HD21	1:B:188[A]:MET:SD	2.57	0.44
1:A:152:PHE:CZ	1:A:188[A]:MET:SD	3.11	0.44
1:A:259:LYS:NZ	5:A:1360:HOH:O	2.37	0.44
1:A:104[A]:PRO:HD2	5:A:1229:HOH:O	2.19	0.43
1:A:223:GLU:HG3	5:A:1167:HOH:O	2.18	0.43
3:B:302:CB3:C6	3:B:302:CB3:C13	2.96	0.43
1:A:228:PRO:HB2	1:A:249:ILE:HG23	2.01	0.43
3:B:302:CB3:C14	3:B:302:CB3:C5	2.96	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:902:CB3:C6	3:A:902:CB3:H15	2.49	0.42
3:A:902:CB3:C14	3:A:902:CB3:C5	2.98	0.41
1:A:61:TRP:CD1	1:A:66:ASP:HB3	2.55	0.41
1:A:243:ARG:HD2	5:A:1348:HOH:O	2.19	0.41
1:A:50:CYS:CB	5:A:1323:HOH:O	2.54	0.41
3:B:302:CB3:C5	3:B:302:CB3:C13	2.99	0.41
1:B:168:CYS:SG	1:B:174:LEU:HB2	2.61	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1369:HOH:O	5:B:701:HOH:O[6_544]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/264 (103%)	268 (99%)	2 (1%)	1 (0%)	38	11
1	B	268/264 (102%)	263 (98%)	4 (2%)	1 (0%)	38	11
All	All	539/528 (102%)	531 (98%)	6 (1%)	2 (0%)	38	11

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	93	VAL
1	B	93	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/232 (104%)	239 (99%)	2 (1%)	85	56
1	B	238/232 (103%)	238 (100%)	0	100	100
All	All	479/464 (103%)	477 (100%)	2 (0%)	94	75

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105[A]	ASP
1	A	105[B]	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	217	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CXM	A	1	1	7,10,11	0.96	1 (14%)	5,11,13	1.40	1 (20%)
1	CXM	B	1	1	7,10,11	0.86	0	5,11,13	1.28	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CXM	A	1	1	-	0/6/10/12	0/0/0/0
1	CXM	B	1	1	-	0/6/10/12	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	CXM	CA-C	2.21	1.53	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	CXM	CB-CA-C	-2.34	107.80	111.65
1	B	1	CXM	O-C-CA	-2.06	120.34	125.15

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	UMC	A	901	1	21,21,21	0.94	1 (4%)	30,31,31	1.23	2 (6%)
3	CB3	A	902	-	29,37,37	1.50	5 (17%)	38,51,51	1.86	5 (13%)
2	UMC	B	301	1	21,21,21	0.87	0	30,31,31	1.23	3 (10%)
3	CB3	B	302	-	29,37,37	1.42	4 (13%)	38,51,51	1.76	7 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UMC	A	901	1	-	0/10/35/35	0/2/2/2
3	CB3	A	902	-	-	0/21/28/28	0/3/3/3
2	UMC	B	301	1	-	0/10/35/35	0/2/2/2
3	CB3	B	302	-	-	0/21/28/28	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	902	CB3	C4A-C8A	-3.12	1.39	1.41
2	A	901	UMC	C2-N3	-2.23	1.33	1.38
3	A	902	CB3	C4-N3	2.66	1.37	1.33
3	A	902	CB3	CP1-CP2	2.72	1.50	1.47
3	B	302	CB3	C4-N3	2.94	1.38	1.33
3	B	302	CB3	CP1-CP2	3.15	1.50	1.47
3	A	902	CB3	O-C	3.20	1.30	1.23
3	A	902	CB3	C2-N3	3.45	1.37	1.33
3	B	302	CB3	O-C	3.46	1.30	1.23
3	B	302	CB3	C2-N3	3.50	1.37	1.33

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	902	CB3	C4A-C4-N3	-8.34	118.57	124.45
3	B	302	CB3	C4A-C4-N3	-7.31	119.29	124.45
2	A	901	UMC	C6-N1-C1'	-4.00	114.85	119.75

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	CB3	C6-C9-N10	-3.89	108.05	114.27
3	A	902	CB3	C6-C9-N10	-3.83	108.15	114.27
2	B	301	UMC	C6-N1-C1'	-3.17	115.86	119.75
2	B	301	UMC	C5-C6-N1	-2.82	107.77	110.70
3	A	902	CB3	C8-C8A-C4A	-2.42	120.18	121.67
2	B	301	UMC	O2-C2-N1	-2.30	120.24	123.12
3	B	302	CB3	CB-CA-CT	-2.23	109.05	112.28
3	B	302	CB3	C8-C8A-C4A	-2.18	120.33	121.67
3	B	302	CB3	C15-C14-N10	-2.18	118.33	121.39
2	A	901	UMC	C5-C4-N3	2.52	119.23	116.72
3	B	302	CB3	C9-N10-C14	2.86	126.10	120.89
3	A	902	CB3	C9-N10-C14	3.16	126.65	120.89
3	B	302	CB3	C4-N3-C2	3.36	121.23	116.17
3	A	902	CB3	C4-N3-C2	3.82	121.92	116.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	902	CB3	2	0
3	B	302	CB3	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	263/264 (99%)	0.20	4 (1%) 74 74	10, 14, 24, 32	0
1	B	263/264 (99%)	0.55	29 (11%) 6 6	11, 18, 31, 46	0
All	All	526/528 (99%)	0.38	33 (6%) 21 20	10, 16, 27, 46	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	104	PRO	10.1
1	B	105	ASP	7.2
1	B	103	THR	6.9
1	A	104[A]	PRO	6.2
1	B	107	ARG	5.1
1	B	86	GLU	4.9
1	B	106	GLY	4.3
1	B	108	HIS	4.0
1	B	70	ALA	3.7
1	B	244	PHE	3.4
1	B	231	ILE	3.3
1	A	103[A]	THR	3.2
1	A	105[A]	ASP	3.1
1	B	88	GLY	2.9
1	A	51[A]	HIS	2.9
1	B	249	ILE	2.9
1	B	232	ILE	2.7
1	B	74	GLU	2.6
1	B	193	ASP	2.6
1	B	102	PRO	2.6
1	B	76	ASN	2.4
1	B	73	HIS	2.4
1	B	52[A]	LEU	2.3
1	B	109	ILE	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	233	LYS	2.3
1	B	71	TYR	2.3
1	B	156	ASP	2.2
1	B	68	ASN	2.2
1	B	87	ASN	2.1
1	B	53	ARG	2.1
1	B	248	GLU	2.1
1	B	195	GLU	2.1
1	B	69	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	CXM	A	1	11/12	0.99	0.08	-	11,14,19,19	0
1	CXM	B	1	11/12	0.98	0.07	-	15,18,22,22	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	NA	A	903	1/1	0.93	0.31	3.11	29,29,29,29	0
3	CB3	B	302	35/35	0.94	0.09	0.54	14,19,28,30	0
2	UMC	A	901	20/20	0.98	0.09	0.33	10,11,13,13	0
2	UMC	B	301	20/20	0.97	0.09	0.20	12,13,16,16	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CB3	A	902	35/35	0.95	0.09	0.02	11,14,23,26	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.