



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 10:18 pm GMT

PDB ID : 5KPP  
Title : Structure of human PARP1 catalytic domain bound to a quinazoline-2,4(1H,3H)-dione inhibitor  
Authors : Cao, R.; Wang, Y.L.; Zhou, J.; Huang, N.; Xu, B.L.  
Deposited on : 2016-07-05  
Resolution : 2.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtrriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1570 (2.36-2.32)
Clashscore	112137	1673 (2.36-2.32)
Ramachandran outliers	110173	1654 (2.36-2.32)
Sidechain outliers	110143	1655 (2.36-2.32)
RSRZ outliers	101464	1576 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	350	
1	B	350	



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			34	23	4	4	3		
2	B	1	Total	C	F	N	O	0	0
			34	23	4	4	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	111	Total	O	0	0
			111	111		
3	B	63	Total	O	0	0
			63	63		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Poly [ADP-ribose] polymerase 1



- Molecule 1: Poly [ADP-ribose] polymerase 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.66Å 92.69Å 163.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	81.74 – 2.33 27.25 – 2.33	Depositor EDS
% Data completeness (in resolution range)	98.4 (81.74-2.33) 98.5 (27.25-2.33)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.34 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 5.8.0123	Depositor
R, $R_{free}$	0.239 , 0.284 0.240 , 0.281	Depositor DCC
$R_{free}$ test set	1578 reflections (5.22%)	DCC
Wilson B-factor (Å <sup>2</sup> )	40.4	Xtrriage
Anisotropy	0.550	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 33.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5752	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6WZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/2818	0.57	0/3802
1	B	0.35	0/2818	0.56	0/3802
All	All	0.35	0/5636	0.56	0/7604

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	3
All	All	0	5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	746	MET	Peptide
1	A	747	LYS	Peptide
1	B	745	GLY	Peptide
1	B	746	MET	Peptide
1	B	747	LYS	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2755	0	2790	1	0
1	B	2755	0	2790	2	0
2	A	34	0	0	0	0
2	B	34	0	0	0	0
3	A	111	0	0	0	0
3	B	63	0	0	0	0
All	All	5752	0	5580	3	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:962:ILE:HD12	1:A:971:LEU:HD11	1.87	0.57
1:B:872:ILE:HG21	1:B:920:LEU:HD11	1.96	0.46
1:B:744:PHE:CD2	1:B:747:LYS:HB3	2.51	0.45

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	348/350 (99%)	342 (98%)	4 (1%)	2 (1%)	28	30
1	B	348/350 (99%)	334 (96%)	12 (3%)	2 (1%)	28	30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	696/700 (99%)	676 (97%)	16 (2%)	4 (1%)	28 30

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	663	SER
1	A	663	SER
1	B	747	LYS
1	A	748	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	309/307 (101%)	305 (99%)	4 (1%)	73 84
1	B	309/307 (101%)	304 (98%)	5 (2%)	68 79
All	All	618/614 (101%)	609 (98%)	9 (2%)	70 80

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	700	LYS
1	A	817	TYR
1	A	832	GLU
1	A	971	LEU
1	B	667	LYS
1	B	744	PHE
1	B	817	TYR
1	B	940	LYS
1	B	968	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	868	ASN
1	A	961	ASN
1	B	868	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	6WZ	A	1101	-	33,37,37	1.74	5 (15%)	44,55,55	2.07	6 (13%)
2	6WZ	B	1101	-	33,37,37	1.72	4 (12%)	44,55,55	2.27	7 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6WZ	A	1101	-	-	0/16/30/30	0/4/4/4
2	6WZ	B	1101	-	-	0/16/30/30	0/4/4/4

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	6WZ	CAY-CAV	-6.37	1.40	1.50
2	B	1101	6WZ	CAY-CAV	-6.18	1.41	1.50
2	A	1101	6WZ	CAR-CAW	-4.42	1.41	1.51
2	B	1101	6WZ	CAR-CAW	-4.06	1.42	1.51
2	A	1101	6WZ	CBC-NBG	-2.02	1.37	1.40
2	A	1101	6WZ	CAT-CBH	2.50	1.53	1.50
2	B	1101	6WZ	CAT-CBH	2.69	1.54	1.50
2	B	1101	6WZ	CAZ-NAU	3.60	1.39	1.33
2	A	1101	6WZ	CAZ-NAU	3.62	1.39	1.33

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1101	6WZ	CBB-CAZ-NAU	-8.93	118.15	124.45
2	A	1101	6WZ	CBB-CAZ-NAU	-8.54	118.43	124.45
2	B	1101	6WZ	CAL-CAX-CAY	-2.53	120.15	123.13
2	A	1101	6WZ	CAL-CAX-CAY	-2.46	120.22	123.13
2	B	1101	6WZ	OAB-CAV-CAY	-2.19	115.71	120.07
2	A	1101	6WZ	CAP-CAQ-NBF	2.44	115.52	110.63
2	A	1101	6WZ	CAR-NBG-CBC	2.58	123.81	120.17
2	B	1101	6WZ	CAP-CAQ-NBF	2.91	116.47	110.63
2	B	1101	6WZ	CAY-CAV-NBE	3.19	123.58	118.39
2	A	1101	6WZ	CAM-CBB-CBC	4.94	120.32	118.04
2	A	1101	6WZ	CAZ-NAU-CBA	5.95	120.36	115.16
2	B	1101	6WZ	CAM-CBB-CBC	6.10	120.85	118.04
2	B	1101	6WZ	CAZ-NAU-CBA	6.28	120.65	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	349/350 (99%)	0.46	24 (6%) 18 25	26, 39, 64, 83	0
1	B	349/350 (99%)	1.20	74 (21%) 1 2	32, 53, 87, 126	0
All	All	698/700 (99%)	0.83	98 (14%) 3 5	26, 45, 81, 126	0

All (98) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	746	MET	16.2
1	B	783	ASP	8.2
1	B	785	SER	7.6
1	B	1011	THR	7.5
1	B	726	ASP	7.1
1	B	723	GLY	6.4
1	B	853	GLN	6.3
1	B	745	GLY	6.2
1	B	747	LYS	5.9
1	B	725	SER	5.9
1	B	662	LYS	5.7
1	B	748	LYS	5.3
1	B	976	SER	4.9
1	B	705	GLN	4.8
1	A	747	LYS	4.8
1	A	807	ASP	4.7
1	B	849	LYS	4.6
1	B	963	SER	4.2
1	B	744	PHE	4.2
1	B	846	GLN	4.2
1	B	829	TYR	4.2
1	A	727	SER	4.1
1	B	787	LYS	4.0
1	A	980	ASN	3.9

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	1009	PHE	3.9
1	B	897	PHE	3.8
1	B	912	GLN	3.8
1	A	846	GLN	3.7
1	B	975	ILE	3.7
1	A	726	ASP	3.6
1	B	959	SER	3.6
1	B	807	ASP	3.6
1	A	920	LEU	3.6
1	B	786	SER	3.5
1	B	855	HIS	3.5
1	A	748	LYS	3.4
1	B	784	ASP	3.4
1	B	980	ASN	3.3
1	B	664	LYS	3.3
1	B	782	SER	3.3
1	B	974	GLY	3.3
1	B	827	ASN	3.3
1	B	922	GLY	3.2
1	B	961	ASN	3.2
1	B	920	LEU	3.2
1	B	825	THR	3.1
1	B	914	ASP	3.1
1	B	913	GLY	3.1
1	B	781	GLY	3.0
1	B	1002	LEU	3.0
1	B	1010	LYS	3.0
1	B	981	ASP	3.0
1	B	921	LEU	3.0
1	B	860	LEU	2.9
1	A	959	SER	2.9
1	B	939	SER	2.9
1	B	730	LEU	2.9
1	A	721	SER	2.9
1	B	704	ARG	2.9
1	A	853	GLN	2.9
1	A	785	SER	2.8
1	A	922	GLY	2.7
1	B	958	PRO	2.7
1	A	725	SER	2.7
1	A	861	TRP	2.7
1	B	832	GLU	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	830	ASP	2.6
1	B	759	GLN	2.6
1	B	852	LYS	2.6
1	B	940	LYS	2.6
1	B	756	ASP	2.6
1	A	860	LEU	2.5
1	B	850	PRO	2.5
1	B	954	THR	2.5
1	B	982	THR	2.5
1	B	826	HIS	2.4
1	A	919	ILE	2.4
1	B	978	GLY	2.4
1	B	843	GLY	2.4
1	B	872	ILE	2.3
1	B	778	LEU	2.3
1	B	962	ILE	2.3
1	B	910	THR	2.3
1	B	751	LEU	2.3
1	B	960	ALA	2.3
1	A	849	LYS	2.2
1	A	809	GLU	2.2
1	A	730	LEU	2.2
1	A	744	PHE	2.2
1	A	898	ALA	2.1
1	B	979	VAL	2.1
1	B	822	HIS	2.1
1	B	845[A]	CYS	2.1
1	B	861	TRP	2.1
1	A	806	ARG	2.0
1	B	953	LYS	2.0
1	A	981	ASP	2.0
1	B	684	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	6WZ	B	1101	34/34	0.92	0.14	-0.56	35,40,46,48	0
2	6WZ	A	1101	34/34	0.93	0.14	-0.72	25,27,30,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.