



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 17, 2017 – 07:42 AM EDT

PDB ID : 5KQH  
Title : Crystal structure of the V293D variant of catalase-peroxidase from *B. pseudo-mallei*  
Authors : Loewen, P.C.  
Deposited on : unknown  
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	5868 (1.84-1.80)
Clashscore	112137	6856 (1.84-1.80)
Ramachandran outliers	110173	6780 (1.84-1.80)
Sidechain outliers	110143	6780 (1.84-1.80)
RSRZ outliers	101464	5947 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	728	<div> <div>%</div> <div> <div></div> <div>87%</div> <div>10% ..</div> </div> </div>
1	B	728	<div> <div>%</div> <div> <div></div> <div>89%</div> <div>8% ...</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	MPD	A	806	-	-	-	X
7	MPD	A	807	-	-	-	X
7	MPD	B	805	-	-	-	X

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 12795 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Catalase-peroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	713	Total	C	N	O	S	0	8	0
			5566	3510	991	1051	14			
1	B	713	Total	C	N	O	S	0	6	0
			5544	3498	986	1046	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	293	ASP	VAL	conflict	UNP Q3JNW6
B	293	ASP	VAL	conflict	UNP Q3JNW6

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	Fe	N	O	
			43	34	1	4	4	
							0	0


- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na		
			1	1	0	0
3	A	1	Total	Na		
			1	1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl		
			1	1	0	0

- Molecule 5 is OXYGEN MOLECULE (three-letter code: OXY) (formula: O<sub>2</sub>).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O		
			2	2	0	0
5	B	1	Total	O		
			2	2	0	0

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	P	0	0
			5	4	1		
6	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 7 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula:  $C_6H_{14}O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			8	6	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			8	6	2		
7	A	1	Total	C	O	0	0
			8	6	2		
7	B	1	Total	C	O	0	0
			8	6	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	786	Total	O	0	0
			786	786		
8	B	764	Total	O	0	0
			764	764		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Catalase-peroxidase



#### • Molecule 1: Catalase-peroxidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.79Å 113.43Å 174.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.56 – 1.82 47.56 – 1.82	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.56-1.82) 99.8 (47.56-1.82)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.67 (at 1.82Å)	Xtriage
Refinement program	REFMAC 5.8.0151	Depositor
R, $R_{free}$	0.154 , 0.184 0.165 , 0.189	Depositor DCC
$R_{free}$ test set	8889 reflections (5.23%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.4	Xtriage
Anisotropy	0.222	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 46.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12795	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: TOX, OXY, CL, NA, PO4, MPD, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.37	22/5694 (0.4%)	1.24	55/7737 (0.7%)
1	B	1.36	15/5672 (0.3%)	1.24	52/7709 (0.7%)
All	All	1.36	37/11366 (0.3%)	1.24	107/15446 (0.7%)

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	288	GLY	C-O	11.56	1.42	1.23
1	B	288	GLY	C-O	10.73	1.40	1.23
1	B	410	GLU	CG-CD	9.17	1.65	1.51
1	A	544	GLN	CD-OE1	8.24	1.42	1.24
1	B	512	ASP	CG-OD2	7.70	1.43	1.25
1	A	362	TRP	CE3-CZ3	7.60	1.51	1.38
1	A	407	GLU	CD-OE1	7.58	1.33	1.25
1	B	36	GLY	N-CA	7.51	1.57	1.46
1	B	512	ASP	CB-CG	7.44	1.67	1.51
1	B	485	TRP	N-CA	7.05	1.60	1.46
1	B	726	GLU	CG-CD	6.99	1.62	1.51
1	A	343	GLU	CD-OE1	6.92	1.33	1.25
1	A	407	GLU	CG-CD	6.90	1.62	1.51
1	A	198	GLU	CD-OE1	6.84	1.33	1.25
1	B	161[A]	ARG	CZ-NH1	6.65	1.41	1.33
1	B	161[B]	ARG	CZ-NH1	6.65	1.41	1.33
1	A	725	GLN	CG-CD	6.58	1.66	1.51
1	A	294	GLY	N-CA	-6.55	1.36	1.46
1	B	208	TRP	CG-CD1	6.39	1.45	1.36
1	A	442	GLU	CD-OE1	-6.34	1.18	1.25
1	A	454	ASP	CB-CG	6.31	1.65	1.51
1	B	117	TYR	CE1-CZ	-6.23	1.30	1.38
1	A	584	GLU	CG-CD	5.94	1.60	1.51
1	B	410	GLU	CD-OE2	5.83	1.32	1.25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	675	TRP	CE3-CZ3	5.79	1.48	1.38
1	A	68	TYR	CE1-CZ	5.71	1.46	1.38
1	A	499	GLY	C-O	5.66	1.32	1.23
1	A	46	GLN	CG-CD	5.63	1.64	1.51
1	A	238	TYR	CE1-CZ	5.63	1.45	1.38
1	B	544	GLN	CD-NE2	5.50	1.46	1.32
1	A	365	LYS	N-CA	5.40	1.57	1.46
1	B	351	GLU	CD-OE1	5.38	1.31	1.25
1	A	159	TYR	CE2-CZ	-5.34	1.31	1.38
1	A	407	GLU	CD-OE2	5.27	1.31	1.25
1	A	183	LYS	C-O	5.14	1.33	1.23
1	B	675	TRP	CG-CD1	5.06	1.43	1.36
1	A	102	TYR	CD1-CE1	5.05	1.47	1.39

All (107) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	161[A]	ARG	NE-CZ-NH2	-13.65	113.47	120.30
1	B	161[B]	ARG	NE-CZ-NH2	-13.65	113.47	120.30
1	B	532	ARG	NE-CZ-NH1	11.81	126.20	120.30
1	B	512	ASP	CB-CG-OD2	11.54	128.69	118.30
1	A	532	ARG	NE-CZ-NH1	10.97	125.78	120.30
1	B	591	MET	CG-SD-CE	-9.95	84.28	100.20
1	B	512	ASP	CB-CG-OD1	-9.93	109.36	118.30
1	A	532	ARG	NE-CZ-NH2	-9.88	115.36	120.30
1	A	82	ARG	NE-CZ-NH2	-9.17	115.71	120.30
1	B	532	ARG	NE-CZ-NH2	-9.03	115.79	120.30
1	A	504	ARG	NE-CZ-NH1	8.99	124.80	120.30
1	A	649	ARG	NE-CZ-NH1	8.56	124.58	120.30
1	A	404	ARG	NE-CZ-NH2	-8.31	116.14	120.30
1	B	76	ASP	CB-CG-OD2	-8.30	110.83	118.30
1	A	82	ARG	NE-CZ-NH1	8.16	124.38	120.30
1	B	504	ARG	NE-CZ-NH2	-7.90	116.35	120.30
1	B	40	ARG	NE-CZ-NH1	7.82	124.21	120.30
1	B	268	ASP	CB-CG-OD1	7.60	125.14	118.30
1	A	591	MET	CG-SD-CE	-7.59	88.05	100.20
1	A	123	ARG	NE-CZ-NH2	-7.57	116.52	120.30
1	A	54	ARG	NE-CZ-NH1	7.39	124.00	120.30
1	A	504	ARG	NE-CZ-NH2	-7.29	116.66	120.30
1	B	293	ASP	N-CA-CB	7.22	123.60	110.60
1	B	713	ARG	NE-CZ-NH1	7.18	123.89	120.30
1	A	54	ARG	CG-CD-NE	-7.06	96.97	111.80

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	76	ASP	CB-CG-OD2	-7.06	111.95	118.30
1	A	161	ARG	NE-CZ-NH1	7.05	123.83	120.30
1	A	587	ASP	CB-CG-OD1	7.02	124.62	118.30
1	A	191	ARG	NE-CZ-NH2	-6.98	116.81	120.30
1	B	726	GLU	OE1-CD-OE2	-6.82	115.11	123.30
1	B	60	ASP	CB-CG-OD1	6.80	124.42	118.30
1	A	293	ASP	N-CA-CB	6.76	122.77	110.60
1	A	731	ASP	CB-CG-OD2	-6.74	112.23	118.30
1	B	649	ARG	NE-CZ-NH2	-6.64	116.98	120.30
1	A	578	ARG	NE-CZ-NH2	-6.53	117.04	120.30
1	A	431	ARG	NE-CZ-NH2	6.52	123.56	120.30
1	A	60	ASP	CB-CG-OD1	6.47	124.13	118.30
1	B	82	ARG	NE-CZ-NH2	-6.46	117.07	120.30
1	A	76	ASP	CB-CG-OD1	6.46	124.11	118.30
1	A	268	ASP	CB-CG-OD2	-6.45	112.50	118.30
1	A	454	ASP	CB-CG-OD2	6.44	124.09	118.30
1	B	719	TYR	CB-CG-CD2	-6.44	117.14	121.00
1	A	161	ARG	NE-CZ-NH2	-6.37	117.12	120.30
1	B	150	ARG	NE-CZ-NH1	6.32	123.46	120.30
1	A	40	ARG	NE-CZ-NH1	6.31	123.45	120.30
1	A	688	ARG	NE-CZ-NH1	6.31	123.45	120.30
1	A	591	MET	CA-CB-CG	6.29	123.99	113.30
1	A	268	ASP	CB-CG-OD1	6.27	123.94	118.30
1	A	343	GLU	CA-CB-CG	6.25	127.14	113.40
1	B	433	ARG	NE-CZ-NH1	6.24	123.42	120.30
1	B	688	ARG	NE-CZ-NH2	-6.24	117.18	120.30
1	B	540	ARG	NE-CZ-NH1	6.24	123.42	120.30
1	A	407	GLU	CG-CD-OE1	6.20	130.69	118.30
1	B	578	ARG	NE-CZ-NH2	-6.17	117.21	120.30
1	A	689	ASP	CB-CG-OD2	-6.06	112.85	118.30
1	A	723	ASP	CB-CG-OD2	6.05	123.75	118.30
1	B	150	ARG	NE-CZ-NH2	-6.05	117.28	120.30
1	A	433	ARG	NE-CZ-NH1	6.05	123.32	120.30
1	A	540	ARG	NE-CZ-NH1	6.03	123.32	120.30
1	B	393	ARG	NE-CZ-NH1	6.03	123.31	120.30
1	B	504	ARG	NE-CZ-NH1	6.01	123.30	120.30
1	A	404	ARG	NE-CZ-NH1	5.97	123.29	120.30
1	B	161[A]	ARG	NE-CZ-NH1	5.96	123.28	120.30
1	B	161[B]	ARG	NE-CZ-NH1	5.96	123.28	120.30
1	B	108	ARG	NE-CZ-NH2	-5.93	117.33	120.30
1	A	223	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	B	434	TYR	CB-CG-CD1	5.83	124.50	121.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	719	TYR	CB-CG-CD1	5.82	124.49	121.00
1	B	314	ARG	NE-CZ-NH2	-5.82	117.39	120.30
1	B	670	ASP	CB-CG-OD1	5.81	123.53	118.30
1	A	434	TYR	CB-CG-CD1	5.74	124.44	121.00
1	A	149	ARG	NE-CZ-NH1	5.73	123.17	120.30
1	B	541	GLY	N-CA-C	-5.73	98.78	113.10
1	A	167	ASP	CB-CG-OD1	5.72	123.45	118.30
1	A	433	ARG	NE-CZ-NH2	-5.71	117.45	120.30
1	A	293	ASP	CB-CG-OD2	-5.67	113.20	118.30
1	A	587	ASP	CB-CG-OD2	-5.67	113.20	118.30
1	A	263	ARG	NE-CZ-NH2	-5.65	117.47	120.30
1	B	293	ASP	CB-CG-OD2	-5.64	113.22	118.30
1	B	223	ARG	NE-CZ-NH1	5.55	123.08	120.30
1	A	656	ARG	NE-CZ-NH1	5.53	123.07	120.30
1	B	649	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	A	393	ARG	NE-CZ-NH1	5.49	123.04	120.30
1	B	46	GLN	CA-CB-CG	5.48	125.45	113.40
1	B	683	ASP	CB-CG-OD1	5.43	123.18	118.30
1	A	649	ARG	NE-CZ-NH2	-5.38	117.61	120.30
1	B	587	ASP	CB-CG-OD1	5.38	123.14	118.30
1	B	735	VAL	CG1-CB-CG2	-5.36	102.33	110.90
1	A	719	TYR	CB-CG-CD2	-5.32	117.81	121.00
1	B	198	GLU	CA-CB-CG	5.31	125.09	113.40
1	B	149	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	B	549	ASP	CB-CG-OD1	5.25	123.02	118.30
1	B	587	ASP	CB-CG-OD2	-5.24	113.58	118.30
1	B	54	ARG	NE-CZ-NH1	5.24	122.92	120.30
1	B	393	ARG	NE-CZ-NH2	-5.23	117.69	120.30
1	B	60	ASP	CB-CG-OD2	-5.19	113.63	118.30
1	B	670	ASP	CB-CG-OD2	-5.15	113.66	118.30
1	A	223	ARG	NE-CZ-NH2	-5.14	117.73	120.30
1	A	225	LEU	CB-CG-CD2	-5.14	102.26	111.00
1	B	433	ARG	NE-CZ-NH2	-5.12	117.74	120.30
1	A	222	ASP	CB-CG-OD2	-5.11	113.70	118.30
1	B	98	ASP	CB-CG-OD1	-5.10	113.71	118.30
1	A	382	ARG	NE-CZ-NH2	-5.10	117.75	120.30
1	A	46	GLN	CA-CB-CG	5.09	124.59	113.40
1	B	46	GLN	CB-CG-CD	5.04	124.72	111.60
1	A	132	ARG	NE-CZ-NH2	-5.04	117.78	120.30
1	A	605	LEU	CB-CG-CD1	-5.01	102.48	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5566	0	5359	19	0
1	B	5544	0	5343	22	0
2	A	43	0	30	0	0
2	B	43	0	30	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
5	A	2	0	0	0	0
5	B	2	0	0	1	0
6	A	5	0	0	1	0
6	B	5	0	0	0	0
7	A	24	0	42	7	0
7	B	8	0	14	2	0
8	A	786	0	0	9	0
8	B	764	0	0	11	0
All	All	12795	0	10818	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:808:MPD:H11	8:A:1330:HOH:O	1.71	0.90
1:B:647:GLN:HG2	8:B:1417:HOH:O	1.75	0.85
1:B:289:PRO:HD2	8:B:1562:HOH:O	1.77	0.85
1:B:512:ASP:OD1	8:B:901:HOH:O	1.94	0.85
1:B:76:ASP:OD2	8:B:902:HOH:O	1.96	0.82
1:B:519:GLU:OE1	8:B:903:HOH:O	1.97	0.82
1:A:119[B]:THR:HG21	8:A:1074:HOH:O	1.80	0.81
1:B:198:GLU:OE1	8:B:904:HOH:O	2.08	0.71
7:B:805:MPD:H52	7:B:805:MPD:HM2	1.81	0.63

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:803:OXY:O2	8:B:905:HOH:O	2.16	0.63
1:B:54:ARG:HB3	1:B:55:HIS:CD2	2.35	0.61
7:A:806:MPD:O4	7:A:806:MPD:H12	2.00	0.61
1:A:610:ARG:HG3	1:A:611:VAL:HG23	1.82	0.60
1:A:633[A]:VAL:CG2	1:A:719:TYR:CZ	2.86	0.58
1:B:540:ARG:HA	1:B:540:ARG:NE	2.18	0.58
1:A:255[B]:ARG:HG2	8:A:905:HOH:O	2.04	0.57
7:A:806:MPD:O4	7:A:806:MPD:C1	2.52	0.56
1:B:633[A]:VAL:CG2	1:B:719:TYR:CZ	2.90	0.55
1:A:593:VAL:HG13	8:A:1042:HOH:O	2.07	0.55
1:B:178:GLU:OE1	8:B:906:HOH:O	2.18	0.54
7:A:808:MPD:C1	8:A:1330:HOH:O	2.45	0.51
1:B:540:ARG:CZ	1:B:540:ARG:HA	2.40	0.51
1:A:455:HIS:CE1	1:A:544:GLN:HG3	2.46	0.51
1:A:400:LYS:NZ	8:A:916:HOH:O	2.44	0.50
1:A:119[A]:THR:CG2	1:A:265:ALA:HB2	2.42	0.50
1:B:284:THR:HG22	2:B:801:HEM:HAA1	1.94	0.50
1:B:255:ARG:HB2	8:B:1604:HOH:O	2.11	0.50
1:B:119[B]:THR:HG23	1:B:593:VAL:HG11	1.95	0.47
1:A:343:GLU:HG3	8:A:1508:HOH:O	2.14	0.47
1:B:119[A]:THR:CG2	1:B:265:ALA:HB2	2.45	0.46
1:A:158:LYS:HD2	7:A:808:MPD:H13	1.97	0.46
1:A:76:ASP:OD2	8:A:903:HOH:O	2.20	0.46
7:B:805:MPD:H52	7:B:805:MPD:CM	2.44	0.46
1:A:119[A]:THR:HG22	1:A:265:ALA:HB2	1.99	0.45
1:B:431:ARG:HD2	1:B:447:GLN:OE1	2.17	0.44
1:A:40:ARG:NH1	8:A:930:HOH:O	2.51	0.44
7:A:807:MPD:O4	7:A:807:MPD:HM1	2.18	0.44
7:A:808:MPD:H52	7:A:808:MPD:H11	2.00	0.44
1:A:119[B]:THR:HG23	1:A:593:VAL:HG11	1.99	0.44
1:B:512:ASP:CB	8:B:901:HOH:O	2.66	0.43
1:A:313:TYR:CZ	1:A:314:ARG:HD3	2.54	0.43
1:B:74:LYS:HD3	8:B:1443:HOH:O	2.19	0.42
1:B:264:MET:O	1:B:265:ALA:HB3	2.20	0.42
1:A:381:HIS:ND1	6:A:805:PO4:O4	2.42	0.41
1:A:510:GLN:HG2	1:A:513:TRP:CH2	2.56	0.41
1:A:88:MET:HB3	1:A:101:HIS:CE1	2.55	0.41
1:B:633[A]:VAL:HG22	1:B:719:TYR:CZ	2.55	0.41
1:B:633[B]:VAL:HG12	1:B:728:PHE:CD1	2.56	0.40
1:B:455:HIS:CE1	1:B:544:GLN:HG3	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	717/728 (98%)	708 (99%)	8 (1%)	1 (0%)	55	38
1	B	715/728 (98%)	707 (99%)	8 (1%)	0	100	100
All	All	1432/1456 (98%)	1415 (99%)	16 (1%)	1 (0%)	55	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	292	ASN

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	556/560 (99%)	545 (98%)	11 (2%)	60	47
1	B	554/560 (99%)	543 (98%)	11 (2%)	60	47
All	All	1110/1120 (99%)	1088 (98%)	22 (2%)	62	47

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	ARG
1	A	118	ARG
1	A	141	ASP
1	A	289	PRO
1	A	428	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	544	GLN
1	A	591	MET
1	A	608	LYS
1	A	725	GLN
1	A	726[A]	GLU
1	A	726[B]	GLU
1	B	40	ARG
1	B	54	ARG
1	B	74	LYS
1	B	118	ARG
1	B	141	ASP
1	B	198	GLU
1	B	289	PRO
1	B	540	ARG
1	B	544	GLN
1	B	608	LYS
1	B	690	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	GLN
1	A	647	GLN
1	B	406	HIS
1	B	544	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	TOX	A	111[A]	2	12,17,18	1.73	3 (25%)	11,23,25	2.00	4 (36%)
1	TOX	A	111[B]	-	12,17,18	1.73	3 (25%)	11,23,25	2.00	4 (36%)
1	TOX	B	111[A]	2	12,17,18	2.08	4 (33%)	11,23,25	3.07	6 (54%)
1	TOX	B	111[B]	-	12,17,18	2.08	4 (33%)	11,23,25	3.07	6 (54%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TOX	A	111[A]	2	-	0/3/8/10	0/2/2/2
1	TOX	A	111[B]	-	-	0/3/8/10	0/2/2/2
1	TOX	B	111[A]	2	-	0/3/8/10	0/2/2/2
1	TOX	B	111[B]	-	-	0/3/8/10	0/2/2/2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	111[A]	TOX	CD1-NE1	-4.25	1.35	1.39
1	A	111[B]	TOX	CD1-NE1	-4.25	1.35	1.39
1	B	111[B]	TOX	CD1-NE1	-3.04	1.36	1.39
1	B	111[A]	TOX	CD1-NE1	-3.04	1.36	1.39
1	B	111[B]	TOX	CA-C	-2.79	1.46	1.50
1	B	111[A]	TOX	CA-C	-2.79	1.46	1.50
1	A	111[A]	TOX	CZ3-CH2	2.24	1.43	1.38
1	A	111[B]	TOX	CZ3-CH2	2.24	1.43	1.38
1	A	111[A]	TOX	O-C	2.75	1.31	1.19
1	A	111[B]	TOX	O-C	2.75	1.31	1.19
1	B	111[B]	TOX	O-C	3.39	1.34	1.19
1	B	111[A]	TOX	O-C	3.39	1.34	1.19
1	B	111[B]	TOX	CH2-CZ2	4.10	1.46	1.36
1	B	111[A]	TOX	CH2-CZ2	4.10	1.46	1.36

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	111[B]	TOX	CB-CG-CD1	-5.99	120.57	127.97
1	B	111[A]	TOX	CB-CG-CD1	-5.99	120.57	127.97
1	A	111[A]	TOX	CZ3-CE3-CD2	-3.79	115.45	120.88
1	A	111[B]	TOX	CZ3-CE3-CD2	-3.79	115.45	120.88

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	111[B]	TOX	CZ3-CH2-CZ2	-3.58	115.34	120.45
1	B	111[A]	TOX	CZ3-CH2-CZ2	-3.58	115.34	120.45
1	A	111[A]	TOX	CB-CG-CD1	-2.54	124.83	127.97
1	A	111[B]	TOX	CB-CG-CD1	-2.54	124.83	127.97
1	B	111[B]	TOX	CZ3-CE3-CD2	-2.04	117.96	120.88
1	B	111[A]	TOX	CZ3-CE3-CD2	-2.04	117.96	120.88
1	A	111[A]	TOX	CB-CG-CD2	2.21	129.69	126.25
1	A	111[B]	TOX	CB-CG-CD2	2.21	129.69	126.25
1	B	111[B]	TOX	CB-CA-C	2.74	116.69	111.41
1	B	111[A]	TOX	CB-CA-C	2.74	116.69	111.41
1	A	111[A]	TOX	CB-CA-C	3.22	117.61	111.41
1	A	111[B]	TOX	CB-CA-C	3.22	117.61	111.41
1	B	111[B]	TOX	CH2-CZ3-CE3	3.47	125.40	120.45
1	B	111[A]	TOX	CH2-CZ3-CE3	3.47	125.40	120.45
1	B	111[B]	TOX	CB-CG-CD2	4.71	133.57	126.25
1	B	111[A]	TOX	CB-CG-CD2	4.71	133.57	126.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 3 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	HEM	A	801	1	28,50,50	1.67	6 (21%)	17,82,82	2.58	8 (47%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	OXY	A	804	-	1,1,1	0.07	0	0,0,0	0.00	-
6	PO4	A	805	-	4,4,4	0.80	0	6,6,6	1.11	0
7	MPD	A	806	-	7,7,7	0.94	0	9,10,10	0.76	0
7	MPD	A	807	-	7,7,7	0.62	0	9,10,10	1.17	0
7	MPD	A	808	-	7,7,7	0.89	0	9,10,10	1.51	2 (22%)
2	HEM	B	801	1	28,50,50	1.63	6 (21%)	17,82,82	3.13	11 (64%)
5	OXY	B	803	-	1,1,1	0.22	0	0,0,0	0.00	-
6	PO4	B	804	-	4,4,4	0.64	0	6,6,6	1.92	1 (16%)
7	MPD	B	805	-	7,7,7	0.95	0	9,10,10	2.08	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	801	1	-	0/6/54/54	0/0/8/8
5	OXY	A	804	-	-	0/0/0/0	0/0/0/0
6	PO4	A	805	-	-	0/0/0/0	0/0/0/0
7	MPD	A	806	-	-	0/5/5/5	0/0/0/0
7	MPD	A	807	-	-	0/5/5/5	0/0/0/0
7	MPD	A	808	-	-	0/5/5/5	0/0/0/0
2	HEM	B	801	1	-	0/6/54/54	0/0/8/8
5	OXY	B	803	-	-	0/0/0/0	0/0/0/0
6	PO4	B	804	-	-	0/0/0/0	0/0/0/0
7	MPD	B	805	-	-	0/5/5/5	0/0/0/0

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	HEM	C4D-ND	-4.14	1.31	1.36
2	B	801	HEM	C3B-C2B	-3.31	1.36	1.40
2	B	801	HEM	CMD-C2D	-3.30	1.44	1.51
2	B	801	HEM	C4B-CHC	-3.16	1.31	1.40
2	B	801	HEM	C1B-NB	-3.13	1.33	1.36
2	A	801	HEM	C1C-NC	-2.96	1.33	1.36
2	A	801	HEM	C3B-C2B	-2.64	1.36	1.40
2	A	801	HEM	C3C-C2C	-2.07	1.37	1.40
2	B	801	HEM	C4A-NA	2.16	1.40	1.36
2	A	801	HEM	C1D-CHD	2.31	1.46	1.40
2	B	801	HEM	CMA-C3A	3.13	1.58	1.51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	HEM	C4C-NC	3.93	1.41	1.36

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	HEM	CAA-CBA-CGA	-5.72	102.89	112.66
2	B	801	HEM	C4C-C3C-C2C	-5.57	103.01	106.90
2	B	801	HEM	CMA-C3A-C4A	-4.65	121.32	128.46
2	B	801	HEM	C1D-C2D-C3D	-3.97	104.23	107.00
2	B	801	HEM	CAA-CBA-CGA	-3.50	106.68	112.66
7	B	805	MPD	O2-C2-C3	-3.40	96.37	109.88
2	A	801	HEM	C3C-C4C-NC	-3.37	104.58	110.94
7	A	808	MPD	O4-C4-C3	-3.20	96.64	111.28
7	B	805	MPD	O2-C2-CM	-2.79	98.33	108.00
2	A	801	HEM	C1D-C2D-C3D	-2.63	105.17	107.00
2	B	801	HEM	CAD-C3D-C2D	-2.58	121.64	129.00
2	B	801	HEM	CAA-C2A-C3A	-2.38	122.20	129.00
7	A	808	MPD	O2-C2-CM	-2.06	100.88	108.00
2	A	801	HEM	CMC-C2C-C3C	2.05	128.70	124.89
2	A	801	HEM	CAD-CBD-CGD	2.25	116.51	112.66
2	B	801	HEM	CMD-C2D-C1D	2.69	132.60	128.46
2	B	801	HEM	CMB-C2B-C3B	2.99	130.44	124.89
2	A	801	HEM	C4C-C3C-C2C	3.01	109.00	106.90
2	B	801	HEM	CMA-C3A-C2A	3.06	130.71	124.94
2	B	801	HEM	CAD-CBD-CGD	3.27	118.25	112.66
2	A	801	HEM	CMB-C2B-C3B	3.42	131.23	124.89
6	B	804	PO4	O3-P-O2	4.06	122.81	107.90
7	B	805	MPD	CM-C2-C1	4.07	119.51	110.42
2	A	801	HEM	CBD-CAD-C3D	4.26	120.60	112.47
2	B	801	HEM	CBD-CAD-C3D	5.22	122.43	112.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	805	PO4	1	0
7	A	806	MPD	2	0
7	A	807	MPD	1	0
7	A	808	MPD	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	HEM	1	0
5	B	803	OXY	1	0
7	B	805	MPD	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	712/728 (97%)	-0.52	9 (1%) 77 74	12, 20, 38, 73	0
1	B	712/728 (97%)	-0.53	9 (1%) 77 74	13, 19, 36, 72	0
All	All	1424/1456 (97%)	-0.52	18 (1%) 77 74	12, 20, 37, 73	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	541	GLY	4.3
1	A	292	ASN	4.2
1	A	540	ARG	3.9
1	B	748	ALA	3.8
1	A	748	ALA	3.8
1	B	679	ALA	3.6
1	A	679	ALA	3.3
1	B	540	ARG	3.1
1	B	292	ASN	2.9
1	A	608	LYS	2.8
1	B	454	ASP	2.7
1	A	680	ALA	2.6
1	B	608	LYS	2.6
1	B	291	SER	2.4
1	B	610	ARG	2.3
1	A	610	ARG	2.3
1	A	291	SER	2.3
1	B	680	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	TOX	A	111[A]	16/17	0.97	0.12	-	13,15,21,27	1
1	TOX	A	111[B]	16/17	0.97	0.12	-	13,15,18,21	1
1	TOX	B	111[B]	16/17	0.97	0.13	-	12,15,18,22	1
1	TOX	B	111[A]	16/17	0.97	0.13	-	12,15,22,22	1

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
7	MPD	A	807	8/8	0.71	0.33	7.17	72,78,82,85	0
7	MPD	B	805	8/8	0.81	0.27	6.46	46,57,77,77	0
7	MPD	A	806	8/8	0.88	0.15	2.31	27,36,46,50	0
2	HEM	B	801	43/43	0.98	0.11	0.98	13,14,16,17	0
2	HEM	A	801	43/43	0.98	0.08	0.03	12,15,18,18	0
4	CL	A	803	1/1	0.98	0.05	-1.74	32,32,32,32	0
3	NA	B	802	1/1	1.00	0.03	-2.42	15,15,15,15	0
3	NA	A	802	1/1	1.00	0.04	-3.14	15,15,15,15	0
5	OXY	B	803	2/2	0.95	0.16	-	30,30,30,32	0
6	PO4	A	805	5/5	0.78	0.21	-	53,75,81,85	0
6	PO4	B	804	5/5	0.92	0.20	-	37,54,55,59	0
7	MPD	A	808	8/8	0.92	0.15	-	33,36,53,55	0
5	OXY	A	804	2/2	0.95	0.14	-	37,37,37,47	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.