W O R L D W I D E
wwPDB
PROTEIN DATA BANK

Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2017 – 01:16 PM EST

PDB ID	:	4KUE
Title	:	Crystal structure of 3-hydroxybutylryl-CoA dehydrogenase from Clostridium butyricum
Authors	:	Kim, E.J.; Kim, S.; Kim, K.J.
Deposited on	:	unknown
Resolution	:	2.00 Å(reported)

The following versions of software and data (see references ⓘ) were used in the production of this report:

|		
MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Percentile Ranks	Value
Rfree		0.240
Clashscore		5
Ramachandran outliers		0.2%
Sidechain outliers		4.8%
RSRZ outliers		2.9%

Worse .... *Better*

■ Percentile relative to all X-ray structures

▯ Percentile relative to X-ray structures of similar resolution

Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	5% ... 85% ... 10% · ·
1	B	288	4% ... 78% ... 18% · ·
1	C	288	2% ... 86% ... 10% · ·
1	D	288	83% ... 11% · ·

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 8855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-hydroxybutyryl-CoA dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	280	Total	C	N	O	S	0	0	0
			2114	1343	350	406	15			
1	B	280	Total	C	N	O	S	0	0	0
			2114	1343	350	406	15			
1	C	280	Total	C	N	O	S	0	0	0
			2114	1343	350	406	15			
1	D	280	Total	C	N	O	S	0	0	0
			2114	1343	350	406	15			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	281	SER	-	EXPRESSION TAG	UNP C4IEM5
A	282	LYS	-	EXPRESSION TAG	UNP C4IEM5
A	283	HIS	-	EXPRESSION TAG	UNP C4IEM5
A	284	HIS	-	EXPRESSION TAG	UNP C4IEM5
A	285	HIS	-	EXPRESSION TAG	UNP C4IEM5
A	286	HIS	-	EXPRESSION TAG	UNP C4IEM5
A	287	HIS	-	EXPRESSION TAG	UNP C4IEM5
A	288	HIS	-	EXPRESSION TAG	UNP C4IEM5
B	281	SER	-	EXPRESSION TAG	UNP C4IEM5
B	282	LYS	-	EXPRESSION TAG	UNP C4IEM5
B	283	HIS	-	EXPRESSION TAG	UNP C4IEM5
B	284	HIS	-	EXPRESSION TAG	UNP C4IEM5
B	285	HIS	-	EXPRESSION TAG	UNP C4IEM5
B	286	HIS	-	EXPRESSION TAG	UNP C4IEM5
B	287	HIS	-	EXPRESSION TAG	UNP C4IEM5
B	288	HIS	-	EXPRESSION TAG	UNP C4IEM5
C	281	SER	-	EXPRESSION TAG	UNP C4IEM5
C	282	LYS	-	EXPRESSION TAG	UNP C4IEM5
C	283	HIS	-	EXPRESSION TAG	UNP C4IEM5
C	284	HIS	-	EXPRESSION TAG	UNP C4IEM5
C	285	HIS	-	EXPRESSION TAG	UNP C4IEM5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	286	HIS	-	EXPRESSION TAG	UNP C4IEM5
C	287	HIS	-	EXPRESSION TAG	UNP C4IEM5
C	288	HIS	-	EXPRESSION TAG	UNP C4IEM5
D	281	SER	-	EXPRESSION TAG	UNP C4IEM5
D	282	LYS	-	EXPRESSION TAG	UNP C4IEM5
D	283	HIS	-	EXPRESSION TAG	UNP C4IEM5
D	284	HIS	-	EXPRESSION TAG	UNP C4IEM5
D	285	HIS	-	EXPRESSION TAG	UNP C4IEM5
D	286	HIS	-	EXPRESSION TAG	UNP C4IEM5
D	287	HIS	-	EXPRESSION TAG	UNP C4IEM5
D	288	HIS	-	EXPRESSION TAG	UNP C4IEM5

- Molecule 2 is water.

| Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	77	Total O 77 77	0	0
2	B	71	Total O 71 71	0	0
2	C	111	Total O 111 111	0	0
2	D	140	Total O 140 140	0	0

3 Residue-property plots ⓘ

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-hydroxybutyryl-CoA dehydrogenase

- Molecule 1: 3-hydroxybutyryl-CoA dehydrogenase

- Molecule 1: 3-hydroxybutyryl-CoA dehydrogenase

- Molecule 1: 3-hydroxybutyryl-CoA dehydrogenase

4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α, β, γ	144.11Å　144.11Å　205.00Å 90.00°　　90.00°　　120.00°	Depositor
Resolution (Å)	29.87　–　2.00 29.85　–　2.00	Depositor EDS
% Data completeness (in resolution range)	80.9 (29.87-2.00) 81.0 (29.85-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$<I/\sigma(I)>$ [1]	3.49 (at 2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.187　,　0.234 0.194　,　0.240	Depositor DCC
R_{free} test set	4393 reflections (5.33%)	DCC
Wilson B-factor (Å^2)	24.4	Xtriage
Anisotropy	0.251	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.39 , 52.0	EDS
L-test for twinning[2]	$<|L|> = 0.48, <L^2> = 0.31$	Xtriage
Estimated twinning fraction	0.029 for h,-h-k,-l	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	8855	wwPDB-VP
Average B, all atoms (Å^2)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.23 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.0225e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

---

[1]Intensities estimated from amplitudes.
[2]Theoretical values of $<|L|>$, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$|Z| > 5$	RMSZ	#$|Z| > 5$
1	A	0.76	0/2140	0.91	3/2882 (0.1%)
1	B	0.80	0/2140	0.91	4/2882 (0.1%)
1	C	0.93	0/2140	0.98	5/2882 (0.2%)
1	D	0.96	2/2140 (0.1%)	1.01	6/2882 (0.2%)
All	All	0.87	2/8560 (0.0%)	0.95	18/11528 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	90	GLU	CD-OE2	-5.98	1.19	1.25
1	D	210	GLU	CG-CD	5.09	1.59	1.51

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	271	ARG	NE-CZ-NH1	-8.20	116.20	120.30
1	D	255	ARG	NE-CZ-NH2	-8.11	116.24	120.30
1	D	271	ARG	NE-CZ-NH2	7.65	124.13	120.30
1	C	279	ASP	CB-CG-OD1	-7.21	111.81	118.30
1	D	30	ARG	NE-CZ-NH1	7.17	123.88	120.30
1	C	261	ARG	NE-CZ-NH1	6.85	123.72	120.30
1	B	255	ARG	NE-CZ-NH1	6.84	123.72	120.30
1	C	68	ARG	NE-CZ-NH1	6.71	123.65	120.30
1	C	68	ARG	NE-CZ-NH2	-6.69	116.96	120.30
1	D	255	ARG	NE-CZ-NH1	6.57	123.58	120.30
1	A	112	LEU	CA-CB-CG	5.87	128.81	115.30
1	B	182	ALA	C-N-CD	5.70	140.37	128.40
1	A	182	ALA	C-N-CD	5.51	139.96	128.40
1	A	232	LEU	CA-CB-CG	5.39	127.69	115.30
1	B	236	ASP	CB-CG-OD1	5.34	123.11	118.30
1	C	236	ASP	CB-CG-OD1	5.14	122.92	118.30
1	B	255	ARG	NE-CZ-NH2	-5.06	117.77	120.30
1	D	68	ARG	NE-CZ-NH2	-5.02	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2114	0	2193	20	0
1	B	2114	0	2193	34	0
1	C	2114	0	2193	16	0
1	D	2114	0	2193	28	0
2	A	77	0	0	1	0
2	B	71	0	0	3	0
2	C	111	0	0	0	0
2	D	140	0	0	3	0
All	All	8855	0	8772	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:194:MET:HE2	1:D:194:MET:HE2	1.50	0.90
1:B:275:LYS:HE2	2:B:366:HOH:O	1.88	0.73
1:A:121:ILE:HD12	2:A:363:HOH:O	1.90	0.71
1:D:58:THR:HG23	1:D:61:ASP:H	1.56	0.70
1:D:58:THR:HG22	1:D:61:ASP:CG	2.11	0.70
1:C:194:MET:HE2	1:D:194:MET:CE	2.24	0.68
1:D:130:ARG:HG2	1:D:133:LYS:HB2	1.79	0.65
1:A:177:VAL:HG22	1:B:211:ASP:HB3	1.78	0.64
1:D:112:LEU:HD22	1:D:130:ARG:HD2	1.83	0.60
1:B:237:VAL:O	1:B:241:ILE:HG12	2.02	0.59
1:C:194:MET:CE	1:D:194:MET:CE	2.80	0.59
1:C:113:ALA:HA	1:C:135:ILE:O	2.05	0.57
1:D:58:THR:CG2	1:D:61:ASP:H	2.18	0.56
1:D:52:VAL:HG21	1:D:59:GLU:HG2	1.87	0.56
1:B:191:LEU:HA	1:B:194:MET:HE3	1.88	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:THR:HG22	1:A:61:ASP:OD2	2.06	0.54
1:A:194:MET:HE2	1:B:194:MET:HE2	1.89	0.54
1:B:52:VAL:HG22	1:B:57:ILE:HG13	1.90	0.54
1:D:113:ALA:HA	1:D:135:ILE:O	2.08	0.54
1:D:89:ILE:HD11	1:D:94:ILE:HG21	1.91	0.52
1:A:177:VAL:HG22	1:B:211:ASP:CB	2.39	0.52
1:A:135:ILE:HD12	1:A:157:THR:HG21	1.92	0.51
1:B:252:THR:HG21	2:B:368:HOH:O	2.10	0.51
1:C:1:MET:HE2	1:C:168:MET:HB2	1.92	0.51
1:A:194:MET:HE2	1:B:194:MET:CE	2.40	0.51
1:C:1:MET:CE	1:C:168:MET:HB2	2.40	0.50
1:B:275:LYS:CD	1:B:279:ASP:OD1	2.59	0.50
1:C:194:MET:CE	1:D:194:MET:HE2	2.30	0.50
1:B:143:ALA:N	1:B:144:PRO:CD	2.75	0.50
1:C:33:LYS:HD3	1:C:35:GLU:H	1.77	0.50
1:A:91:ASN:OD1	1:A:93:LYS:HG2	2.12	0.49
1:A:138:HIS:HB3	1:A:150:GLU:HB2	1.94	0.49
1:C:199:THR:HG22	1:C:277:PHE:CZ	2.48	0.49
1:A:113:ALA:HA	1:A:135:ILE:O	2.12	0.49
1:B:76:LYS:HA	1:B:105:ILE:HD13	1.95	0.49
1:D:58:THR:HG21	2:D:394:HOH:O	2.12	0.48
1:A:64:GLU:HG2	1:A:68:ARG:NH2	2.28	0.48
1:D:119:LEU:HD12	1:D:119:LEU:N	2.29	0.48
1:D:89:ILE:CD1	1:D:94:ILE:HG21	2.45	0.47
1:B:84:VAL:HG11	1:B:102:LEU:HD13	1.96	0.47
1:B:7:LEU:N	1:B:7:LEU:HD12	2.30	0.47
1:B:80:ASP:HA	1:B:107:LYS:HE2	1.95	0.47
1:B:135:ILE:HG21	1:B:165:VAL:HG21	1.97	0.47
1:C:192:ILE:HB	1:C:193:PRO:HD3	1.97	0.47
1:C:217:LYS:O	1:C:221:ASN:HA	2.15	0.46
1:A:1:MET:HE2	1:A:168:MET:HB2	1.96	0.46
1:A:150:GLU:HA	1:A:177:VAL:HG12	1.98	0.46
1:A:129:LYS:HE3	1:A:129:LYS:HA	1.97	0.46
1:B:22:ALA:HA	1:B:65:ILE:HD12	1.98	0.46
1:B:58:THR:O	1:B:61:ASP:HB2	2.16	0.45
1:A:112:LEU:CD2	1:A:130:ARG:HD2	2.47	0.45
1:B:275:LYS:HD3	1:B:279:ASP:OD1	2.16	0.45
1:D:30:ARG:HD3	1:D:31:ASP:N	2.32	0.45
1:A:191:LEU:HA	1:A:194:MET:HE3	1.97	0.45
1:A:194:MET:CE	1:B:194:MET:CE	2.95	0.45
1:B:113:ALA:HA	1:B:135:ILE:O	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:39:ARG:O	1:B:42:ALA:HB3	2.17	0.45
1:B:49:SER:HB3	1:B:62:LYS:NZ	2.32	0.44
1:D:247:ASN:ND2	2:D:399:HOH:O	2.43	0.44
1:B:112:LEU:HD22	1:B:130:ARG:HD2	1.99	0.44
1:B:138:HIS:HB3	1:B:150:GLU:HB2	2.00	0.44
1:D:23:LYS:NZ	2:D:415:HOH:O	2.51	0.43
1:B:163:ASP:O	1:B:167:GLU:HG3	2.17	0.43
1:C:259:LEU:HD11	1:C:263:TYR:CE1	2.53	0.43
1:B:33:LYS:HG3	1:B:35:GLU:H	1.84	0.43
1:D:119:LEU:CD1	1:D:119:LEU:N	2.81	0.43
1:D:259:LEU:HD11	1:D:263:TYR:CE1	2.53	0.43
1:C:74:ASP:HB3	1:C:77:LEU:HD13	1.99	0.43
1:B:48:LEU:O	1:B:51:LEU:HB2	2.18	0.42
1:C:96:LYS:HG2	1:C:127:ALA:HB2	2.00	0.42
1:B:275:LYS:CE	2:B:366:HOH:O	2.58	0.42
1:D:194:MET:HB3	1:D:194:MET:HE3	1.88	0.42
1:A:58:THR:HG23	1:A:61:ASP:H	1.85	0.42
1:C:143:ALA:N	1:C:144:PRO:CD	2.83	0.42
1:D:64:GLU:HG2	1:D:68:ARG:NH2	2.35	0.42
1:B:217:LYS:O	1:B:221:ASN:HA	2.20	0.41
1:C:30:ARG:HG2	1:C:31:ASP:N	2.36	0.41
1:D:191:LEU:HA	1:D:194:MET:HE3	2.01	0.41
1:D:31:ASP:O	1:D:73:THR:HA	2.20	0.41
1:B:80:ASP:OD1	1:B:107:LYS:HE2	2.21	0.41
1:B:5:PHE:CZ	1:B:30:ARG:HB2	2.56	0.41
1:A:190:ILE:CG2	1:B:194:MET:HG3	2.51	0.41
1:B:58:THR:OG1	1:B:59:GLU:N	2.54	0.40
1:D:115:ASN:O	1:D:115:ASN:CG	2.60	0.40
1:D:201:ILE:HG23	1:D:206:VAL:HB	2.04	0.40
1:D:213:ASP:O	1:D:217:LYS:HG3	2.22	0.40
1:A:268:TRP:CD1	1:A:268:TRP:N	2.87	0.40
1:D:232:LEU:HA	1:D:271:ARG:HD2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	278/288 (96%)	270 (97%)	7 (2%)	1 (0%)	38	33
1	B	278/288 (96%)	267 (96%)	10 (4%)	1 (0%)	38	33
1	C	278/288 (96%)	268 (96%)	10 (4%)	0	100	100
1	D	278/288 (96%)	272 (98%)	6 (2%)	0	100	100
All	All	1112/1152 (96%)	1077 (97%)	33 (3%)	2 (0%)	51	48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	54	LYS
1	B	10	GLY

5.3.2　Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/234 (97%)	212 (94%)	14 (6%)	21	16
1	B	226/234 (97%)	216 (96%)	10 (4%)	33	28
1	C	226/234 (97%)	219 (97%)	7 (3%)	45	44
1	D	226/234 (97%)	214 (95%)	12 (5%)	26	21
All	All	904/936 (97%)	861 (95%)	43 (5%)	30	25

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	LYS
1	A	30	ARG
1	A	33	LYS
1	A	56	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	70	SER
1	A	76	LYS
1	A	89	ILE
1	A	93	LYS
1	A	108	PRO
1	A	112	LEU
1	A	129	LYS
1	A	177	VAL
1	A	232	LEU
1	A	269	LEU
1	B	11	THR
1	B	33	LYS
1	B	34	GLU
1	B	47	SER
1	B	70	SER
1	B	76	LYS
1	B	112	LEU
1	B	133	LYS
1	B	177	VAL
1	B	262	LYS
1	C	2	LYS
1	C	33	LYS
1	C	52	VAL
1	C	107	LYS
1	C	112	LEU
1	C	160	GLU
1	C	194	MET
1	D	12	MET
1	D	34	GLU
1	D	52	VAL
1	D	89	ILE
1	D	112	LEU
1	D	129	LYS
1	D	142	PRO
1	D	168	MET
1	D	177	VAL
1	D	194	MET
1	D	232	LEU
1	D	269	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	203	GLN
1	C	203	GLN
1	D	203	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	280/288 (97%)	-0.08	14 (5%)	30	30	26, 41, 66, 75	0
1	B	280/288 (97%)	-0.12	12 (4%)	36	36	25, 39, 71, 97	0
1	C	280/288 (97%)	-0.31	5 (1%)	69	68	19, 32, 58, 73	0
1	D	280/288 (97%)	-0.44	1 (0%)	92	92	18, 28, 46, 62	0
All	All	1120/1152 (97%)	-0.24	32 (2%)	52	52	18, 35, 62, 97	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	55	GLU	5.1
1	B	53	ALA	4.6
1	B	52	VAL	4.5
1	B	54	LYS	4.0
1	B	39	ARG	3.9
1	C	36	PHE	3.9
1	B	49	SER	3.9
1	A	32	ILE	3.6
1	B	56	LYS	3.5
1	A	36	PHE	3.3
1	A	52	VAL	3.3
1	A	55	GLU	3.0
1	C	93	LYS	2.9
1	B	50	LYS	2.6
1	B	51	LEU	2.6
1	A	39	ARG	2.6
1	B	57	ILE	2.6
1	A	53	ALA	2.4
1	B	46	LYS	2.4
1	C	56	LYS	2.4
1	A	54	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	6	VAL	2.3
1	A	279	ASP	2.2
1	D	39	ARG	2.2
1	B	32	ILE	2.2
1	C	34	GLU	2.2
1	A	50	LYS	2.2
1	A	85	VAL	2.2
1	C	32	ILE	2.1
1	A	56	LYS	2.1
1	A	35	GLU	2.0
1	A	93	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.