



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 05:58 am GMT

PDB ID : 3KX9
Title : Engineering a closed form of the Archaeoglobus fulgidus ferritin by site directed mutagenesis
Authors : Johnson, E.; Cascio, D.; Sawaya, M.R.
Deposited on : 2009-12-02
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





















Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	173	 77% 17% • 6%
1	B	173	 83% 11% 6%
1	C	173	 81% 13% • 6%
1	D	173	 79% 16% 5%
1	E	173	 82% 13% 6%
1	F	173	 83% 10% • 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	173	
1	H	173	
1	I	173	
1	J	173	
1	K	173	
1	L	173	
1	M	173	
1	N	173	
1	O	173	
1	P	173	
1	Q	173	
1	R	173	
1	S	173	
1	T	173	
1	U	173	
1	V	173	
1	W	173	
1	X	173	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	174	-	-	-	X
2	GOL	B	174	-	-	-	X
2	GOL	C	174	-	-	X	X
2	GOL	D	174	-	-	-	X
2	GOL	F	174	-	-	X	-
2	GOL	H	174	-	-	X	-
2	GOL	I	174	-	-	X	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	J	174	-	-	-	X
2	GOL	L	174	-	-	-	X
2	GOL	M	174	-	-	X	-
2	GOL	P	174	-	-	-	X
2	GOL	Q	174	-	-	-	X
2	GOL	R	174	-	-	-	X
2	GOL	S	174	-	-	-	X
2	GOL	T	174	-	-	-	X
2	GOL	U	174	-	-	-	X
2	GOL	X	174	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 33805 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ferritin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	163	Total	C	N	O	S	0	2	0
			1358	873	217	260	8			
1	B	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	C	163	Total	C	N	O	S	0	2	0
			1358	873	217	260	8			
1	D	164	Total	C	N	O	S	0	2	0
			1362	876	217	261	8			
1	E	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	F	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	G	163	Total	C	N	O	S	0	2	0
			1358	873	217	260	8			
1	H	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	I	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	J	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	K	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	L	163	Total	C	N	O	S	0	2	0
			1358	873	217	260	8			
1	M	163	Total	C	N	O	S	0	2	0
			1358	873	217	260	8			
1	N	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	O	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	P	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Q	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	R	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	S	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	T	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	U	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	V	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			
1	W	163	Total	C	N	O	S	0	2	0
			1358	873	217	260	8			
1	X	163	Total	C	N	O	S	0	2	0
			1355	872	216	259	8			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
A	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
B	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
B	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
C	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
C	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
D	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
D	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
E	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
E	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
F	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
F	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
G	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
G	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
H	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
H	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
I	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
I	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
J	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
J	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
K	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
K	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
L	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
L	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
M	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
M	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
N	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
N	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
O	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
O	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
P	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
P	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
Q	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
Q	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
R	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
R	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
S	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
S	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
T	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
T	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
U	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
U	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
V	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
V	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
W	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
W	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424
X	150	ALA	LYS	ENGINEERED MUTATION	UNP O29424
X	151	ALA	ARG	ENGINEERED MUTATION	UNP O29424

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	E	1	Total	C	O	0	0
			6	3	3		
2	F	1	Total	C	O	0	0
			6	3	3		
2	G	1	Total	C	O	0	0
			6	3	3		
2	H	1	Total	C	O	0	0
			6	3	3		
2	I	1	Total	C	O	0	0
			6	3	3		
2	J	1	Total	C	O	0	0
			6	3	3		
2	K	1	Total	C	O	0	0
			6	3	3		
2	L	1	Total	C	O	0	0
			6	3	3		
2	M	1	Total	C	O	0	0
			6	3	3		
2	N	1	Total	C	O	0	0
			6	3	3		
2	O	1	Total	C	O	0	0
			6	3	3		
2	P	1	Total	C	O	0	0
			6	3	3		
2	Q	1	Total	C	O	0	0
			6	3	3		
2	R	1	Total	C	O	0	0
			6	3	3		
2	S	1	Total	C	O	0	0
			6	3	3		
2	T	1	Total	C	O	0	0
			6	3	3		
2	U	1	Total	C	O	0	0
			6	3	3		
2	V	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	W	1	Total	C	O	0	0
			6	3	3		
2	X	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	48	Total	O	0	0
			48	48		
3	B	59	Total	O	0	0
			59	59		
3	C	44	Total	O	0	0
			44	44		
3	D	46	Total	O	0	0
			46	46		
3	E	51	Total	O	0	0
			51	51		
3	F	49	Total	O	0	0
			49	49		
3	G	41	Total	O	0	0
			41	41		
3	H	42	Total	O	0	0
			42	42		
3	I	43	Total	O	0	0
			43	43		
3	J	45	Total	O	0	0
			45	45		
3	K	50	Total	O	0	0
			50	50		
3	L	41	Total	O	0	0
			41	41		
3	M	52	Total	O	0	0
			52	52		
3	N	56	Total	O	0	0
			56	56		
3	O	40	Total	O	0	0
			40	40		
3	P	38	Total	O	0	0
			38	38		
3	Q	54	Total	O	0	0
			54	54		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	R	41	Total 41	O 41	0	0
3	S	34	Total 34	O 34	0	0
3	T	54	Total 54	O 54	0	0
3	U	53	Total 53	O 53	0	0
3	V	43	Total 43	O 43	0	0
3	W	44	Total 44	O 44	0	0
3	X	48	Total 48	O 48	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Ferritin




• Molecule 1: Ferritin




• Molecule 1: Ferritin





• Molecule 1: Ferritin




• Molecule 1: Ferritin




● Molecule 1: Ferritin

Chain F:  83% 10% 6%


● Molecule 1: Ferritin

Chain G:  83% 12% 6%


● Molecule 1: Ferritin

Chain H:  78% 16% 6%



● Molecule 1: Ferritin

Chain I:  75% 20% 6%

● Molecule 1: Ferritin

Chain J:  83% 9% 6%


● Molecule 1: Ferritin

Chain K:  79% 15% 6%

● Molecule 1: Ferritin


Chain L:  80% 13% 6%

- Molecule 1: Ferritin

Chain M: 




- Molecule 1: Ferritin

Chain N: 





- Molecule 1: Ferritin

Chain O: 





- Molecule 1: Ferritin

Chain P: 




- Molecule 1: Ferritin

Chain Q: 




- Molecule 1: Ferritin

Chain R: 




- Molecule 1: Ferritin


Chain S: 




● Molecule 1: Ferritin




● Molecule 1: Ferritin




● Molecule 1: Ferritin



● Molecule 1: Ferritin



● Molecule 1: Ferritin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	123.59Å 123.60Å 350.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.95 – 2.10 49.95 – 2.10	Depositor EDS
% Data completeness (in resolution range)	92.0 (49.95-2.10) 92.0 (49.95-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.65 (at 2.10Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
R, R_{free}	0.183 , 0.227 0.176 , 0.221	Depositor DCC
R_{free} test set	14477 reflections (5.06%)	DCC
Wilson B-factor (Å ²)	29.8	Xtriage
Anisotropy	0.162	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 53.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.034 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	33805	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1392	0.47	0/1879
1	B	0.38	0/1392	0.48	0/1879
1	C	0.38	0/1392	0.50	0/1879
1	D	0.38	0/1399	0.49	0/1889
1	E	0.39	0/1392	0.49	0/1879
1	F	0.40	0/1392	0.49	0/1879
1	G	0.40	0/1392	0.48	0/1879
1	H	0.37	0/1392	0.47	0/1879
1	I	0.39	0/1392	0.49	0/1879
1	J	0.38	0/1392	0.47	0/1879
1	K	0.37	0/1392	0.47	0/1879
1	L	0.40	0/1392	0.47	0/1879
1	M	0.37	0/1392	0.46	0/1879
1	N	0.38	0/1392	0.47	0/1879
1	O	0.38	0/1392	0.50	0/1879
1	P	0.38	0/1392	0.46	0/1879
1	Q	0.38	0/1392	0.50	0/1879
1	R	0.37	0/1392	0.48	0/1879
1	S	0.37	0/1392	0.48	0/1879
1	T	0.38	0/1392	0.49	0/1879
1	U	0.37	0/1392	0.48	0/1879
1	V	0.36	0/1392	0.48	0/1879
1	W	0.36	0/1392	0.49	0/1879
1	X	0.37	0/1392	0.47	0/1879
All	All	0.38	0/33415	0.48	0/45106

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1358	0	1304	30	0
1	B	1355	0	1305	11	0
1	C	1358	0	1304	20	0
1	D	1362	0	1312	19	0
1	E	1355	0	1305	13	0
1	F	1355	0	1305	21	0
1	G	1358	0	1304	13	0
1	H	1355	0	1305	19	0
1	I	1355	0	1305	24	0
1	J	1355	0	1305	12	0
1	K	1355	0	1305	19	0
1	L	1358	0	1304	15	0
1	M	1358	0	1304	20	0
1	N	1355	0	1305	19	0
1	O	1355	0	1305	19	0
1	P	1355	0	1305	24	0
1	Q	1355	0	1305	16	0
1	R	1355	0	1305	26	0
1	S	1355	0	1305	22	0
1	T	1355	0	1305	22	0
1	U	1355	0	1305	12	0
1	V	1355	0	1305	18	0
1	W	1358	0	1304	21	0
1	X	1355	0	1305	16	0
2	A	6	0	8	3	0
2	B	6	0	8	0	0
2	C	6	0	7	5	0
2	D	6	0	8	2	0
2	E	6	0	8	2	0
2	F	6	0	8	7	0
2	G	6	0	8	1	0
2	H	6	0	8	4	0
2	I	6	0	7	4	0
2	J	6	0	8	2	0
2	K	6	0	8	1	0
2	L	6	0	8	1	0
2	M	6	0	7	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	N	6	0	8	0	0
2	O	6	0	8	3	0
2	P	6	0	8	3	0
2	Q	6	0	7	1	0
2	R	6	0	8	2	0
2	S	6	0	8	0	0
2	T	6	0	8	0	0
2	U	6	0	8	1	0
2	V	6	0	8	1	0
2	W	6	0	8	2	0
2	X	6	0	7	3	0
3	A	48	0	0	1	0
3	B	59	0	0	0	0
3	C	44	0	0	1	0
3	D	46	0	0	3	0
3	E	51	0	0	4	0
3	F	49	0	0	1	0
3	G	41	0	0	1	0
3	H	42	0	0	3	0
3	I	43	0	0	1	0
3	J	45	0	0	2	0
3	K	50	0	0	2	0
3	L	41	0	0	2	0
3	M	52	0	0	2	0
3	N	56	0	0	7	0
3	O	40	0	0	0	0
3	P	38	0	0	4	0
3	Q	54	0	0	2	0
3	R	41	0	0	2	0
3	S	34	0	0	1	0
3	T	54	0	0	2	0
3	U	53	0	0	2	0
3	V	43	0	0	0	0
3	W	44	0	0	3	0
3	X	48	0	0	1	0
All	All	33805	0	31508	409	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 409 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:O:7:LYS:H	1:O:7:LYS:HD3	1.06	1.21
1:S:69:ARG:HG3	1:S:69:ARG:HH11	1.06	1.09
1:A:98:ASN:HD22	2:A:174:GOL:H11	1.20	1.03
1:O:7:LYS:HD3	1:O:7:LYS:N	1.74	1.02
1:C:7:LYS:HE3	1:C:7:LYS:H	1.26	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	163/173 (94%)	162 (99%)	1 (1%)	0	100	100
1	B	163/173 (94%)	162 (99%)	1 (1%)	0	100	100
1	C	163/173 (94%)	161 (99%)	2 (1%)	0	100	100
1	D	164/173 (95%)	163 (99%)	1 (1%)	0	100	100
1	E	163/173 (94%)	160 (98%)	2 (1%)	1 (1%)	28	24
1	F	163/173 (94%)	162 (99%)	1 (1%)	0	100	100
1	G	163/173 (94%)	162 (99%)	1 (1%)	0	100	100
1	H	163/173 (94%)	163 (100%)	0	0	100	100
1	I	163/173 (94%)	162 (99%)	1 (1%)	0	100	100
1	J	163/173 (94%)	162 (99%)	1 (1%)	0	100	100
1	K	163/173 (94%)	161 (99%)	2 (1%)	0	100	100
1	L	163/173 (94%)	161 (99%)	2 (1%)	0	100	100
1	M	163/173 (94%)	161 (99%)	2 (1%)	0	100	100
1	N	163/173 (94%)	161 (99%)	2 (1%)	0	100	100
1	O	163/173 (94%)	162 (99%)	1 (1%)	0	100	100
1	P	163/173 (94%)	160 (98%)	3 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Q	163/173 (94%)	162 (99%)	1 (1%)	0	100	100
1	R	163/173 (94%)	160 (98%)	2 (1%)	1 (1%)	28	24
1	S	163/173 (94%)	161 (99%)	2 (1%)	0	100	100
1	T	163/173 (94%)	162 (99%)	1 (1%)	0	100	100
1	U	163/173 (94%)	162 (99%)	1 (1%)	0	100	100
1	V	163/173 (94%)	161 (99%)	2 (1%)	0	100	100
1	W	163/173 (94%)	161 (99%)	2 (1%)	0	100	100
1	X	163/173 (94%)	163 (100%)	0	0	100	100
All	All	3913/4152 (94%)	3877 (99%)	34 (1%)	2 (0%)	55	57

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	R	4	ILE
1	E	149	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	144/150 (96%)	140 (97%)	4 (3%)	49	52
1	B	144/150 (96%)	139 (96%)	5 (4%)	41	42
1	C	144/150 (96%)	140 (97%)	4 (3%)	49	52
1	D	145/150 (97%)	141 (97%)	4 (3%)	49	52
1	E	144/150 (96%)	140 (97%)	4 (3%)	49	52
1	F	144/150 (96%)	138 (96%)	6 (4%)	34	33
1	G	144/150 (96%)	141 (98%)	3 (2%)	59	64
1	H	144/150 (96%)	137 (95%)	7 (5%)	29	26
1	I	144/150 (96%)	140 (97%)	4 (3%)	49	52
1	J	144/150 (96%)	138 (96%)	6 (4%)	34	33

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	K	144/150 (96%)	140 (97%)	4 (3%)	49	52
1	L	144/150 (96%)	140 (97%)	4 (3%)	49	52
1	M	144/150 (96%)	138 (96%)	6 (4%)	34	33
1	N	144/150 (96%)	140 (97%)	4 (3%)	49	52
1	O	144/150 (96%)	138 (96%)	6 (4%)	34	33
1	P	144/150 (96%)	140 (97%)	4 (3%)	49	52
1	Q	144/150 (96%)	139 (96%)	5 (4%)	41	42
1	R	144/150 (96%)	138 (96%)	6 (4%)	34	33
1	S	144/150 (96%)	136 (94%)	8 (6%)	25	21
1	T	144/150 (96%)	138 (96%)	6 (4%)	34	33
1	U	144/150 (96%)	138 (96%)	6 (4%)	34	33
1	V	144/150 (96%)	138 (96%)	6 (4%)	34	33
1	W	144/150 (96%)	140 (97%)	4 (3%)	49	52
1	X	144/150 (96%)	139 (96%)	5 (4%)	41	42
All	All	3457/3600 (96%)	3336 (96%)	121 (4%)	41	42

5 of 121 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	160	LEU
1	O	51	GLU
1	V	160	LEU
1	M	49	TRP
1	N	7	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 51 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	98	ASN
1	L	120	ASN
1	V	55	HIS
1	K	55	HIS
1	M	98	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	174	-	5,5,5	0.37	0	5,5,5	0.38	0
2	GOL	B	174	-	5,5,5	0.44	0	5,5,5	0.48	0
2	GOL	C	174	-	5,5,5	1.27	0	5,5,5	1.07	1 (20%)
2	GOL	D	174	-	5,5,5	0.67	0	5,5,5	0.90	0
2	GOL	E	174	-	5,5,5	0.35	0	5,5,5	0.25	0
2	GOL	F	174	-	5,5,5	1.18	1 (20%)	5,5,5	0.92	0
2	GOL	G	174	-	5,5,5	0.36	0	5,5,5	0.26	0
2	GOL	H	174	-	5,5,5	1.01	0	5,5,5	0.75	0
2	GOL	I	174	-	5,5,5	1.53	1 (20%)	5,5,5	0.67	0
2	GOL	J	174	-	5,5,5	0.79	0	5,5,5	0.81	0
2	GOL	K	174	-	5,5,5	0.42	0	5,5,5	0.30	0
2	GOL	L	174	-	5,5,5	1.25	1 (20%)	5,5,5	1.20	1 (20%)
2	GOL	M	174	-	5,5,5	1.29	1 (20%)	5,5,5	0.82	0
2	GOL	N	174	-	5,5,5	0.35	0	5,5,5	0.26	0
2	GOL	O	174	-	5,5,5	0.48	0	5,5,5	0.55	0
2	GOL	P	174	-	5,5,5	0.44	0	5,5,5	0.41	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	Q	174	-	5,5,5	1.30	1 (20%)	5,5,5	0.90	0
2	GOL	R	174	-	5,5,5	0.31	0	5,5,5	0.35	0
2	GOL	S	174	-	5,5,5	0.42	0	5,5,5	0.54	0
2	GOL	T	174	-	5,5,5	1.12	1 (20%)	5,5,5	1.05	0
2	GOL	U	174	-	5,5,5	1.14	1 (20%)	5,5,5	0.78	0
2	GOL	V	174	-	5,5,5	0.36	0	5,5,5	0.65	0
2	GOL	W	174	-	5,5,5	0.46	0	5,5,5	0.55	0
2	GOL	X	174	-	5,5,5	1.32	1 (20%)	5,5,5	0.93	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	174	-	-	0/4/4/4	0/0/0/0
2	GOL	B	174	-	-	0/4/4/4	0/0/0/0
2	GOL	C	174	-	-	0/4/4/4	0/0/0/0
2	GOL	D	174	-	-	0/4/4/4	0/0/0/0
2	GOL	E	174	-	-	0/4/4/4	0/0/0/0
2	GOL	F	174	-	-	0/4/4/4	0/0/0/0
2	GOL	G	174	-	-	0/4/4/4	0/0/0/0
2	GOL	H	174	-	-	0/4/4/4	0/0/0/0
2	GOL	I	174	-	-	0/4/4/4	0/0/0/0
2	GOL	J	174	-	-	0/4/4/4	0/0/0/0
2	GOL	K	174	-	-	0/4/4/4	0/0/0/0
2	GOL	L	174	-	-	0/4/4/4	0/0/0/0
2	GOL	M	174	-	-	0/4/4/4	0/0/0/0
2	GOL	N	174	-	-	0/4/4/4	0/0/0/0
2	GOL	O	174	-	-	0/4/4/4	0/0/0/0
2	GOL	P	174	-	-	0/4/4/4	0/0/0/0
2	GOL	Q	174	-	-	0/4/4/4	0/0/0/0
2	GOL	R	174	-	-	0/4/4/4	0/0/0/0
2	GOL	S	174	-	-	0/4/4/4	0/0/0/0
2	GOL	T	174	-	-	0/4/4/4	0/0/0/0
2	GOL	U	174	-	-	0/4/4/4	0/0/0/0
2	GOL	V	174	-	-	0/4/4/4	0/0/0/0
2	GOL	W	174	-	-	0/4/4/4	0/0/0/0
2	GOL	X	174	-	-	0/4/4/4	0/0/0/0

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	174	GOL	O2-C2	-2.97	1.34	1.43
2	X	174	GOL	O2-C2	-2.78	1.35	1.43
2	Q	174	GOL	O2-C2	-2.46	1.36	1.43
2	M	174	GOL	O2-C2	-2.44	1.36	1.43
2	T	174	GOL	O2-C2	-2.17	1.37	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	174	GOL	O3-C3-C2	-2.58	97.09	110.07
2	C	174	GOL	O1-C1-C2	-2.05	99.74	110.07

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

20 monomers are involved in 52 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	174	GOL	3	0
2	C	174	GOL	5	0
2	D	174	GOL	2	0
2	E	174	GOL	2	0
2	F	174	GOL	7	0
2	G	174	GOL	1	0
2	H	174	GOL	4	0
2	I	174	GOL	4	0
2	J	174	GOL	2	0
2	K	174	GOL	1	0
2	L	174	GOL	1	0
2	M	174	GOL	4	0
2	O	174	GOL	3	0
2	P	174	GOL	3	0
2	Q	174	GOL	1	0
2	R	174	GOL	2	0
2	U	174	GOL	1	0
2	V	174	GOL	1	0
2	W	174	GOL	2	0
2	X	174	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	163/173 (94%)	-0.29	0 100 100	21, 30, 48, 71	1 (0%)
1	B	163/173 (94%)	-0.24	0 100 100	21, 32, 51, 68	1 (0%)
1	C	163/173 (94%)	-0.21	0 100 100	22, 32, 51, 69	1 (0%)
1	D	164/173 (94%)	-0.24	0 100 100	21, 31, 51, 73	1 (0%)
1	E	163/173 (94%)	-0.21	0 100 100	22, 31, 50, 64	1 (0%)
1	F	163/173 (94%)	-0.26	0 100 100	22, 31, 48, 65	1 (0%)
1	G	163/173 (94%)	-0.23	1 (0%) 89 91	22, 32, 50, 66	1 (0%)
1	H	163/173 (94%)	-0.23	0 100 100	21, 32, 49, 70	1 (0%)
1	I	163/173 (94%)	-0.23	1 (0%) 89 91	22, 32, 51, 63	1 (0%)
1	J	163/173 (94%)	-0.20	1 (0%) 89 91	20, 31, 52, 80	1 (0%)
1	K	163/173 (94%)	-0.12	2 (1%) 79 82	21, 33, 54, 70	1 (0%)
1	L	163/173 (94%)	-0.24	0 100 100	21, 32, 48, 63	1 (0%)
1	M	163/173 (94%)	-0.27	0 100 100	22, 31, 52, 68	1 (0%)
1	N	163/173 (94%)	-0.22	0 100 100	21, 32, 52, 70	1 (0%)
1	O	163/173 (94%)	-0.24	0 100 100	21, 32, 51, 66	1 (0%)
1	P	163/173 (94%)	-0.18	0 100 100	21, 33, 53, 72	1 (0%)
1	Q	163/173 (94%)	-0.21	1 (0%) 89 91	22, 32, 53, 68	1 (0%)
1	R	163/173 (94%)	-0.20	1 (0%) 89 91	22, 32, 52, 66	1 (0%)
1	S	163/173 (94%)	-0.18	1 (0%) 89 91	23, 33, 53, 81	1 (0%)
1	T	163/173 (94%)	-0.20	1 (0%) 89 91	21, 32, 53, 69	1 (0%)
1	U	163/173 (94%)	-0.21	0 100 100	22, 33, 51, 69	1 (0%)
1	V	163/173 (94%)	-0.18	0 100 100	24, 34, 52, 68	1 (0%)
1	W	163/173 (94%)	-0.15	2 (1%) 79 82	21, 34, 52, 62	1 (0%)
1	X	163/173 (94%)	-0.33	0 100 100	22, 32, 48, 64	1 (0%)

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
All	All	3913/4152 (94%)	-0.22	11 (0%) 93 94	20, 32, 52, 81	24 (0%)

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	K	3	SER	4.1
1	R	3	SER	3.6
1	J	3	SER	3.3
1	W	151	ALA	2.5
1	Q	3	SER	2.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GOL	X	174	6/6	0.82	0.28	21.86	44,54,58,60	0
2	GOL	D	174	6/6	0.82	0.24	8.55	52,58,64,67	0
2	GOL	T	174	6/6	0.93	0.23	7.21	46,48,53,54	0
2	GOL	S	174	6/6	0.87	0.27	7.01	46,47,56,56	0
2	GOL	J	174	6/6	0.84	0.23	6.15	61,66,74,82	0
2	GOL	U	174	6/6	0.88	0.23	5.91	40,46,49,52	0
2	GOL	C	174	6/6	0.92	0.22	4.67	39,48,55,66	0
2	GOL	A	174	6/6	0.91	0.22	4.63	45,54,57,59	0
2	GOL	L	174	6/6	0.87	0.23	4.36	48,58,64,65	0
2	GOL	R	174	6/6	0.79	0.21	4.11	48,50,51,56	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GOL	Q	174	6/6	0.89	0.23	3.36	43,46,52,54	0
2	GOL	I	174	6/6	0.89	0.21	2.87	39,49,56,58	0
2	GOL	B	174	6/6	0.92	0.20	2.85	45,49,59,62	0
2	GOL	P	174	6/6	0.92	0.22	2.12	45,48,58,59	0
2	GOL	E	174	6/6	0.92	0.19	2.00	45,53,54,57	0
2	GOL	H	174	6/6	0.94	0.15	1.59	59,63,65,68	0
2	GOL	W	174	6/6	0.97	0.17	1.09	51,52,57,61	0
2	GOL	K	174	6/6	0.93	0.19	-	55,57,61,67	0
2	GOL	F	174	6/6	0.88	0.23	-	59,60,64,64	0
2	GOL	G	174	6/6	0.89	0.22	-	44,52,53,61	0
2	GOL	V	174	6/6	0.84	0.23	-	42,52,54,63	0
2	GOL	M	174	6/6	0.95	0.18	-	47,51,54,57	0
2	GOL	N	174	6/6	0.91	0.24	-	46,52,56,60	0
2	GOL	O	174	6/6	0.92	0.23	-	48,49,61,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.