



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 03:29 pm GMT

PDB ID : 4KYT  
Title : The structure of superinhibitory phospholamban bound to the calcium pump SERCA1a  
Authors : Hurley, T.D.; Akin, B.L.; Jones, L.R.  
Deposited on : 2013-05-29  
Resolution : 2.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3466 (2.88-2.80)
Clashscore	112137	3975 (2.88-2.80)
Ramachandran outliers	110173	3902 (2.88-2.80)
Sidechain outliers	110143	3905 (2.88-2.80)
RSRZ outliers	101464	3501 (2.88-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	994	<div> <div>4%</div> <div>81%</div> <div>16%</div> <div>••</div> </div>
2	B	52	<div> <div>17%</div> <div>52%</div> <div>44%</div> <div>•</div> </div>
2	C	52	<div> <div>4%</div> <div>25%</div> <div>71%</div> <div>•</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7908 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERCA1a.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	975	Total	C	N	O	S	0	0	0
			7520	4786	1262	1415	57			


- Molecule 2 is a protein called Cardiac phospholamban.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	29	Total	C	N	O	S	0	0	0
			223	148	37	34	4			
2	C	15	Total	C	N	O	S	0	0	0
			118	80	18	18	2			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	27	ALA	ASN	ENGINEERED MUTATION	UNP P61012
B	30	CYS	ASN	ENGINEERED MUTATION	UNP P61012
B	37	ALA	LEU	ENGINEERED MUTATION	UNP P61012
B	49	GLY	VAL	ENGINEERED MUTATION	UNP P61012
C	27	ALA	ASN	ENGINEERED MUTATION	UNP P61012
C	30	CYS	ASN	ENGINEERED MUTATION	UNP P61012
C	37	ALA	LEU	ENGINEERED MUTATION	UNP P61012
C	49	GLY	VAL	ENGINEERED MUTATION	UNP P61012

- Molecule 3 is SUGAR (MALTOSE) (three-letter code: MAL) (formula: C<sub>12</sub>H<sub>22</sub>O<sub>11</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			23	12	11		
3	A	1	Total	C	O	0	0
			23	12	11		


- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	K	0	0
			1	1		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: SERCA1a




#### • Molecule 2: Cardiac phospholamban



#### • Molecule 2: Cardiac phospholamban





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.59Å 93.09Å 316.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.83 48.86 – 2.83	Depositor EDS
% Data completeness (in resolution range)	96.7 (50.00-2.83) 96.7 (48.86-2.83)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.98 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.239 , 0.284 0.236 , 0.283	Depositor DCC
$R_{free}$ test set	2167 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	75.4	Xtriage
Anisotropy	0.342	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 54.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	7908	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	92.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K, MAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/7654	0.58	0/10376
2	B	0.35	0/225	0.43	0/303
2	C	0.45	0/119	0.42	0/160
All	All	0.45	0/7998	0.58	0/10839

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7520	0	7623	107	0
2	B	223	0	246	1	0
2	C	118	0	125	1	0
3	A	46	0	44	0	0
4	A	1	0	0	0	0
All	All	7908	0	8038	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:748:GLU:HA	1:A:817:MET:HE3	1.47	0.96
1:A:788:ILE:HG13	1:A:789:PRO:HD2	1.53	0.91
1:A:751:ARG:HD2	1:A:817:MET:HE1	1.66	0.78
1:A:419:LEU:HD11	1:A:479:MET:SD	2.31	0.71
1:A:895:GLU:H	1:A:896:PRO:HD2	1.56	0.70
1:A:748:GLU:HA	1:A:817:MET:CE	2.20	0.69
1:A:810:ASN:HB3	1:A:930:ASN:HD22	1.58	0.68
1:A:909:MET:HE2	1:A:937:ILE:HA	1.74	0.68
1:A:810:ASN:HB2	1:A:811:PRO:HD2	1.77	0.67
1:A:748:GLU:CA	1:A:817:MET:HE3	2.23	0.66
1:A:24:LEU:HD12	1:A:149:ASP:HB3	1.79	0.64
1:A:521:VAL:CG2	1:A:563:ALA:HB3	2.28	0.64
1:A:749:GLU:O	1:A:753:ILE:HG12	1.97	0.64
1:A:751:ARG:HD2	1:A:817:MET:CE	2.27	0.64
1:A:811:PRO:HG3	1:A:929:VAL:HG13	1.80	0.64
1:A:521:VAL:HG21	1:A:563:ALA:HB3	1.81	0.63
1:A:32:HIS:CB	1:A:146:VAL:HG11	2.30	0.62
1:A:32:HIS:HB3	1:A:146:VAL:HG11	1.83	0.60
1:A:177:GLN:HE22	1:A:189:LYS:NZ	2.00	0.59
1:A:22:THR:HG22	1:A:23:GLY:O	2.02	0.59
1:A:32:HIS:HB3	1:A:146:VAL:CG1	2.31	0.59
1:A:762:ARG:HH21	1:A:918:GLU:HA	1.68	0.59
1:A:322:GLY:HA3	1:A:753:ILE:HD12	1.85	0.58
1:A:524:ARG:HD2	1:A:588:GLU:O	2.03	0.58
1:A:791:GLN:NE2	1:A:958:LYS:HG2	2.19	0.58
1:A:895:GLU:N	1:A:896:PRO:HD2	2.18	0.58
1:A:322:GLY:HA3	1:A:753:ILE:CD1	2.33	0.58
1:A:419:LEU:CD1	1:A:479:MET:SD	2.92	0.57
1:A:987:ILE:O	1:A:987:ILE:HG22	2.05	0.57
1:A:874:MET:HG3	1:A:875:GLN:HG3	1.86	0.56
1:A:368:ILE:HD13	1:A:409:GLY:HA3	1.88	0.56
1:A:104:VAL:HG11	1:A:801:GLY:HA2	1.86	0.56
1:A:227:GLY:O	1:A:230:THR:HG22	2.06	0.56
1:A:273:LEU:O	1:A:276:ILE:HG23	2.06	0.55
1:A:49:LEU:HG	1:A:110:ARG:HH12	1.72	0.55
1:A:978:ILE:O	1:A:982:GLU:HG2	2.05	0.55
1:A:501:ALA:O	1:A:503:SER:N	2.39	0.55
1:A:352:LYS:HD3	1:A:625:THR:CG2	2.37	0.54
1:A:125:GLU:OE1	1:A:158:LYS:HE3	2.07	0.54
1:A:865:VAL:HG12	1:A:867:TYR:H	1.72	0.54
1:A:926:PRO:HD2	1:A:929:VAL:HG23	1.88	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:762:ARG:HG3	1:A:837:TYR:HE2	1.72	0.53
1:A:352:LYS:HD3	1:A:625:THR:HG21	1.91	0.53
1:A:650:ASP:O	1:A:672:ARG:NH1	2.42	0.53
1:A:898:THR:HG23	1:A:958:LYS:HB2	1.90	0.52
1:A:473:SER:HA	1:A:476:ARG:HB2	1.90	0.52
1:A:342:LEU:CD2	1:A:733:MET:HE1	2.40	0.52
1:A:513:PHE:HD1	1:A:566:THR:HG22	1.75	0.51
1:A:352:LYS:CB	1:A:625:THR:HG22	2.40	0.51
1:A:651:ARG:HH11	1:A:651:ARG:HG2	1.76	0.51
1:A:230:THR:HG23	1:A:233:GLY:H	1.76	0.50
1:A:501:ALA:C	1:A:503:SER:H	2.14	0.50
1:A:73:PHE:HB2	1:A:91:PRO:HG3	1.93	0.50
1:A:147:PRO:HA	1:A:223:VAL:HG12	1.93	0.49
1:A:762:ARG:HG3	1:A:837:TYR:CE2	2.47	0.49
1:A:24:LEU:CD1	1:A:149:ASP:HB3	2.42	0.49
1:A:177:GLN:HE22	1:A:189:LYS:HZ1	1.60	0.49
1:A:528:VAL:HG11	1:A:541:VAL:HG11	1.95	0.49
1:A:352:LYS:HB3	1:A:625:THR:HG22	1.95	0.49
1:A:795:VAL:HG21	1:A:904:LEU:HD23	1.95	0.48
1:A:5:HIS:HE1	1:A:196:ASP:O	1.95	0.48
1:A:753:ILE:O	1:A:757:MET:HG2	2.13	0.48
1:A:762:ARG:NH2	1:A:918:GLU:HA	2.27	0.48
1:A:810:ASN:HB3	1:A:930:ASN:ND2	2.25	0.48
1:A:969:MET:HA	1:A:972:LYS:HG2	1.95	0.48
1:A:721:GLY:O	1:A:728:LYS:NZ	2.47	0.48
1:A:932:TRP:CZ3	2:B:28:LEU:HG	2.49	0.47
1:A:512:MET:HE3	1:A:570:PRO:HB2	1.97	0.46
1:A:32:HIS:HB2	1:A:146:VAL:HG11	1.96	0.46
1:A:802:LEU:HB2	1:A:803:PRO:CD	2.45	0.46
1:A:680:GLU:HB2	1:A:683:HIS:CE1	2.51	0.46
1:A:840:ILE:HD13	1:A:980:LEU:HD23	1.98	0.46
1:A:909:MET:CE	1:A:937:ILE:HA	2.44	0.46
1:A:495:SER:HB3	1:A:514:VAL:HG22	1.96	0.46
1:A:986:PHE:C	1:A:988:ALA:H	2.19	0.46
1:A:735:LEU:HD22	1:A:742:THR:HB	1.98	0.45
1:A:895:GLU:N	1:A:896:PRO:CD	2.80	0.45
1:A:367:PHE:HD2	1:A:552:TRP:CH2	2.34	0.45
1:A:342:LEU:HD21	1:A:733:MET:HE1	1.97	0.45
1:A:895:GLU:H	1:A:896:PRO:CD	2.27	0.45
1:A:802:LEU:HB3	1:A:936:SER:HB2	1.99	0.45
1:A:39:ASN:HB3	1:A:142:ALA:O	2.17	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:722:SER:OG	1:A:738:ASP:OD2	2.31	0.44
1:A:762:ARG:NH1	1:A:914:ASN:O	2.48	0.44
1:A:802:LEU:HB2	1:A:803:PRO:HD3	2.00	0.44
1:A:533:THR:HG22	1:A:534:ARG:N	2.32	0.44
1:A:791:GLN:HE22	1:A:958:LYS:HG2	1.81	0.44
1:A:840:ILE:O	1:A:844:VAL:HG12	2.18	0.44
1:A:423:SER:OG	1:A:438:GLY:HA3	2.18	0.43
1:A:646:GLU:HG2	1:A:647:GLU:O	2.19	0.42
1:A:270:ALA:O	1:A:274:ILE:HG12	2.20	0.42
1:A:795:VAL:HG21	1:A:904:LEU:CD2	2.49	0.42
1:A:853:ALA:O	1:A:857:MET:HG2	2.19	0.42
1:A:512:MET:HB2	1:A:567:ARG:HB3	2.01	0.42
1:A:127:GLY:HA3	1:A:145:ILE:HD11	2.03	0.41
1:A:515:LYS:HD2	1:A:562:LEU:HD13	2.02	0.41
1:A:701:THR:HA	1:A:718:ILE:O	2.20	0.41
1:A:403:ARG:O	1:A:406:GLN:HG2	2.21	0.41
1:A:605:LYS:HG2	1:A:606:GLU:OE1	2.21	0.41
1:A:788:ILE:CG1	1:A:789:PRO:HD2	2.38	0.41
2:C:30:CYS:O	2:C:33:ILE:HG22	2.21	0.41
1:A:533:THR:HG22	1:A:534:ARG:H	1.87	0.40
1:A:848:THR:HG21	1:A:904:LEU:HD13	2.03	0.40
1:A:177:GLN:NE2	1:A:189:LYS:NZ	2.68	0.40
1:A:326:MET:CE	1:A:342:LEU:HD23	2.51	0.40
1:A:162:ASP:CG	1:A:230:THR:OG1	2.59	0.40
1:A:89:VAL:O	1:A:93:VAL:HG23	2.20	0.40
1:A:446:THR:HG23	1:A:472:ASN:ND2	2.37	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	965/994 (97%)	904 (94%)	53 (6%)	8 (1%)	22	53
2	B	27/52 (52%)	25 (93%)	2 (7%)	0	100	100
2	C	13/52 (25%)	13 (100%)	0	0	100	100
All	All	1005/1098 (92%)	942 (94%)	55 (6%)	8 (1%)	22	53

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	867	TYR
1	A	502	LYS
1	A	507	ALA
1	A	504	SER
1	A	802	LEU
1	A	987	ILE
1	A	503	SER
1	A	276	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	824/840 (98%)	787 (96%)	37 (4%)	32	64
2	B	25/46 (54%)	24 (96%)	1 (4%)	36	69
2	C	13/46 (28%)	13 (100%)	0	100	100
All	All	862/932 (92%)	824 (96%)	38 (4%)	33	65

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	LEU
1	A	77	TRP
1	A	106	VAL
1	A	108	GLN
1	A	125	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	146	VAL
1	A	237	ASP
1	A	276	ILE
1	A	309	GLU
1	A	316	THR
1	A	318	CYS
1	A	324	ARG
1	A	394	GLU
1	A	402	ILE
1	A	406	GLN
1	A	430	THR
1	A	535	VAL
1	A	560	ARG
1	A	573	ARG
1	A	606	GLU
1	A	656	ARG
1	A	680	GLU
1	A	685	SER
1	A	689	GLU
1	A	712	LYS
1	A	767	SER
1	A	774	CYS
1	A	778	THR
1	A	813	ASP
1	A	818	ASP
1	A	844	VAL
1	A	876	CYS
1	A	898	THR
1	A	929	VAL
1	A	951	ASP
1	A	962	LEU
1	A	991	TYR
2	B	46	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	HIS
1	A	177	GLN
1	A	278	HIS
1	A	380	ASN
1	A	406	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	768	ASN
1	A	868	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	MAL	A	1001	-	24,24,24	0.64	0	35,35,35	1.57	7 (20%)
3	MAL	A	1002	-	24,24,24	0.55	0	35,35,35	1.19	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAL	A	1001	-	-	0/8/48/48	0/2/2/2
3	MAL	A	1002	-	-	0/8/48/48	0/2/2/2

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1002	MAL	C3'-C4'-C5'	-2.24	106.12	110.88
3	A	1001	MAL	O5'-C5'-C4'	-2.21	105.23	109.75
3	A	1001	MAL	C2'-C3'-C4'	2.02	113.80	109.61
3	A	1001	MAL	O1-C4'-C3'	2.05	112.14	107.19
3	A	1001	MAL	O5-C5-C4	2.69	114.62	109.66
3	A	1001	MAL	O5'-C1'-C2'	3.34	115.58	110.04
3	A	1001	MAL	C1-O5-C5	3.45	120.21	113.72
3	A	1001	MAL	C1'-C2'-C3'	3.65	117.25	110.65
3	A	1002	MAL	C1'-O5'-C5'	3.82	120.28	113.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	975/994 (98%)	0.32	42 (4%) 36 27	44, 77, 153, 208	0
2	B	29/52 (55%)	1.23	9 (31%) 0 0	115, 128, 177, 199	0
2	C	15/52 (28%)	1.03	2 (13%) 4 2	111, 176, 205, 208	0
All	All	1019/1098 (92%)	0.36	53 (5%) 28 20	44, 80, 161, 208	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	881	PRO	7.0
1	A	877	THR	6.4
1	A	878	GLU	5.0
1	A	867	TYR	4.8
1	A	964	LEU	4.7
1	A	868	HIS	4.5
1	A	874	MET	4.5
2	C	27	ALA	4.4
1	A	960	LYS	4.2
1	A	959	LEU	3.9
1	A	883	PHE	3.8
1	A	890	ILE	3.8
1	A	506	ALA	3.6
2	B	23	GLN	3.5
1	A	88	PHE	3.4
1	A	882	HIS	3.2
1	A	875	GLN	3.2
1	A	955	MET	3.1
1	A	990	ASN	3.0
1	A	871	THR	2.9
2	B	43	LEU	2.9
1	A	277	GLY	2.9
1	A	245	ASP	2.8

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	24	ALA	2.8
2	B	40	ILE	2.8
1	A	865	VAL	2.8
2	C	28	LEU	2.8
1	A	951	ASP	2.8
2	B	28	LEU	2.7
1	A	235	ILE	2.6
1	A	107	TRP	2.6
1	A	981	ASP	2.6
1	A	869	GLN	2.6
1	A	52	LEU	2.5
1	A	502	LYS	2.5
1	A	573	ARG	2.4
1	A	60	LEU	2.4
1	A	505	ARG	2.4
1	A	895	GLU	2.4
2	B	22	GLN	2.3
2	B	25	ARG	2.3
1	A	876	CYS	2.3
2	B	47	ILE	2.2
1	A	507	ALA	2.2
1	A	880	HIS	2.2
1	A	251	GLN	2.2
1	A	894	PRO	2.2
1	A	198	ARG	2.2
1	A	50	TRP	2.1
1	A	240	ALA	2.1
1	A	241	ALA	2.1
2	B	46	CYS	2.0
1	A	932	TRP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	MAL	A	1001	23/23	0.95	0.20	-0.02	76,80,82,83	0
3	MAL	A	1002	23/23	0.93	0.18	-0.96	89,90,90,91	0
4	K	A	1003	1/1	0.96	0.13	-2.87	81,81,81,81	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.