



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 01:45 am GMT

PDB ID : 3L0F  
Title : High resolution structure of C-Phycocyanin from *Thermosynechococcus elongatus*  
Authors : Fromme, R.; Brune, D.; Fromme, P.  
Deposited on : 2009-12-09  
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1024 (1.38-1.34)
Clashscore	112137	1063 (1.38-1.34)
Ramachandran outliers	110173	1048 (1.38-1.34)
Sidechain outliers	110143	1048 (1.38-1.34)
RSRZ outliers	101464	1025 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	162	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 98%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>%</span> <span>90%</span> <span>10%</span> </div> </div>
2	B	172	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 2%, yellow 12%, green 86%);"></div> <div style="display: flex; justify-content: space-between; width: 86%; margin: 0 auto;"> <span>2%</span> <span>86%</span> <span>12%</span> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3219 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called C-phycocyanin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	3	3	0
			1239	780	207	245	7			

- Molecule 2 is a protein called C-phycocyanin beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	172	Total	C	N	O	S	0	1	0
			1278	792	229	249	8			

- Molecule 3 is PHYCOCYANOBILIN (three-letter code: CYC) (formula:  $C_{33}H_{40}N_4O_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			43	33	4	6		
3	B	1	Total	C	N	O	0	0
			43	33	4	6		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			43	33	4	6		



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	274	Total	O	0	0
			274	274		
4	B	299	Total	O	0	0
			299	299		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: C-phycocyanin alpha chain



- Molecule 2: C-phycocyanin beta chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	187.10Å 187.10Å 59.82Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.50 – 1.35 19.05 – 1.35	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.50-1.35) 99.8 (19.05-1.35)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.05 (at 1.35Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.133 , 0.175 0.136 , 0.174	Depositor DCC
$R_{free}$ test set	4367 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.5	Xtriage
Anisotropy	0.506	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 55.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	3219	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CYC, MEN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.30	8/1267 (0.6%)	1.11	6/1720 (0.3%)
2	B	1.23	6/1285 (0.5%)	1.12	8/1740 (0.5%)
All	All	1.27	14/2552 (0.5%)	1.11	14/3460 (0.4%)

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	68	GLN	CD-NE2	7.51	1.51	1.32
1	A	39	GLU	CG-CD	6.75	1.62	1.51
2	B	115	GLU	CG-CD	6.72	1.62	1.51
2	B	132	GLN	CG-CD	6.44	1.65	1.51
1	A	117	GLU	CG-CD	6.34	1.61	1.51
1	A	149	GLU	CD-OE2	5.88	1.32	1.25
2	B	62	GLU	CD-OE1	5.82	1.32	1.25
1	A	91	TYR	CE2-CZ	-5.78	1.31	1.38
1	A	156	TYR	CD2-CE2	-5.76	1.30	1.39
1	A	32	LYS	CD-CE	5.35	1.64	1.51
1	A	47	ASN	CG-OD1	-5.33	1.12	1.24
2	B	115	GLU	CD-OE1	5.27	1.31	1.25
2	B	68	GLN	CG-CD	5.20	1.63	1.51
1	A	42	ARG	CZ-NH1	5.08	1.39	1.33

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	30	ARG	NE-CZ-NH1	11.67	126.14	120.30
2	B	57	ARG	NE-CZ-NH1	8.38	124.49	120.30
1	A	30	ARG	NE-CZ-NH2	-7.47	116.57	120.30
1	A	42	ARG	NE-CZ-NH2	-7.21	116.69	120.30
2	B	166	ARG	NE-CZ-NH2	-6.64	116.98	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	78	ARG	NE-CZ-NH2	-6.09	117.25	120.30
2	B	15	ARG	NE-CZ-NH2	-5.80	117.40	120.30
1	A	90	TYR	CB-CG-CD1	5.64	124.38	121.00
2	B	77	ARG	NE-CZ-NH2	-5.62	117.49	120.30
2	B	114	ARG	NE-CZ-NH2	-5.45	117.58	120.30
1	A	97	TYR	CB-CG-CD1	5.31	124.19	121.00
2	B	19	LEU	CB-CG-CD1	5.31	120.02	111.00
2	B	84	ARG	NE-CZ-NH2	-5.21	117.70	120.30
1	A	13	ASP	CB-CG-OD2	5.20	122.98	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1239	0	1219	7	0
2	B	1278	0	1282	20	0
3	A	43	0	35	1	0
3	B	86	0	72	18	0
4	A	274	0	0	2	0
4	B	299	0	0	8	0
All	All	3219	0	2608	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:153:CYS:SG	3:B:555:CYC:HAC1	1.35	1.62
2:B:153:CYS:SG	3:B:555:CYC:CAC	2.20	1.29
2:B:153:CYS:SG	3:B:555:CYC:C3C	2.61	0.88
3:B:584:CYC:HMC1	3:B:584:CYC:HBC2	1.57	0.86
2:B:153:CYS:CB	3:B:555:CYC:HAC1	2.14	0.77

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57[B]:GLN:HE21	1:A:61:GLN:HE21	1.33	0.74
3:B:584:CYC:HMB1	3:B:584:CYC:HBB2	1.70	0.73
2:B:72:MEN:HE22	3:B:584:CYC:HBD2	1.72	0.71
1:A:57[B]:GLN:HG3	1:A:61:GLN:NE2	2.07	0.70
2:B:15:ARG:NH1	4:B:577:HOH:O	2.25	0.69
2:B:109:CYS:HA	3:B:584:CYC:HBB1	1.81	0.62
1:A:79:GLU:HG3	4:A:313:HOH:O	2.01	0.59
2:B:119:ALA:HB3	4:B:625:HOH:O	2.03	0.58
3:B:584:CYC:HC	3:B:584:CYC:HMD2	1.68	0.58
3:B:584:CYC:CMB	3:B:584:CYC:HBB2	2.36	0.55
2:B:153:CYS:SG	3:B:555:CYC:CBC	2.93	0.54
1:A:57[B]:GLN:NE2	1:A:61:GLN:HE21	2.04	0.54
2:B:27:LEU:HD23	2:B:30:LEU:HD12	1.89	0.54
2:B:109:CYS:HA	3:B:584:CYC:CBB	2.39	0.53
2:B:132:GLN:NE2	4:B:560:HOH:O	2.40	0.53
3:A:484:CYC:HMD2	3:A:484:CYC:HC	1.73	0.53
2:B:29:ASN:HB2	4:B:466:HOH:O	2.08	0.53
2:B:172:ALA:HB2	4:B:548:HOH:O	2.11	0.50
1:A:57[B]:GLN:CG	1:A:61:GLN:NE2	2.76	0.48
2:B:108:ARG:O	3:B:584:CYC:HBB1	2.13	0.48
3:B:555:CYC:NB	3:B:555:CYC:HMA1	2.30	0.47
2:B:36:LYS:HE2	3:B:555:CYC:HMD3	1.97	0.46
2:B:23:GLN:NE2	4:B:259:HOH:O	2.48	0.45
1:A:32:LYS:CE	4:A:499:HOH:O	2.65	0.44
2:B:62:GLU:HG3	4:B:289:HOH:O	2.17	0.44
2:B:128:ALA:O	2:B:132:GLN:HG3	2.18	0.43
3:B:555:CYC:NC	3:B:555:CYC:HMD2	2.34	0.43
3:B:555:CYC:HB	3:B:555:CYC:HMA1	1.84	0.42
3:B:584:CYC:HMC1	3:B:584:CYC:CBC	2.40	0.41
1:A:43:ALA:HB3	1:A:142:LEU:HD21	2.02	0.40
2:B:172:ALA:CB	4:B:548:HOH:O	2.69	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	163/162 (101%)	160 (98%)	3 (2%)	0	100	100
2	B	170/172 (99%)	167 (98%)	3 (2%)	0	100	100
All	All	333/334 (100%)	327 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/123 (102%)	126 (100%)	0	100	100
2	B	128/127 (101%)	128 (100%)	0	100	100
All	All	254/250 (102%)	254 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	MEN	B	72	2	8,8,9	1.39	1 (12%)	8,9,11	2.00	3 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MEN	B	72	2	-	0/6/8/10	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	72	MEN	CA-C	3.39	1.54	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	72	MEN	OD1-CG-CB	-3.33	116.39	121.42
2	B	72	MEN	O-C-CA	-3.14	116.35	125.02
2	B	72	MEN	CB-CA-C	-2.78	106.05	111.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	72	MEN	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	CYC	A	484	-	36,46,46	2.19	9 (25%)	44,67,67	3.34	17 (38%)
3	CYC	B	555	-	36,46,46	2.60	12 (33%)	44,67,67	4.06	28 (63%)
3	CYC	B	584	2	36,46,46	3.36	17 (47%)	44,67,67	3.20	21 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CYC	A	484	-	-	2/21/74/74	0/4/4/4
3	CYC	B	555	-	-	2/21/74/74	0/4/4/4
3	CYC	B	584	2	-	2/21/74/74	0/4/4/4

All (38) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	584	CYC	C2C-C1C	-8.08	1.44	1.52
3	B	555	CYC	C2C-C1C	-6.24	1.46	1.52
3	B	584	CYC	CMC-C2C	-4.43	1.43	1.53
3	B	555	CYC	CAC-C3C	-3.74	1.46	1.54
3	B	555	CYC	CMC-C2C	-3.64	1.45	1.53
3	B	584	CYC	C3C-C4C	-3.48	1.45	1.50
3	B	584	CYC	C1C-NC	-2.73	1.34	1.37
3	B	584	CYC	C2C-C3C	-2.57	1.47	1.54
3	A	484	CYC	C2C-C1C	-2.55	1.49	1.52
3	A	484	CYC	C2C-C3C	-2.36	1.47	1.54
3	A	484	CYC	CBB-CAB	-2.11	1.41	1.51
3	A	484	CYC	C1C-NC	-2.07	1.35	1.37
3	B	584	CYC	C1B-C2B	2.01	1.48	1.45

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	584	CYC	CHD-C4C	2.02	1.43	1.38
3	B	584	CYC	CBC-CAC	2.04	1.63	1.50
3	B	584	CYC	OB-C4B	2.22	1.27	1.23
3	B	584	CYC	CHB-C4A	2.28	1.45	1.40
3	B	555	CYC	CHD-C4C	2.29	1.44	1.38
3	A	484	CYC	CHB-C1B	2.44	1.43	1.37
3	A	484	CYC	C3D-C2D	2.47	1.45	1.37
3	B	584	CYC	C3D-C2D	2.54	1.45	1.37
3	B	555	CYC	CHB-C1B	2.96	1.44	1.37
3	B	555	CYC	OC-C1C	3.13	1.29	1.23
3	B	584	CYC	CAC-C3C	3.16	1.60	1.54
3	B	584	CYC	C2A-C3A	3.46	1.44	1.36
3	B	555	CYC	C3D-C2D	3.48	1.48	1.37
3	B	555	CYC	OB-C4B	3.63	1.30	1.23
3	B	555	CYC	CHB-C4A	3.83	1.49	1.40
3	B	555	CYC	C2A-C3A	4.03	1.45	1.36
3	B	584	CYC	OC-C1C	4.04	1.31	1.23
3	B	555	CYC	C3B-C2B	4.09	1.45	1.36
3	B	584	CYC	CHB-C1B	4.69	1.49	1.37
3	A	484	CYC	OB-C4B	5.03	1.33	1.23
3	A	484	CYC	C3B-C2B	5.73	1.49	1.36
3	B	584	CYC	C3B-C2B	6.02	1.49	1.36
3	A	484	CYC	CHA-C1A	6.71	1.41	1.35
3	B	555	CYC	CHA-C1A	7.37	1.41	1.35
3	B	584	CYC	CHA-C1A	12.46	1.46	1.35

All (66) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	555	CYC	C4B-C3B-C2B	-8.64	103.05	108.01
3	A	484	CYC	C4B-C3B-C2B	-6.65	104.19	108.01
3	A	484	CYC	OB-C4B-C3B	-6.05	121.36	128.11
3	B	584	CYC	C1B-C2B-C3B	-5.81	101.46	107.83
3	B	584	CYC	C4B-C3B-C2B	-5.53	104.83	108.01
3	A	484	CYC	C1B-NB-C4B	-5.46	103.44	110.70
3	B	555	CYC	C4A-C3A-C2A	-5.27	100.39	106.51
3	B	555	CYC	OB-C4B-C3B	-4.84	122.71	128.11
3	B	555	CYC	OC-C1C-NC	-4.77	119.18	124.87
3	A	484	CYC	C1B-C2B-C3B	-4.47	102.93	107.83
3	B	555	CYC	C1B-NB-C4B	-4.39	104.86	110.70
3	B	584	CYC	CBB-CAB-C3B	-4.13	100.69	112.41
3	A	484	CYC	OC-C1C-NC	-4.13	119.95	124.87

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	584	CYC	CAC-C3C-C2C	-4.11	103.91	114.24
3	B	584	CYC	OC-C1C-C2C	-3.90	123.10	126.25
3	B	555	CYC	CHB-C4A-NA	-3.67	117.92	124.97
3	B	555	CYC	CHA-C1A-NA	-3.42	123.61	128.79
3	B	584	CYC	CBD-CAD-C3D	-3.41	105.97	112.48
3	A	484	CYC	C4A-C3A-C2A	-3.07	102.95	106.51
3	B	584	CYC	C1B-NB-C4B	-3.03	106.66	110.70
3	B	584	CYC	C1A-C2A-C3A	-2.86	103.60	106.81
3	B	555	CYC	C1B-C2B-C3B	-2.68	104.89	107.83
3	B	584	CYC	CBC-CAC-C3C	-2.62	107.57	113.51
3	B	555	CYC	CHB-C1B-C2B	-2.53	121.77	126.83
3	B	584	CYC	CHA-C1A-C2A	-2.34	119.84	125.39
3	B	584	CYC	OB-C4B-NB	-2.33	119.43	125.11
3	A	484	CYC	CHB-C1B-C2B	-2.21	122.40	126.83
3	B	584	CYC	CAA-CBA-CGA	-2.09	109.09	112.66
3	B	555	CYC	CAA-C2A-C1A	2.25	129.03	125.00
3	B	584	CYC	C2A-C1A-NA	2.27	113.39	109.93
3	B	555	CYC	OC-C1C-C2C	2.31	128.12	126.25
3	B	555	CYC	C4D-CHA-C1A	2.34	131.61	128.77
3	A	484	CYC	CMB-C2B-C1B	2.40	127.30	124.17
3	B	555	CYC	CAA-CBA-CGA	2.41	116.78	112.66
3	B	555	CYC	CAD-CBD-CGD	2.50	116.93	112.66
3	B	555	CYC	CMC-C2C-C3C	2.56	124.18	113.77
3	B	555	CYC	C2A-C1A-NA	2.59	113.88	109.93
3	B	555	CYC	C3A-C4A-NA	2.61	116.23	110.53
3	A	484	CYC	CMC-C2C-C3C	2.64	124.49	113.77
3	B	555	CYC	CMD-C2D-C3D	2.65	129.94	124.94
3	A	484	CYC	CAC-C3C-C4C	2.77	119.79	112.67
3	B	555	CYC	CAC-C3C-C2C	2.98	121.73	114.24
3	A	484	CYC	CAC-C3C-C2C	3.14	122.13	114.24
3	B	555	CYC	C2B-C1B-NB	3.47	112.20	107.04
3	B	584	CYC	CMC-C2C-C3C	3.53	128.09	113.77
3	A	484	CYC	C3C-C4C-NC	3.65	111.40	107.97
3	A	484	CYC	CMA-C3A-C4A	3.73	130.86	125.04
3	B	584	CYC	C3C-C4C-NC	4.37	112.08	107.97
3	B	584	CYC	C2B-C1B-NB	4.42	113.63	107.04
3	B	584	CYC	CMA-C3A-C4A	4.80	132.52	125.04
3	B	555	CYC	C2C-C1C-NC	4.88	112.69	108.28
3	A	484	CYC	C2B-C1B-NB	5.27	114.89	107.04
3	A	484	CYC	C2C-C1C-NC	5.83	113.54	108.28
3	B	584	CYC	C2C-C1C-NC	5.85	113.57	108.28
3	B	584	CYC	CMC-C2C-C1C	5.86	124.74	112.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	555	CYC	CMC-C2C-C1C	6.38	125.83	112.43
3	B	555	CYC	CMA-C3A-C4A	7.01	135.96	125.04
3	B	555	CYC	C3C-C4C-NC	7.02	114.58	107.97
3	B	584	CYC	CAB-C3B-C4B	7.16	127.56	121.53
3	B	555	CYC	CAB-C3B-C4B	7.26	127.65	121.53
3	B	555	CYC	CBC-CAC-C3C	7.36	130.18	113.51
3	B	584	CYC	C3B-C4B-NB	7.60	113.18	106.75
3	B	555	CYC	CAC-C3C-C4C	7.82	132.74	112.67
3	A	484	CYC	C3B-C4B-NB	8.49	113.93	106.75
3	B	555	CYC	C3B-C4B-NB	9.75	114.99	106.75
3	A	484	CYC	CAB-C3B-C4B	10.39	130.28	121.53

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	584	CYC	C1B-CHB-C4A-C3A
3	A	484	CYC	C1B-CHB-C4A-C3A
3	B	584	CYC	C1B-CHB-C4A-NA
3	B	555	CYC	C1B-CHB-C4A-C3A
3	A	484	CYC	C1B-CHB-C4A-NA
3	B	555	CYC	C1B-CHB-C4A-NA

There are no ring outliers.

3 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	484	CYC	1	0
3	B	555	CYC	9	0
3	B	584	CYC	9	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	162/162 (100%)	-0.17	2 (1%) 79 83	12, 18, 28, 34	0
2	B	171/172 (99%)	0.04	4 (2%) 61 67	15, 21, 32, 38	0
All	All	333/334 (99%)	-0.06	6 (1%) 69 73	12, 19, 30, 38	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	111	ASN	3.0
2	B	1	MET	2.8
1	A	14	THR	2.6
1	A	70	GLN	2.6
2	B	172	ALA	2.6
2	B	108	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	MEN	B	72	9/10	0.96	0.07	-	17,18,22,29	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	CYC	B	584	43/43	0.89	0.12	1.50	15,25,38,43	0
3	CYC	A	484	43/43	0.98	0.08	0.87	11,13,17,19	0
3	CYC	B	555	43/43	0.95	0.07	-0.22	18,22,28,34	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.