



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 13, 2017 – 08:30 am GMT

PDB ID : 3L2O  
Title : Structure-Based Mechanism of Dimerization-Dependent Ubiquitination by the SCFFbx4 Ubiquitin Ligase  
Authors : Li, Y.; Hao, B.  
Deposited on : 2009-12-15  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	149	
2	B	312	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3425 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called S-phase kinase-associated protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	145	Total	C	N	O	S	0	0	0
			1142	723	186	227	6			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1002	ALA	PRO	ENGINEERED	UNP P63208
A	?	-	ASP	DELETION	UNP P63208
A	?	-	ASP	DELETION	UNP P63208
A	?	-	GLU	DELETION	UNP P63208
A	?	-	GLY	DELETION	UNP P63208
A	?	-	ASP	DELETION	UNP P63208
A	?	-	ASP	DELETION	UNP P63208
A	?	-	PRO	DELETION	UNP P63208
A	?	-	PRO	DELETION	UNP P63208
A	?	-	PRO	DELETION	UNP P63208
A	?	-	GLU	DELETION	UNP P63208
A	?	-	ASP	DELETION	UNP P63208
A	?	-	ASP	DELETION	UNP P63208
A	?	-	GLU	DELETION	UNP P63208
A	?	-	ASN	DELETION	UNP P63208
A	?	-	LYS	DELETION	UNP P63208
A	?	-	GLU	DELETION	UNP P63208
A	?	-	LYS	DELETION	UNP P63208
A	?	-	ARG	DELETION	UNP P63208
A	1078	GLY	-	INSERTION	UNP P63208
A	1079	GLY	-	INSERTION	UNP P63208
A	1080	SER	-	INSERTION	UNP P63208
A	1081	GLY	-	INSERTION	UNP P63208

- Molecule 2 is a protein called F-box only protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	274	Total	C	N	O	S	0	0	0
			2207	1423	364	406	14			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	THR	DELETION	UNP Q9UKT5
B	?	-	ARG	DELETION	UNP Q9UKT5
B	?	-	ARG	DELETION	UNP Q9UKT5
B	?	-	ALA	DELETION	UNP Q9UKT5
B	?	-	SER	DELETION	UNP Q9UKT5
B	?	-	LYS	DELETION	UNP Q9UKT5
B	?	-	SER	DELETION	UNP Q9UKT5
B	?	-	SER	DELETION	UNP Q9UKT5
B	?	-	ARG	DELETION	UNP Q9UKT5
B	?	-	PRO	DELETION	UNP Q9UKT5
B	?	-	MET	DELETION	UNP Q9UKT5
B	?	-	TYR	DELETION	UNP Q9UKT5
B	?	-	GLY	DELETION	UNP Q9UKT5
B	?	-	ALA	DELETION	UNP Q9UKT5
B	?	-	VAL	DELETION	UNP Q9UKT5
B	?	-	THR	DELETION	UNP Q9UKT5
B	?	-	SER	DELETION	UNP Q9UKT5
B	?	-	PHE	DELETION	UNP Q9UKT5
B	?	-	LEU	DELETION	UNP Q9UKT5
B	?	-	HIS	DELETION	UNP Q9UKT5
B	?	-	SER	DELETION	UNP Q9UKT5


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	20	Total	O	0	0
			20	20		
3	B	56	Total	O	0	0
			56	56		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: S-phase kinase-associated protein 1



#### • Molecule 2: F-box only protein 4



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.19Å 92.19Å 148.07Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.80 27.14 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (50.00-2.80) 99.7 (27.14-2.80)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.87 (at 2.80Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.249 , 0.295 0.267 , 0.311	Depositor DCC
$R_{free}$ test set	915 reflections (5.24%)	DCC
Wilson B-factor (Å <sup>2</sup> )	77.0	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 79.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.019 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3425	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.78	0/1161	1.05	7/1572 (0.4%)
2	B	0.81	0/2266	0.96	6/3087 (0.2%)
All	All	0.80	0/3427	0.99	13/4659 (0.3%)

There are no bond length outliers.

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1144	ASP	CB-CA-C	-10.26	89.88	110.40
1	A	1133	GLU	CB-CA-C	-9.82	90.75	110.40
2	B	106	ASP	CB-CA-C	-8.57	93.27	110.40
1	A	1136	ARG	NE-CZ-NH1	8.28	124.44	120.30
2	B	106	ASP	CB-CG-OD2	7.34	124.90	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1142	0	1139	85	0
2	B	2207	0	2148	122	0
3	A	20	0	0	1	0
3	B	56	0	0	1	0
All	All	3425	0	3287	201	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 201 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1136:ARG:HA	1:A:1141:ILE:HD11	1.26	1.13
1:A:1132:PRO:O	1:A:1134:GLU:N	1.84	1.10
2:B:198:MET:CE	2:B:221:VAL:HG21	1.81	1.09
1:A:1136:ARG:CA	1:A:1141:ILE:HD11	1.85	1.06
2:B:198:MET:HE1	2:B:221:VAL:HG21	1.05	1.05

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	143/149 (96%)	111 (78%)	17 (12%)	15 (10%)	0	1
2	B	268/312 (86%)	231 (86%)	24 (9%)	13 (5%)	2	8
All	All	411/461 (89%)	342 (83%)	41 (10%)	28 (7%)	1	3

5 of 28 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1012	GLU
1	A	1013	ILE
1	A	1037	ASP
1	A	1068	ASP
1	A	1080	SER



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	129/133 (97%)	117 (91%)	12 (9%)	10	30
2	B	247/280 (88%)	222 (90%)	25 (10%)	9	25
All	All	376/413 (91%)	339 (90%)	37 (10%)	9	27

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	130	GLU
2	B	173	ILE
2	B	314	THR
2	B	136	PHE
2	B	137	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	86	ASN
2	B	87	HIS
2	B	278	GLN
2	B	81	GLN
2	B	280	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	145/149 (97%)	0.32	16 (11%) <b>6</b> <b>3</b>	30, 43, 69, 72	0
2	B	274/312 (87%)	0.42	26 (9%) <b>9</b> <b>5</b>	33, 45, 54, 63	0
All	All	419/461 (90%)	0.38	42 (10%) <b>8</b> <b>4</b>	30, 45, 61, 72	0

The worst 5 of 42 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1078	GLY	6.5
2	B	273	TYR	6.0
2	B	172	ILE	5.6
2	B	55	ALA	5.4
1	A	1079	GLY	5.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.