



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 16, 2017 – 04:59 PM EDT

PDB ID : 5L25
Title : Crystal Structure of Arabidopsis thaliana Bor1
Authors : Thurtle-Schmidt, B.H.; Stroud, R.M.
Deposited on : unknown
Resolution : 4.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1156 (4.62-3.60)
Clashscore	112137	1003 (4.58-3.66)
Ramachandran outliers	110173	1012 (4.62-3.62)
Sidechain outliers	110143	1194 (4.62-3.60)
RSRZ outliers	101464	1168 (4.62-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	652	<div> <div style="width: 44%; background-color: green;"></div> <div style="width: 17%; background-color: yellow;"></div> <div style="width: 39%; background-color: grey;"></div> </div> <div>44% 17% . 39%</div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2873 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Boron transporter 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	399	Total	C	N	O	S	0	0	0
			2873	1897	461	495	20			


There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	646	ALA	-	expression tag	UNP Q8VYR7
A	647	ALA	-	expression tag	UNP Q8VYR7
A	648	ALA	-	expression tag	UNP Q8VYR7
A	649	LEU	-	expression tag	UNP Q8VYR7
A	650	VAL	-	expression tag	UNP Q8VYR7
A	651	PRO	-	expression tag	UNP Q8VYR7
A	652	ARG	-	expression tag	UNP Q8VYR7

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Boron transporter 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 ₂ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	184.56 Å 184.56 Å 89.83 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.07 – 4.11 20.07 – 4.11	Depositor EDS
% Data completeness (in resolution range)	94.8 (20.07-4.11) 95.8 (20.07-4.11)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 4.07 Å)	Xtriage
Refinement program	REFMAC 5.8.0151	Depositor
R, R_{free}	0.359 , 0.391 0.360 , 0.393	Depositor DCC
R_{free} test set	565 reflections (4.99%)	DCC
Wilson B-factor (Å ²)	223.8	Xtriage
Anisotropy	0.296	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.17 , 218.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	2873	wwPDB-VP
Average B, all atoms (Å ²)	258.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	1/2935 (0.0%)	0.68	5/4005 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	471	SER	CB-OG	5.49	1.49	1.42

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	194	PRO	N-CA-CB	8.28	113.23	103.30
1	A	539	PRO	N-CA-CB	6.45	111.05	103.30
1	A	254	PRO	N-CA-CB	5.88	110.36	103.30
1	A	182	PRO	N-CA-CB	5.75	110.20	103.30
1	A	482	PRO	N-CA-CB	5.05	109.36	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2873	0	2721	95	0
All	All	2873	0	2721	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 17.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:553:TYR:CE1	1:A:577:LEU:HD12	1.46	1.50
1:A:553:TYR:OH	1:A:574:ILE:N	1.62	1.32
1:A:553:TYR:CD1	1:A:577:LEU:CD1	2.35	1.09
1:A:553:TYR:OH	1:A:573:MET:C	1.91	1.09
1:A:553:TYR:CD1	1:A:577:LEU:HD12	1.88	1.09
1:A:553:TYR:CE1	1:A:577:LEU:CD1	2.36	1.08
1:A:553:TYR:CD1	1:A:577:LEU:HD13	2.08	0.89
1:A:546:PHE:CD1	1:A:549:PHE:HD1	1.90	0.89
1:A:553:TYR:OH	1:A:574:ILE:CA	2.20	0.89
1:A:546:PHE:CD1	1:A:549:PHE:CD1	2.64	0.86
1:A:553:TYR:HE1	1:A:577:LEU:HD12	1.05	0.85
1:A:169:GLN:O	1:A:172:ILE:HG12	1.78	0.83
1:A:480:ALA:O	1:A:484:LEU:CB	2.26	0.83
1:A:553:TYR:HD1	1:A:577:LEU:CD1	1.92	0.82
1:A:546:PHE:O	1:A:548:LEU:N	2.15	0.79
1:A:553:TYR:HH	1:A:574:ILE:N	1.81	0.78
1:A:553:TYR:HD1	1:A:577:LEU:HD13	1.46	0.78
1:A:93:LEU:HD23	1:A:94:GLY:N	2.01	0.75
1:A:54:PHE:HB3	1:A:100:VAL:HG21	1.69	0.75
1:A:34:PHE:HA	1:A:37:LEU:HD12	1.73	0.70
1:A:553:TYR:OH	1:A:574:ILE:HA	1.90	0.70
1:A:91:LEU:O	1:A:93:LEU:N	2.25	0.70
1:A:169:GLN:O	1:A:172:ILE:CG1	2.40	0.69
1:A:574:ILE:HG23	1:A:578:ILE:HD12	1.75	0.67
1:A:201:GLY:O	1:A:205:LEU:HB2	1.97	0.65
1:A:78:CYS:C	1:A:93:LEU:HD11	2.16	0.65
1:A:56:GLU:HA	1:A:59:GLU:HB2	1.78	0.65
1:A:291:ILE:O	1:A:294:ILE:HG22	1.97	0.64
1:A:546:PHE:O	1:A:547:THR:C	2.36	0.64
1:A:57:GLN:HE21	1:A:496:PHE:HE1	1.47	0.63
1:A:560:LEU:HB3	1:A:566:ALA:HB1	1.78	0.63
1:A:552:THR:O	1:A:556:ILE:HG13	1.98	0.63
1:A:577:LEU:O	1:A:581:ARG:HG3	1.98	0.62
1:A:214:THR:HG1	1:A:247:TRP:HE1	1.43	0.62
1:A:553:TYR:CE2	1:A:578:ILE:HD11	2.36	0.61
1:A:163:ILE:O	1:A:167:PHE:HB2	2.01	0.60
1:A:174:GLY:HA3	1:A:493:TRP:CD2	2.36	0.60
1:A:357:VAL:HG12	1:A:359:PRO:HD2	1.83	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:477:CYS:HA	1:A:480:ALA:HB3	1.84	0.60
1:A:58:LEU:HD22	1:A:66:LEU:HD23	1.86	0.58
1:A:79:GLY:HA2	1:A:93:LEU:HD13	1.86	0.57
1:A:158:LEU:HD23	1:A:553:TYR:HD2	1.70	0.57
1:A:241:PRO:O	1:A:244:VAL:HB	2.04	0.57
1:A:200:ASN:HA	1:A:203:PHE:HB2	1.87	0.56
1:A:553:TYR:HE1	1:A:577:LEU:CD1	1.96	0.55
1:A:546:PHE:CZ	1:A:549:PHE:HB2	2.42	0.55
1:A:546:PHE:CE1	1:A:549:PHE:HB2	2.42	0.55
1:A:78:CYS:CB	1:A:93:LEU:HD11	2.39	0.52
1:A:137:PHE:CD2	1:A:337:LEU:HD11	2.44	0.52
1:A:337:LEU:O	1:A:339:LEU:N	2.43	0.52
1:A:546:PHE:C	1:A:548:LEU:N	2.60	0.52
1:A:93:LEU:HD23	1:A:93:LEU:C	2.31	0.51
1:A:569:MET:O	1:A:573:MET:HG2	2.10	0.50
1:A:79:GLY:N	1:A:93:LEU:HD11	2.28	0.49
1:A:79:GLY:CA	1:A:93:LEU:HD13	2.43	0.49
1:A:78:CYS:HB2	1:A:93:LEU:HD11	1.93	0.48
1:A:195:SER:HA	1:A:199:ALA:HB3	1.96	0.48
1:A:581:ARG:O	1:A:585:LEU:HG	2.13	0.48
1:A:174:GLY:HA3	1:A:493:TRP:CE3	2.49	0.48
1:A:38:ALA:HB3	1:A:39:PRO:HD3	1.96	0.48
1:A:123:ALA:O	1:A:126:GLY:N	2.48	0.47
1:A:504:LEU:HB2	1:A:505:PRO:HD3	1.97	0.47
1:A:546:PHE:CE1	1:A:549:PHE:CD1	3.03	0.47
1:A:221:ALA:HB3	1:A:236:ALA:HB2	1.97	0.46
1:A:172:ILE:HG13	1:A:173:LYS:N	2.30	0.46
1:A:169:GLN:HA	1:A:172:ILE:HG12	1.98	0.46
1:A:510:TRP:O	1:A:513:ILE:HG22	2.16	0.46
1:A:492:LEU:O	1:A:496:PHE:HD2	1.99	0.45
1:A:546:PHE:O	1:A:549:PHE:N	2.34	0.45
1:A:55:GLY:HA2	1:A:71:THR:OG1	2.17	0.45
1:A:211:LEU:HD21	1:A:495:TYR:CD1	2.52	0.44
1:A:492:LEU:HA	1:A:495:TYR:HD2	1.81	0.44
1:A:491:VAL:HG12	1:A:495:TYR:CE2	2.52	0.44
1:A:492:LEU:O	1:A:495:TYR:HB2	2.17	0.44
1:A:557:CYS:O	1:A:561:THR:HG23	2.18	0.43
1:A:79:GLY:N	1:A:93:LEU:CD1	2.81	0.43
1:A:308:TYR:CE1	1:A:355:ASN:HB3	2.54	0.43
1:A:58:LEU:HD21	1:A:100:VAL:HG13	2.01	0.43
1:A:233:SER:HA	1:A:236:ALA:HB3	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:304:ILE:HG23	1:A:308:TYR:CD2	2.54	0.43
1:A:307:LEU:HD12	1:A:571:PRO:HG3	2.01	0.43
1:A:495:TYR:O	1:A:496:PHE:C	2.57	0.43
1:A:50:PRO:HA	1:A:53:SER:HB3	2.01	0.43
1:A:139:LEU:HB2	1:A:309:TYR:OH	2.19	0.42
1:A:304:ILE:HG23	1:A:308:TYR:CE2	2.54	0.42
1:A:492:LEU:HD23	1:A:492:LEU:HA	1.86	0.42
1:A:127:TRP:HB3	1:A:294:ILE:HD11	2.02	0.42
1:A:514:LEU:HA	1:A:514:LEU:HD23	1.88	0.42
1:A:495:TYR:O	1:A:498:PHE:N	2.53	0.41
1:A:466:SER:HA	1:A:469:LEU:HD12	2.01	0.41
1:A:206:VAL:HA	1:A:209:PHE:CZ	2.56	0.41
1:A:300:PRO:O	1:A:304:ILE:HB	2.21	0.41
1:A:209:PHE:CD1	1:A:210:GLY:N	2.89	0.40
1:A:339:LEU:O	1:A:343:THR:HG23	2.21	0.40
1:A:224:TRP:O	1:A:232:ARG:NH1	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	383/652 (59%)	334 (87%)	40 (10%)	9 (2%)	<div><div>7</div>45</div>

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	92	ILE
1	A	203	PHE
1	A	338	LEU
1	A	535	VAL
1	A	82	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	117	GLY
1	A	369	ALA
1	A	193	LEU
1	A	194	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	265/553 (48%)	260 (98%)	5 (2%)	62 83

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	SER
1	A	127	TRP
1	A	167	PHE
1	A	562	TRP
1	A	583	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	399/652 (61%)	-0.78	1 (0%) 93 90	129, 249, 382, 500	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	142	CYS	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.