



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 7, 2017 – 11:52 AM EDT

PDB ID : 5L2L  
Title : Nab2 Zn fingers 5-7 bound to A11G RNA  
Authors : Stewart, M.; Aibara, S.  
Deposited on : 2016-08-02  
Resolution : 1.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20029077
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20029077


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1088 (1.56-1.56)
Clashscore	112137	1132 (1.56-1.56)
Ramachandran outliers	110173	1110 (1.56-1.56)
Sidechain outliers	110143	1108 (1.56-1.56)
RSRZ outliers	101464	1089 (1.56-1.56)
RNA backbone	2435	1043 (2.60-0.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	77	<div> <div>4%</div> <div>94%</div> <div>6%</div> </div>
1	B	77	<div> <div>5%</div> <div>95%</div> <div>• •</div> </div>
1	E	77	<div> <div>5%</div> <div>92%</div> <div>• 6%</div> </div>
1	F	77	<div> <div>3%</div> <div>97%</div> <div>•</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	C	12	
2	D	12	
2	G	12	
2	H	12	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ZN	A	503	-	-	-	X
3	ZN	B	502	-	-	-	X
3	ZN	E	501	-	-	-	X
3	ZN	F	503	-	-	-	X
3	ZN	F	504	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6787 atoms, of which 2756 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nab2p.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	72	Total	C	H	N	O	S	0	0	0
			1142	357	560	117	98	10			
1	B	74	Total	C	H	N	O	S	0	0	0
			1163	364	570	119	100	10			
1	E	72	Total	C	H	N	O	S	0	0	0
			1145	359	562	117	97	10			
1	F	75	Total	C	H	N	O	S	0	0	0
			1178	369	576	120	103	10			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	407	GLY	PRO	conflict	UNP A0A0C6D5P3
A	408	SER	VAL	conflict	UNP A0A0C6D5P3
B	407	GLY	PRO	conflict	UNP A0A0C6D5P3
B	408	SER	VAL	conflict	UNP A0A0C6D5P3
E	407	GLY	PRO	conflict	UNP A0A0C6D5P3
E	408	SER	VAL	conflict	UNP A0A0C6D5P3
F	407	GLY	PRO	conflict	UNP A0A0C6D5P3
F	408	SER	VAL	conflict	UNP A0A0C6D5P3

- Molecule 2 is a RNA chain called RNA (5'-R(\*AP\*AP\*AP\*AP\*AP\*AP\*AP\*AP\*AP\*G)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	10	Total	C	H	N	O	P	0	0	0
			332	100	111	50	61	10			
2	D	11	Total	C	H	N	O	P	0	1	0
			398	120	133	60	73	12			
2	G	10	Total	C	H	N	O	P	0	1	0
			365	110	122	55	67	11			
2	H	10	Total	C	H	N	O	P	0	1	0
			365	110	122	55	67	11			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total 3	Zn 3	0	0
3	A	4	Total 4	Zn 4	0	0
3	F	4	Total 4	Zn 4	0	0
3	E	3	Total 3	Zn 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	119	Total 119	O 119	0	0
4	B	133	Total 133	O 133	0	0
4	C	54	Total 54	O 54	0	0
4	D	49	Total 49	O 49	0	0
4	E	122	Total 122	O 122	0	0
4	F	130	Total 130	O 130	0	0
4	G	46	Total 46	O 46	0	0
4	H	32	Total 32	O 32	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nab2p




- Molecule 1: Nab2p




- Molecule 1: Nab2p




- Molecule 1: Nab2p




- Molecule 2: RNA (5'-R(\*AP\*AP\*AP\*AP\*AP\*AP\*AP\*AP\*G)-3')



- Molecule 2: RNA (5'-R(\*AP\*AP\*AP\*AP\*AP\*AP\*AP\*AP\*G)-3')



- Molecule 2: RNA (5'-R(\*AP\*AP\*AP\*AP\*AP\*AP\*AP\*AP\*G)-3')



- Molecule 2: RNA (5'-R(\*AP\*AP\*AP\*AP\*AP\*AP\*AP\*AP\*G)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.67Å 85.03Å 64.96Å 90.00° 99.07° 90.00°	Depositor
Resolution (Å)	46.09 – 1.55 46.09 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.4 (46.09-1.55) 99.4 (46.09-1.55)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.11 (at 1.55Å)	Xtriage
Refinement program	PHENIX (1.10.1 _2155: ???)	Depositor
R, $R_{free}$	0.131 , 0.152 0.126 , 0.148	Depositor DCC
$R_{free}$ test set	3567 reflections (4.95%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.3	Xtriage
Anisotropy	0.324	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 52.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	6787	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.46	0/595	0.57	0/795
1	B	0.45	0/607	0.64	1/812 (0.1%)
1	E	0.48	0/597	0.62	0/799
1	F	0.42	0/616	0.60	0/824
2	C	0.73	0/250	1.07	0/388
2	D	0.64	0/300	0.94	0/466
2	G	0.59	0/275	0.92	0/427
2	H	0.55	0/275	0.92	0/427
All	All	0.51	0/3515	0.75	1/4938 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	438	ARG	NE-CZ-NH1	5.42	123.01	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	582	560	560	0	0
1	B	593	570	570	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	583	562	562	1	0
1	F	602	576	576	0	0
2	C	221	111	111	1	0
2	D	265	133	133	2	0
2	G	243	122	122	0	0
2	H	243	122	122	0	0
3	A	4	0	0	0	0
3	B	3	0	0	0	0
3	E	3	0	0	0	0
3	F	4	0	0	0	0
4	A	119	0	0	0	4
4	B	133	0	0	0	3
4	C	54	0	0	1	0
4	D	49	0	0	1	1
4	E	122	0	0	0	7
4	F	130	0	0	0	6
4	G	46	0	0	0	1
4	H	32	0	0	0	0
All	All	4031	2756	2756	3	11

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:3:A:OP1	4:D:101:HOH:O	2.13	0.65
2:D:8[A]:A:C5'	1:E:425:ARG:HH22	2.33	0.41
2:C:8:A:N3	4:C:102:HOH:O	2.37	0.41

The worst 5 of 11 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:602:HOH:O	4:E:707:HOH:O[2_756]	1.86	0.34
4:A:612:HOH:O	4:F:717:HOH:O[1_655]	1.87	0.33
4:A:693:HOH:O	4:E:687:HOH:O[2_756]	1.90	0.30
4:A:697:HOH:O	4:E:633:HOH:O[2_756]	1.94	0.26
4:E:632:HOH:O	4:F:693:HOH:O[2_646]	1.99	0.21

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	70/77 (91%)	70 (100%)	0	0	100	100
1	B	72/77 (94%)	72 (100%)	0	0	100	100
1	E	70/77 (91%)	70 (100%)	0	0	100	100
1	F	73/77 (95%)	73 (100%)	0	0	100	100
All	All	285/308 (92%)	285 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	65/69 (94%)	65 (100%)	0	100	100
1	B	66/69 (96%)	66 (100%)	0	100	100
1	E	65/69 (94%)	65 (100%)	0	100	100
1	F	67/69 (97%)	67 (100%)	0	100	100
All	All	263/276 (95%)	263 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	9/12 (75%)	2 (22%)	0
2	D	9/12 (75%)	2 (22%)	0
2	G	8/12 (66%)	2 (25%)	0
2	H	8/12 (66%)	2 (25%)	0
All	All	34/48 (70%)	8 (23%)	0

5 of 8 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	7	A
2	C	10	A
2	D	7	A
2	D	10	A
2	G	7	A

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 14 ligands modelled in this entry, 14 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	72/77 (93%)	0.09	3 (4%) 37 42	17, 21, 46, 56	0
1	B	74/77 (96%)	-0.04	4 (5%) 26 31	17, 23, 49, 66	0
1	E	72/77 (93%)	0.02	4 (5%) 25 29	16, 21, 44, 68	0
1	F	75/77 (97%)	-0.13	2 (2%) 55 62	17, 23, 37, 61	0
2	C	10/12 (83%)	-0.17	1 (10%) 8 9	18, 24, 29, 52	0
2	D	11/12 (91%)	0.18	1 (9%) 10 12	20, 23, 39, 69	0
2	G	10/12 (83%)	0.05	1 (10%) 8 9	19, 29, 39, 39	0
2	H	10/12 (83%)	-0.18	0 100 100	18, 33, 41, 41	0
All	All	334/356 (93%)	-0.02	16 (4%) 31 37	16, 23, 48, 69	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	478	VAL	7.6
2	D	2	A	7.1
1	A	479	LEU	6.8
1	B	478	VAL	6.2
1	E	478	VAL	5.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ZN	E	501	1/1	1.00	0.11	4.16	17,17,17,17	0
3	ZN	A	503	1/1	1.00	0.10	3.49	17,17,17,17	0
3	ZN	F	503	1/1	1.00	0.09	3.11	17,17,17,17	0
3	ZN	B	502	1/1	1.00	0.10	2.71	19,19,19,19	0
3	ZN	F	504	1/1	1.00	0.10	2.05	19,19,19,19	0
3	ZN	A	501	1/1	1.00	0.10	1.87	17,17,17,17	0
3	ZN	E	503	1/1	1.00	0.11	1.79	16,16,16,16	0
3	ZN	B	501	1/1	1.00	0.10	1.63	16,16,16,16	0
3	ZN	B	503	1/1	1.00	0.09	1.48	20,20,20,20	0
3	ZN	E	502	1/1	1.00	0.10	1.45	20,20,20,20	0
3	ZN	A	502	1/1	1.00	0.10	0.96	17,17,17,17	0
3	ZN	F	501	1/1	1.00	0.10	0.36	17,17,17,17	0
3	ZN	F	502	1/1	1.00	0.07	0.05	23,23,23,23	0
3	ZN	A	504	1/1	1.00	0.05	-	33,33,33,33	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.