



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 15, 2017 – 01:26 am GMT

PDB ID : 4L4V
Title : Structure of human MAIT TCR in complex with human MR1-RL-6-Me-7-OH
Authors : Patel, O.; Kjer-Nielsen, L.; Le Nours, J.; Eckle, S.B.G.; Birkinshaw, R.W.; Beddoe, T.; Corbett, A.J.; Liu, L.; Miles, J.J.; Meehan, B.; Reantragoon, R.; Sandoval-Romero, M.L.; Sullivan, L.C.; Brooks, A.G.; Chen, Z.; Fairlie, D.P.; McCluskey, J.; Rossjohn, J.
Deposited on : 2013-06-09
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	<div> <div>89%</div> <div>7% . .</div> </div>
1	C	271	<div> <div>3%</div> <div>85%</div> <div>10% 6%</div> </div>
2	B	99	<div> <div>96%</div> <div>.</div> </div>
2	F	99	<div> <div>%</div> <div>94%</div> <div>. .</div> </div>
3	D	203	<div> <div>%</div> <div>91%</div> <div>7% .</div> </div>
3	G	203	<div> <div>8%</div> <div>86%</div> <div>7% 7%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	E	245	
4	H	245	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	E	301	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 13688 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major histocompatibility complex class I-related gene protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	262	Total	C	N	O	S	0	0	0
			2139	1374	368	387	10			
1	C	256	Total	C	N	O	S	0	0	0
			2075	1327	363	374	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	EXPRESSION TAG	UNP Q95460
A	261	SER	CYS	ENGINEERED MUTATION	UNP Q95460
C	0	MET	-	EXPRESSION TAG	UNP Q95460
C	261	SER	CYS	ENGINEERED MUTATION	UNP Q95460

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	0	0
			792	507	137	145	3			
2	F	96	Total	C	N	O	S	0	0	0
			769	494	130	143	2			


- Molecule 3 is a protein called MAIT T-cell receptor alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	200	Total	C	N	O	S	0	3	0
			1552	983	247	313	9			
3	G	188	Total	C	N	O	S	0	2	0
			1418	913	223	273	9			

- Molecule 4 is a protein called MAIT T-cell receptor beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	243	Total	C	N	O	S	0	3	0
			1913	1205	331	368	9			
4	H	240	Total	C	N	O	S	0	4	0
			1864	1181	318	356	9			

- Molecule 5 is 1-DEOXY-1-(7-HYDROXY-6-METHYL-2,4-DIOXO-3,4-DIHYDROPTERIDIN-8(2H)-YL)-D-RIBITOL (three-letter code: 1VY) (formula: $C_{12}H_{16}N_4O_7$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			23	12	4	7		
5	C	1	Total	C	N	O	0	0
			23	12	4	7		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	201	Total	O	0	0
			201	201		
7	B	73	Total	O	0	0
			73	73		
7	C	191	Total	O	0	0
			191	191		
7	D	157	Total	O	0	0
			157	157		
7	E	189	Total	O	0	0
			189	189		
7	F	56	Total	O	0	0
			56	56		
7	G	84	Total	O	0	0
			84	84		
7	H	151	Total	O	0	0
			151	151		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major histocompatibility complex class I-related gene protein

Chain A: 



- Molecule 1: Major histocompatibility complex class I-related gene protein

Chain C: 




- Molecule 2: Beta-2-microglobulin

Chain B: 



- Molecule 2: Beta-2-microglobulin

Chain F: 




- Molecule 3: MAIT T-cell receptor alpha chain


Chain D: 




- Molecule 3: MAIT T-cell receptor alpha chain



- Molecule 4: MAIT T-cell receptor beta chain



- Molecule 4: MAIT T-cell receptor beta chain



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	218.37Å 71.20Å 144.64Å 90.00° 104.86° 90.00°	Depositor
Resolution (Å)	40.45 – 1.90 40.45 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (40.45-1.90) 99.7 (40.45-1.90)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 1.89Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.177 , 0.209 0.187 , 0.221	Depositor DCC
R_{free} test set	8496 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	21.8	Xtriage
Anisotropy	0.391	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 47.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13688	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 1VY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.54	0/2203	0.62	0/2995
1	C	0.51	0/2135	0.62	0/2901
2	B	0.47	0/815	0.61	0/1110
2	F	0.44	0/792	0.60	0/1080
3	D	0.52	0/1596	0.64	0/2167
3	G	0.51	0/1456	0.66	0/1978
4	E	0.52	0/1973	0.65	0/2687
4	H	0.50	0/1927	0.66	0/2628
All	All	0.51	0/12897	0.63	0/17546

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2139	0	2024	15	0
1	C	2075	0	1952	12	0
2	B	792	0	737	2	0
2	F	769	0	713	2	0
3	D	1552	0	1464	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	1418	0	1324	7	0
4	E	1913	0	1807	4	0
4	H	1864	0	1748	7	0
5	A	23	0	16	4	0
5	C	23	0	16	2	0
6	B	12	0	16	0	0
6	E	6	0	8	0	0
7	A	201	0	0	2	0
7	B	73	0	0	0	0
7	C	191	0	0	1	0
7	D	157	0	0	0	0
7	E	189	0	0	1	0
7	F	56	0	0	0	0
7	G	84	0	0	0	0
7	H	151	0	0	0	0
All	All	13688	0	11825	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:ARG:HH22	5:A:301:1VY:H14	1.38	0.88
3:G:1:GLY:H2	3:G:27:SER:H	1.17	0.87
3:G:1:GLY:N	3:G:27:SER:H	1.80	0.80
1:C:94:ARG:HH22	5:C:301:1VY:H14	1.48	0.78
1:C:198:LEU:HG	1:C:246:LEU:HD11	1.68	0.75
3:D:1:GLY:HA2	3:D:27:SER:H	1.54	0.72
3:D:19:GLN:HE21	3:D:21:ASN:HD21	1.37	0.70
1:A:198:LEU:HD22	1:A:246:LEU:HD21	1.76	0.68
3:D:150:ASP:HB2	3:D:177:LYS:HD2	1.78	0.66
3:G:146:SER:HB3	3:G:153:ILE:HG12	1.79	0.64
1:C:217:ASN:HD21	1:C:252:ASN:HD22	1.47	0.60
1:A:127:LEU:HG	7:A:553:HOH:O	2.00	0.60
4:E:12:VAL:HG23	7:E:584:HOH:O	2.02	0.59
4:H:128:VAL:HG23	4:H:238:ALA:HB3	1.84	0.58
7:C:581:HOH:O	4:E:97[A]:THR:HG21	2.05	0.56
1:C:215:MET:HG3	1:C:257:HIS:CD2	2.42	0.54
3:G:28:GLY:HA3	3:G:93[B]:SER:OG	2.08	0.53
1:C:217:ASN:HD21	1:C:252:ASN:ND2	2.06	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:28:GLY:HA3	3:D:93[A]:SER:OG	2.08	0.53
1:A:152:TYR:CD1	4:H:100:GLY:HA3	2.44	0.53
1:A:43:LYS:HD2	1:A:62:TYR:HB3	1.91	0.52
1:C:152:TYR:CD1	4:E:100:GLY:HA3	2.44	0.52
1:C:96:ILE:HD13	5:C:301:1VY:H15	1.92	0.51
4:H:136:ILE:HG23	4:H:199:ALA:HB1	1.92	0.51
4:E:128:VAL:HG23	4:E:238:ALA:HB3	1.93	0.51
1:A:96:ILE:HD13	5:A:301:1VY:H15	1.95	0.49
1:A:151:LEU:HD22	3:G:51:LEU:HD12	1.94	0.49
1:A:1:ARG:HB2	7:A:450:HOH:O	2.13	0.48
1:A:186:VAL:HG11	1:A:269:VAL:HG22	1.96	0.48
1:C:77:LEU:HD13	1:C:92:TYR:HB2	1.95	0.47
1:C:113:ALA:HB2	2:F:60:TRP:CE2	2.50	0.47
1:A:4:SER:HB3	1:A:99:GLU:HG2	1.96	0.46
3:G:50:VAL:O	3:G:66:ARG:HD3	2.17	0.45
4:H:154:ASP:CG	4:H:177:PRO:HG3	2.36	0.45
1:C:28:VAL:HG23	1:C:33:ILE:HD13	1.98	0.44
3:G:54:LEU:HD13	3:G:63:PHE:HB2	1.99	0.44
1:A:43:LYS:HZ1	5:A:301:1VY:H1	1.81	0.44
3:D:91:LYS:HG2	3:D:95:TYR:HA	1.99	0.44
1:C:6:ARG:NH2	2:F:58:LYS:HD3	2.33	0.44
1:C:43:LYS:HD2	1:C:62:TYR:HB3	1.99	0.43
4:H:8:PRO:HD2	4:H:21:LEU:HD22	2.00	0.43
3:D:3:ASN:ND2	3:D:5:ASP:OD1	2.51	0.43
1:A:113:ALA:HB2	2:B:60:TRP:CE2	2.54	0.43
2:B:77:GLU:HG2	2:B:97:ARG:HH12	1.83	0.43
3:D:6:GLN:HE21	3:D:100:GLY:HA3	1.83	0.42
1:A:43:LYS:NZ	5:A:301:1VY:H1	2.35	0.42
4:H:46:ILE:HG22	4:H:47:TYR:HD2	1.86	0.41
1:A:246:LEU:HD22	1:A:254:TYR:CE2	2.57	0.40
4:H:155:HIS:HB3	4:H:216:TYR:HB2	2.04	0.40
1:A:100:LEU:HD23	1:A:100:LEU:HA	1.98	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/271 (94%)	252 (98%)	4 (2%)	0	100	100
1	C	248/271 (92%)	246 (99%)	2 (1%)	0	100	100
2	B	97/99 (98%)	96 (99%)	1 (1%)	0	100	100
2	F	94/99 (95%)	93 (99%)	1 (1%)	0	100	100
3	D	201/203 (99%)	197 (98%)	4 (2%)	0	100	100
3	G	184/203 (91%)	179 (97%)	4 (2%)	1 (0%)	32	20
4	E	244/245 (100%)	241 (99%)	3 (1%)	0	100	100
4	H	242/245 (99%)	238 (98%)	4 (2%)	0	100	100
All	All	1566/1636 (96%)	1542 (98%)	23 (2%)	1 (0%)	55	45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	G	2	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/241 (93%)	216 (96%)	8 (4%)	40	29
1	C	216/241 (90%)	207 (96%)	9 (4%)	34	23
2	B	85/94 (90%)	84 (99%)	1 (1%)	75	75
2	F	83/94 (88%)	82 (99%)	1 (1%)	75	75
3	D	173/180 (96%)	172 (99%)	1 (1%)	89	90
3	G	148/180 (82%)	146 (99%)	2 (1%)	71	69
4	E	207/211 (98%)	203 (98%)	4 (2%)	62	57
4	H	199/211 (94%)	194 (98%)	5 (2%)	53	45

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1335/1452 (92%)	1304 (98%)	31 (2%)	56 49

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	ASP
1	A	73	PHE
1	A	75	VAL
1	A	100	LEU
1	A	179	THR
1	A	198	LEU
1	A	221	ILE
1	A	246	LEU
2	B	70	PHE
1	C	73	PHE
1	C	82	ARG
1	C	100	LEU
1	C	169	LEU
1	C	175	THR
1	C	179	THR
1	C	183	LEU
1	C	196	THR
1	C	268	GLN
3	D	146	SER
4	E	97[A]	THR
4	E	97[B]	THR
4	E	185	ASN
4	E	194	ARG
2	F	70	PHE
3	G	78	LEU
3	G	157	CYS
4	H	19	MET
4	H	99	GLU
4	H	155	HIS
4	H	184	LEU
4	H	194	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	264	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	111	GLN
1	C	203	HIS
1	C	252	ASN
1	C	264	HIS
3	D	6	GLN
3	D	21	ASN
4	E	22	GLN
2	F	51	HIS
4	H	11	GLN
4	H	155	HIS
4	H	185	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	1VY	A	301	-	21,24,24	1.62	2 (9%)	19,35,35	2.09	5 (26%)
6	GOL	B	601	-	5,5,5	0.11	0	5,5,5	0.29	0
6	GOL	B	602	-	5,5,5	0.21	0	5,5,5	0.59	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	1VY	C	301	-	21,24,24	1.69	4 (19%)	19,35,35	2.34	6 (31%)
6	GOL	E	301	-	5,5,5	0.19	0	5,5,5	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	1VY	A	301	-	-	0/12/14/14	0/2/2/2
6	GOL	B	601	-	-	0/4/4/4	0/0/0/0
6	GOL	B	602	-	-	0/4/4/4	0/0/0/0
5	1VY	C	301	-	-	0/12/14/14	0/2/2/2
6	GOL	E	301	-	-	0/4/4/4	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	301	1VY	C9-C6	-4.76	1.41	1.50
5	A	301	1VY	C9-C6	-4.56	1.42	1.50
5	C	301	1VY	C8A-N1	2.08	1.36	1.33
5	C	301	1VY	C4A-N5	2.21	1.36	1.33
5	C	301	1VY	C6-N5	3.01	1.35	1.32
5	A	301	1VY	C6-N5	3.31	1.35	1.32

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	301	1VY	C4A-C4-N3	-3.28	118.81	123.48
5	A	301	1VY	C4A-C4-N3	-2.59	119.80	123.48
5	C	301	1VY	O4'-C4'-C3'	-2.25	103.50	109.09
5	A	301	1VY	C4-C4A-N5	2.04	120.92	118.68
5	C	301	1VY	C4-C4A-N5	2.21	121.10	118.68
5	C	301	1VY	C9-C6-N5	2.47	120.65	116.99
5	C	301	1VY	O5'-C5'-C4'	2.65	116.96	111.11
5	A	301	1VY	C9-C6-N5	2.74	121.05	116.99
5	A	301	1VY	C1'-N8-C8A	3.63	122.22	118.50
5	A	301	1VY	C4-N3-C2	5.68	120.12	115.16
5	C	301	1VY	C4-N3-C2	7.31	121.55	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	301	1VY	4	0
5	C	301	1VY	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/271 (96%)	-0.31	1 (0%) 92 93	12, 22, 40, 68	3 (1%)
1	C	256/271 (94%)	-0.17	7 (2%) 55 59	14, 24, 48, 75	1 (0%)
2	B	99/99 (100%)	-0.29	0 100 100	14, 26, 47, 59	0
2	F	96/99 (96%)	0.26	1 (1%) 82 84	18, 40, 61, 82	0
3	D	200/203 (98%)	-0.16	3 (1%) 74 77	14, 24, 50, 60	0
3	G	188/203 (92%)	0.25	17 (9%) 10 11	13, 32, 70, 82	0
4	E	243/245 (99%)	-0.36	5 (2%) 64 67	13, 23, 43, 69	1 (0%)
4	H	240/245 (97%)	-0.15	5 (2%) 64 67	13, 28, 62, 80	1 (0%)
All	All	1584/1636 (96%)	-0.15	39 (2%) 58 62	12, 25, 58, 82	6 (0%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	196	THR	6.2
3	G	149	SER	6.0
3	D	179	ASP	4.7
1	C	246	LEU	4.7
3	G	164	MET	4.3
4	H	205	PRO	4.3
3	G	163	SER	4.0
1	C	244	ILE	3.9
3	G	182	CYS	3.8
3	G	180	PHE	3.8
3	G	190	ILE	3.7
1	A	222	VAL	3.5
4	E	184	LEU	3.1
3	G	146	SER	3.1
1	C	197	ALA	3.0
4	H	221	ASN	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	251	SER	2.8
3	G	112	GLN	2.8
3	G	1	GLY	2.8
4	H	136	ILE	2.8
3	D	1	GLY	2.7
4	E	185	ASN	2.7
4	E	183	ALA	2.6
1	C	18	GLY	2.6
4	E	186	ASP	2.5
4	H	220	GLU	2.5
4	H	240	ALA	2.4
2	F	44	GLU	2.4
3	G	184	ASN	2.4
3	D	149	SER	2.4
3	G	196	PHE	2.4
1	C	16	ILE	2.3
3	G	197	PHE	2.3
4	E	2	ALA	2.3
3	G	191	ILE	2.2
3	G	183	ALA	2.2
3	G	181	ALA	2.1
3	G	185	ALA	2.1
3	G	151	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	GOL	E	301	6/6	0.88	0.18	2.77	27,33,35,36	0
6	GOL	B	602	6/6	0.95	0.11	1.89	23,25,36,41	0
5	1VY	A	301	23/23	0.96	0.14	1.53	15,21,29,30	0
5	1VY	C	301	23/23	0.97	0.11	0.52	14,19,27,29	0
6	GOL	B	601	6/6	0.97	0.08	-0.23	17,22,24,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.