



# Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 11:46 pm GMT

PDB ID : 2L7J  
Title : Solution structure of the third Immunoglobulin-like domain of nectin-1  
Authors : Clausen, O.; Poulsen, F.M.  
Deposited on : 2010-12-13

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : trunk28760  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 50%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	95	 <div>42%</div> <div>46%</div> <div>• 7%</div>

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:244-A:254, A:258-A:334 (88)	0.33	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 4, 5, 6, 8, 9, 11, 12, 13, 15, 16, 17, 18, 20
2	7, 10, 14, 19
Single-model clusters	3

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1466 atoms, of which 718 are hydrogens and 0 are deuteriums.


- Molecule 1 is a protein called Poliovirus receptor-related 1.

Mol	Chain	Residues	Atoms						Trace
1	A	95	Total	C	H	N	O	S	0
			1466	472	718	127	147	2	

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	335	HIS	-	EXPRESSION TAG	UNP Q6P9M9





#### 4.2.3 Score per residue for model 3

- Molecule 1: Poliovirus receptor-related 1


Chain A: 42% 46% 7%



#### 4.2.4 Score per residue for model 4

- Molecule 1: Poliovirus receptor-related 1


Chain A: 44% 42% 6% 7%



#### 4.2.5 Score per residue for model 5


- Molecule 1: Poliovirus receptor-related 1

Chain A: 43% 44% 5% 7%




#### 4.2.6 Score per residue for model 6

- Molecule 1: Poliovirus receptor-related 1




#### 4.2.7 Score per residue for model 7

- Molecule 1: Poliovirus receptor-related 1




#### 4.2.8 Score per residue for model 8

- Molecule 1: Poliovirus receptor-related 1




#### 4.2.9 Score per residue for model 9

- Molecule 1: Poliovirus receptor-related 1




### 4.2.10 Score per residue for model 10

- Molecule 1: Poliovirus receptor-related 1




### 4.2.11 Score per residue for model 11

- Molecule 1: Poliovirus receptor-related 1




### 4.2.12 Score per residue for model 12

- Molecule 1: Poliovirus receptor-related 1



### 4.2.13 Score per residue for model 13


- Molecule 1: Poliovirus receptor-related 1






### 4.2.14 Score per residue for model 14

- Molecule 1: Poliovirus receptor-related 1




### 4.2.15 Score per residue for model 15

- Molecule 1: Poliovirus receptor-related 1




### 4.2.16 Score per residue for model 16

- Molecule 1: Poliovirus receptor-related 1




### 4.2.17 Score per residue for model 17

- Molecule 1: Poliovirus receptor-related 1




### 4.2.18 Score per residue for model 18

- Molecule 1: Poliovirus receptor-related 1




### 4.2.19 Score per residue for model 19

- Molecule 1: Poliovirus receptor-related 1



### 4.2.20 Score per residue for model 20

- Molecule 1: Poliovirus receptor-related 1



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA	refinement	
CYANA	structure solution	
CYANA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2l7j_cs.str
Number of chemical shift lists	1
Total number of shifts	627
Number of shifts mapped to atoms	627
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	50%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	694	669	668	18±3
All	All	13880	13380	13360	361

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:253:GLY:HA3	1:A:267:LEU:HD23	0.78	1.55	6	4
1:A:274:ASN:HB3	1:A:275:PRO:HD3	0.73	1.59	9	19
1:A:311:ALA:HA	1:A:331:VAL:HG22	0.70	1.64	19	6
1:A:258:TRP:HZ2	1:A:267:LEU:HD21	0.68	1.46	18	16
1:A:254:PHE:HB3	1:A:258:TRP:NE1	0.67	2.05	5	5
1:A:313:THR:HG23	1:A:330:GLU:HG2	0.66	1.67	1	2
1:A:306:ILE:HG23	1:A:333:ILE:HD12	0.66	1.67	13	18
1:A:311:ALA:HB2	1:A:333:ILE:HG12	0.65	1.68	10	5
1:A:265:VAL:HB	1:A:306:ILE:HD12	0.62	1.70	6	6
1:A:247:PRO:HB2	1:A:271:ALA:HB1	0.62	1.70	9	13
1:A:313:THR:HG23	1:A:330:GLU:HG3	0.62	1.70	19	3
1:A:265:VAL:HG11	1:A:306:ILE:HB	0.62	1.71	7	17
1:A:258:TRP:CZ2	1:A:267:LEU:HD21	0.61	2.30	18	19
1:A:259:TYR:HD1	1:A:262:ARG:HD2	0.60	1.56	9	1

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:247:PRO:HA	1:A:273:ALA:HB2	0.60	1.73	10	20
1:A:254:PHE:HB3	1:A:258:TRP:CD1	0.59	2.32	5	2
1:A:254:PHE:HB3	1:A:267:LEU:HD23	0.59	1.74	16	1
1:A:258:TRP:CZ3	1:A:331:VAL:HA	0.59	2.33	19	10
1:A:280:TYR:HE2	1:A:298:ARG:HB3	0.59	1.56	17	1
1:A:252:GLU:HB2	1:A:268:THR:HB	0.58	1.75	13	1
1:A:320:ASN:HB3	1:A:322:ILE:HD12	0.55	1.77	19	10
1:A:248:GLU:O	1:A:271:ALA:HA	0.55	2.02	5	7
1:A:275:PRO:HG2	1:A:322:ILE:HD11	0.54	1.80	15	4
1:A:258:TRP:CE3	1:A:331:VAL:HA	0.54	2.38	19	6
1:A:258:TRP:HD1	1:A:265:VAL:HG12	0.54	1.63	7	4
1:A:280:TYR:CE2	1:A:298:ARG:HB3	0.54	2.37	17	1
1:A:266:LYS:HA	1:A:302:PHE:CD1	0.53	2.39	20	11
1:A:244:GLN:HG2	1:A:274:ASN:HD22	0.53	1.62	7	1
1:A:292:GLY:O	1:A:303:ARG:HB2	0.52	2.05	7	6
1:A:279:GLU:CB	1:A:319:THR:HB	0.52	2.35	8	10
1:A:315:ILE:HG23	1:A:328:GLN:HB3	0.52	1.81	2	9
1:A:245:TYR:CE1	1:A:273:ALA:HA	0.52	2.40	18	2
1:A:245:TYR:O	1:A:322:ILE:HD13	0.51	2.06	7	16
1:A:262:ARG:HB3	1:A:265:VAL:HG13	0.51	1.81	10	1
1:A:269:CYS:O	1:A:298:ARG:HD3	0.51	2.06	4	1
1:A:247:PRO:HD3	1:A:320:ASN:HB2	0.51	1.81	2	6
1:A:294:GLU:HB2	1:A:303:ARG:HG2	0.50	1.83	13	1
1:A:316:CYS:O	1:A:326:SER:HA	0.50	2.06	9	10
1:A:296:GLN:O	1:A:296:GLN:HG2	0.50	2.04	8	1
1:A:260:LEU:O	1:A:305:PRO:HB2	0.50	2.07	20	2
1:A:254:PHE:HB2	1:A:265:VAL:HA	0.49	1.82	5	1
1:A:260:LEU:HD23	1:A:306:ILE:O	0.49	2.07	19	1
1:A:314:TYR:O	1:A:328:GLN:HA	0.49	2.08	2	5
1:A:267:LEU:HD13	1:A:329:VAL:HG11	0.48	1.85	10	2
1:A:285:LEU:HD12	1:A:315:ILE:HD11	0.48	1.86	8	1
1:A:262:ARG:O	1:A:305:PRO:HA	0.48	2.09	9	3
1:A:282:TRP:HE1	1:A:300:LEU:HD21	0.47	1.70	11	1
1:A:315:ILE:HA	1:A:327:GLY:O	0.47	2.10	12	10
1:A:259:TYR:CD1	1:A:262:ARG:HD2	0.46	2.44	9	1
1:A:259:TYR:HA	1:A:334:THR:OG1	0.46	2.10	3	4
1:A:297:ASN:OD1	1:A:298:ARG:HD2	0.46	2.10	12	1
1:A:248:GLU:HA	1:A:248:GLU:OE1	0.45	2.11	11	1
1:A:281:HIS:HB3	1:A:317:GLU:OE2	0.45	2.12	15	1
1:A:280:TYR:HB3	1:A:282:TRP:CZ3	0.45	2.47	12	1
1:A:274:ASN:CB	1:A:275:PRO:HD3	0.44	2.39	5	6

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:250:THR:HB	1:A:270:LYS:HG3	0.44	1.89	17	1
1:A:279:GLU:HB2	1:A:319:THR:HB	0.44	1.89	13	5
1:A:260:LEU:HG	1:A:333:ILE:HG22	0.44	1.89	15	1
1:A:252:GLU:O	1:A:267:LEU:HA	0.44	2.12	3	1
1:A:259:TYR:HA	1:A:334:THR:HB	0.44	1.90	2	1
1:A:320:ASN:OD1	1:A:321:PRO:HD2	0.43	2.13	2	1
1:A:269:CYS:O	1:A:298:ARG:HB2	0.43	2.14	17	1
1:A:260:LEU:HG	1:A:333:ILE:CG2	0.43	2.43	5	1
1:A:306:ILE:CG2	1:A:333:ILE:HD12	0.42	2.44	16	3
1:A:279:GLU:HB3	1:A:319:THR:HB	0.42	1.91	8	1
1:A:266:LYS:HB2	1:A:300:LEU:O	0.42	2.13	18	2
1:A:260:LEU:HA	1:A:306:ILE:HG22	0.42	1.91	8	1
1:A:250:THR:HB	1:A:270:LYS:CG	0.42	2.44	17	1
1:A:302:PHE:CE2	1:A:306:ILE:HG13	0.42	2.49	9	3
1:A:265:VAL:HG23	1:A:302:PHE:CB	0.42	2.44	4	1
1:A:258:TRP:CD1	1:A:306:ILE:HD12	0.42	2.49	2	4
1:A:251:ILE:HG13	1:A:329:VAL:HB	0.41	1.92	6	2
1:A:310:LEU:HA	1:A:310:LEU:HD12	0.41	1.78	18	1
1:A:289:LEU:HG	1:A:290:PRO:HD2	0.41	1.92	19	1
1:A:285:LEU:HA	1:A:285:LEU:HD23	0.41	1.78	5	1
1:A:259:TYR:CD2	1:A:262:ARG:HG3	0.41	2.51	16	1
1:A:313:THR:HA	1:A:330:GLU:HA	0.41	1.93	14	1
1:A:296:GLN:OE1	1:A:299:THR:HB	0.41	2.16	8	1
1:A:252:GLU:O	1:A:268:THR:N	0.40	2.51	20	1
1:A:308:TYR:HA	1:A:333:ILE:HG13	0.40	1.92	3	4
1:A:320:ASN:HB3	1:A:321:PRO:HD2	0.40	1.93	11	1
1:A:259:TYR:HB2	1:A:262:ARG:HB2	0.40	1.94	15	1
1:A:250:THR:HB	1:A:270:LYS:HE3	0.40	1.94	15	1

## 6.3 Torsion angles ⓘ

### 6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	88/95 (93%)	82±1 (93±1%)	5±1 (5±1%)	1±0 (1±0%)	20	66

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1760/1900 (93%)	1644 (93%)	95 (5%)	21 (1%)	20 66

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	276	PRO	20
1	A	334	THR	1

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	75/81 (93%)	47±2 (63±3%)	28±2 (37±3%)	1 7
All	All	1500/1620 (93%)	939 (63%)	561 (37%)	1 7

All 56 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	263	THR	20
1	A	307	THR	20
1	A	310	LEU	20
1	A	300	LEU	20
1	A	266	LYS	20
1	A	283	THR	20
1	A	326	SER	20
1	A	280	TYR	19
1	A	309	SER	18
1	A	270	LYS	17
1	A	331	VAL	17
1	A	288	SER	16
1	A	285	LEU	16
1	A	322	ILE	15
1	A	298	ARG	15
1	A	313	THR	15
1	A	334	THR	15

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	303	ARG	14
1	A	289	LEU	14
1	A	279	GLU	13
1	A	301	PHE	13
1	A	262	ARG	13
1	A	316	CYS	13
1	A	282	TRP	11
1	A	264	ASP	11
1	A	250	THR	11
1	A	281	HIS	10
1	A	325	ARG	9
1	A	296	GLN	9
1	A	294	GLU	9
1	A	330	GLU	8
1	A	332	ASN	8
1	A	299	THR	8
1	A	291	LYS	8
1	A	317	GLU	8
1	A	246	GLU	7
1	A	244	GLN	7
1	A	272	ASP	7
1	A	254	PHE	7
1	A	286	ASN	5
1	A	328	GLN	4
1	A	308	TYR	3
1	A	269	CYS	3
1	A	319	THR	3
1	A	259	TYR	3
1	A	260	LEU	3
1	A	274	ASN	3
1	A	329	VAL	2
1	A	261	GLN	2
1	A	315	ILE	2
1	A	248	GLU	2
1	A	306	ILE	1
1	A	293	VAL	1
1	A	267	LEU	1
1	A	297	ASN	1
1	A	268	THR	1



### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 50% for the well-defined parts and 49% for the entire structure.

### 7.1 Chemical shift list 1

File name: 2l7j\_cs.str

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	627
Number of shifts mapped to atoms	627
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	7

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	89	$0.28 \pm 0.12$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	81	$-0.12 \pm 0.18$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
$^{15}\text{N}$	84	$-0.18 \pm 0.56$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 50%, i.e. 526 atoms were assigned a chemical shift out of a possible 1061. 0 out of 11 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	314/428 (73%)	153/170 (90%)	83/176 (47%)	78/82 (95%)
Sidechain	212/534 (40%)	136/312 (44%)	76/198 (38%)	0/24 (0%)

*Continued on next page...*

*Continued from previous page...*

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Aromatic	0/99 (0%)	0/51 (0%)	0/44 (0%)	0/4 (0%)
Overall	526/1061 (50%)	289/533 (54%)	159/418 (38%)	78/110 (71%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 49%, i.e. 558 atoms were assigned a chemical shift out of a possible 1140. 0 out of 13 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	336/463 (73%)	163/184 (89%)	89/190 (47%)	84/89 (94%)
Sidechain	222/570 (39%)	141/332 (42%)	81/212 (38%)	0/26 (0%)
Aromatic	0/107 (0%)	0/55 (0%)	0/46 (0%)	0/6 (0%)
Overall	558/1140 (49%)	304/571 (53%)	170/448 (38%)	84/121 (69%)

#### 7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	307	THR	HB	1.17	5.82 – 2.52	-9.1
1	A	300	LEU	HD21	-1.19	2.14 – -0.66	-6.9
1	A	300	LEU	HD22	-1.19	2.14 – -0.66	-6.9
1	A	300	LEU	HD23	-1.19	2.14 – -0.66	-6.9
1	A	265	VAL	HG22	-0.77	2.20 – -0.60	-5.6
1	A	265	VAL	HG23	-0.77	2.20 – -0.60	-5.6
1	A	265	VAL	HG21	-0.77	2.20 – -0.60	-5.6

#### 7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

