



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 05:38 am GMT

PDB ID : 4L7G  
Title : Diethylaminosulfur Trifluoride-Mediated Intramolecular Cyclization of 2-hydroxy-benzylureas to Fused Bicyclic Aminooxazoline Compounds and Evaluation of Their Biochemical Activity Against Beta-Secretase-1 (BACE1)  
Authors : Huestis, M.P.; Liu, W.; Volgraf, M.; Purkey, H.; Wang, W.; Yu, C.; Wu, P.; Smith, D.; Vigers, G.; Dutcher, D.; Geck Do, M.K.; Hunt, K.W.; Siu, M.  
Deposited on : 2013-06-13  
Resolution : 1.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2133 (1.40-1.36)
Clashscore	112137	2266 (1.40-1.36)
Ramachandran outliers	110173	2215 (1.40-1.36)
Sidechain outliers	110143	2214 (1.40-1.36)
RSRZ outliers	101464	2141 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	409	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	DMS	A	502	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3486 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-secretase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	373	3081	1979	510	575	17	0	18	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	EXPRESSION TAG	UNP P56817
A	-6	GLY	-	EXPRESSION TAG	UNP P56817
A	-5	SER	-	EXPRESSION TAG	UNP P56817
A	393	GLY	-	EXPRESSION TAG	UNP P56817
A	394	ASN	-	EXPRESSION TAG	UNP P56817
A	395	SER	-	EXPRESSION TAG	UNP P56817
A	396	HIS	-	EXPRESSION TAG	UNP P56817
A	397	HIS	-	EXPRESSION TAG	UNP P56817
A	398	HIS	-	EXPRESSION TAG	UNP P56817
A	399	HIS	-	EXPRESSION TAG	UNP P56817
A	400	HIS	-	EXPRESSION TAG	UNP P56817
A	401	HIS	-	EXPRESSION TAG	UNP P56817

- Molecule 2 is (3AS,7AR)-7A-[3-(PYRIMIDIN-5-YL)PHENYL]-3A,6,7,7A-TETRAHYDRO-4H-PYRANO[4,3-D][1,3]OXAZOL-2-AMINE (three-letter code: 1W0) (formula: C<sub>16</sub>H<sub>16</sub>N<sub>4</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			22	16	4	2		

- Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula:  $C_2H_6OS$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	379	Total 379	O 379	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Beta-secretase 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.66Å 104.47Å 50.08Å 90.00° 94.82° 90.00°	Depositor
Resolution (Å)	16.63 – 1.38 19.19 – 1.38	Depositor EDS
% Data completeness (in resolution range)	99.4 (16.63-1.38) 99.4 (19.19-1.38)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.13 (at 1.38Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, $R_{free}$	0.154 , 0.192 0.149 , 0.188	Depositor DCC
$R_{free}$ test set	3836 reflections (5.06%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.4	Xtriage
Anisotropy	0.542	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 60.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3486	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, 1W0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/3154	0.56	0/4282

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3081	0	3007	48	1
2	A	22	0	16	0	0
3	A	4	0	6	0	0
4	A	379	0	0	11	0
All	All	3486	0	3029	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:TYR:HA	4:A:768:HOH:O	1.60	1.00
1:A:2[B]:MET:HE1	1:A:176:ILE:H	1.34	0.92
1:A:288[B]:MET:HE2	4:A:835:HOH:O	1.76	0.84
1:A:267:LEU:HD22	1:A:309:VAL:HG21	1.65	0.78
1:A:236:LEU:HD12	1:A:324[B]:ILE:HD11	1.68	0.73
1:A:108:PHE:HZ	1:A:118[B]:ILE:HD12	1.55	0.71
1:A:208:ILE:C	1:A:209[B]:ASN:HD22	1.94	0.71
1:A:301:LEU:HD12	1:A:361:VAL:HG13	1.74	0.70
1:A:309:VAL:O	1:A:310:GLU:HB2	1.90	0.69
1:A:39:ALA:HB3	1:A:118[B]:ILE:HD11	1.78	0.64
1:A:300[B]:ILE:HD13	1:A:337:ILE:HD12	1.80	0.64
1:A:140:LEU:HG	1:A:146[B]:VAL:HG21	1.80	0.63
1:A:125:GLU:OE2	1:A:195:ARG:NH1	2.30	0.63
1:A:322:PHE:CZ	1:A:324[B]:ILE:CG2	2.86	0.58
1:A:209[B]:ASN:N	1:A:209[B]:ASN:HD22	2.01	0.57
1:A:292:THR:HB	4:A:833:HOH:O	2.03	0.57
1:A:7:ARG:CZ	4:A:975:HOH:O	2.53	0.56
1:A:267:LEU:CD2	1:A:309:VAL:HG21	2.35	0.54
1:A:340:GLY:C	1:A:357:SER:HB3	2.29	0.53
1:A:360:HIS:CE1	4:A:819:HOH:O	2.61	0.52
1:A:381:ASP:N	1:A:381:ASP:OD2	2.42	0.52
1:A:9:LYS:HZ3	1:A:169:SER:HA	1.75	0.52
1:A:300[A]:ILE:HD13	1:A:337:ILE:CD1	2.43	0.48
1:A:303:GLN:HB3	1:A:361:VAL:HG21	1.95	0.48
1:A:215[A]:MET:CE	1:A:239:LYS:HG2	2.44	0.47
1:A:39:ALA:HB3	1:A:118[B]:ILE:CD1	2.44	0.47
1:A:239:LYS:HE2	4:A:826:HOH:O	2.15	0.47
1:A:309:VAL:O	1:A:310:GLU:CB	2.63	0.46
1:A:283:ILE:HB	1:A:300[A]:ILE:HG12	1.98	0.45
1:A:9:LYS:NZ	1:A:9:LYS:HA	2.32	0.44
1:A:187:SER:HB2	4:A:825:HOH:O	2.18	0.44
1:A:303:GLN:CB	1:A:361:VAL:HG21	2.47	0.44
1:A:108:PHE:CZ	1:A:118[B]:ILE:HD12	2.43	0.44
1:A:340:GLY:O	1:A:357:SER:HB3	2.18	0.44
1:A:142[A]:LYS:HG3	4:A:734:HOH:O	2.17	0.44
1:A:223:ASP:O	1:A:385:ASN:HB3	2.18	0.44
1:A:215[A]:MET:HE2	1:A:219:GLU:CB	2.49	0.42
1:A:7:ARG:NH2	4:A:816:HOH:O	2.53	0.42
1:A:263[A]:LEU:HD22	1:A:263[A]:LEU:HA	1.93	0.42
1:A:215[A]:MET:HE3	1:A:239:LYS:HG2	2.02	0.41
1:A:254:THR:HB	4:A:959:HOH:O	2.20	0.41
1:A:381:ASP:N	4:A:918:HOH:O	2.52	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:254:THR:HG22	1:A:255:GLU:HG3	2.02	0.41
1:A:222:TYR:HA	1:A:223:ASP:HA	1.83	0.41
1:A:234:LEU:O	1:A:324[B]:ILE:HA	2.21	0.41
1:A:340:GLY:HA2	1:A:360:HIS:HA	2.02	0.41
1:A:322:PHE:CE2	1:A:324[B]:ILE:HG23	2.56	0.40
1:A:276:PRO:O	1:A:279:ILE:HG12	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:TYR:O	1:A:384:TYR:OH[2_656]	2.05	0.15

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	383/409 (94%)	371 (97%)	10 (3%)	2 (0%)	32 9

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	209[A]	ASN
1	A	209[B]	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	336/349 (96%)	329 (98%)	7 (2%)	59 23

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	VAL
1	A	9	LYS
1	A	197	TRP
1	A	254	THR
1	A	263[A]	LEU
1	A	263[B]	LEU
1	A	381	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	1W0	A	501	-	23,25,25	0.96	1 (4%)	26,36,36	2.56	11 (42%)
3	DMS	A	502	-	3,3,3	2.58	1 (33%)	3,3,3	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1W0	A	501	-	-	0/10/33/33	1/4/4/4
3	DMS	A	502	-	-	0/0/0/0	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	1W0	O13-C12	-2.84	1.42	1.46
3	A	502	DMS	O-S	4.34	1.79	1.50

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	1W0	N16-C14-N15	-4.66	123.49	127.58
2	A	501	1W0	N21-C20-N19	-4.07	121.93	126.92
2	A	501	1W0	C9-C8-C7	-3.95	106.48	111.51
2	A	501	1W0	O10-C11-C12	-2.40	106.24	111.60
2	A	501	1W0	C4-C5-C7	-2.01	119.46	121.22
2	A	501	1W0	C7-N15-C14	2.12	109.74	106.63
2	A	501	1W0	C22-C17-C18	2.18	117.58	114.42
2	A	501	1W0	C9-O10-C11	2.57	114.03	109.90
2	A	501	1W0	C22-N21-C20	3.97	120.46	115.84
2	A	501	1W0	C18-N19-C20	4.60	121.19	115.84
2	A	501	1W0	O13-C14-N16	6.54	121.81	114.69

There are no chirality outliers.

There are no torsion outliers.

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	1W0	C11-C12-C7-C8-C9-O10

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	373/409 (91%)	0.47	41 (10%) 6 7	11, 24, 45, 55	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	316	GLN	8.4
1	A	315	SER	6.6
1	A	362	HIS	6.1
1	A	365	PHE	6.0
1	A	73	GLN	5.8
1	A	254	THR	5.8
1	A	384	TYR	5.7
1	A	271	GLN	5.6
1	A	361	VAL	5.4
1	A	273	GLY	5.1
1	A	145	HIS	5.1
1	A	386	ILE	5.0
1	A	364	GLU	4.9
1	A	274	THR	4.5
1	A	72	THR	4.5
1	A	272	ALA	4.4
1	A	64	ARG	4.3
1	A	265	GLU	4.1
1	A	68	TYR	4.0
1	A	0	VAL	3.7
1	A	376	THR	3.7
1	A	157	ALA	3.7
1	A	74	GLY	3.5
1	A	324[A]	ILE	3.3
1	A	310	GLU	3.2
1	A	165	GLU	3.1
1	A	166	VAL	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	360	HIS	2.7
1	A	270	TRP	2.6
1	A	253	SER	2.6
1	A	317	ASP	2.5
1	A	381	ASP	2.4
1	A	46	PRO	2.4
1	A	256	LYS	2.4
1	A	278	ASN	2.3
1	A	263[A]	LEU	2.2
1	A	104	GLU	2.2
1	A	1	GLU	2.2
1	A	49	HIS	2.1
1	A	385	ASN	2.1
1	A	292	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	DMS	A	502	4/4	0.95	0.10	3.96	37,38,40,42	0
2	1W0	A	501	22/22	0.96	0.06	-0.27	16,19,21,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.