



# Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 11:49 pm GMT

PDB ID : 2L9Q  
Title : Structural Characterization of small heat shock protein (Hsp12)  
Authors : Singarapu, K.K.; Tonelli, M.; Westler, W.M.; Markley, J.L.; Center for Eukaryotic Structural Genomics (CESG)  
Deposited on : 2011-02-22

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : trunk28760  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 83%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	109	

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 11 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:9-A:18 (10)	0.05	11
2	A:25-A:43 (19)	0.12	18
3	A:51-A:61 (11)	0.59	18
4	A:79-A:100 (22)	0.10	11

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	2, 3, 4, 7, 10, 11, 15, 16, 19
2	6, 8, 12, 14, 18, 20
3	5, 9, 13
Single-model clusters	1; 17

### 3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 1452 atoms, of which 632 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called 12 kDa heat shock protein.


Mol	Chain	Residues	Atoms						Trace
1	A	109	Total	C	H	N	O	S	0
			1452	496	632	146	176	2	

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 12 kDa heat shock protein




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1

- Molecule 1: 12 kDa heat shock protein



#### 4.2.2 Score per residue for model 2


- Molecule 1: 12 kDa heat shock protein






### 4.2.3 Score per residue for model 3

- Molecule 1: 12 kDa heat shock protein




### 4.2.4 Score per residue for model 4

- Molecule 1: 12 kDa heat shock protein




### 4.2.5 Score per residue for model 5

- Molecule 1: 12 kDa heat shock protein




### 4.2.6 Score per residue for model 6

- Molecule 1: 12 kDa heat shock protein




#### 4.2.7 Score per residue for model 7

- Molecule 1: 12 kDa heat shock protein




#### 4.2.8 Score per residue for model 8

- Molecule 1: 12 kDa heat shock protein




#### 4.2.9 Score per residue for model 9

- Molecule 1: 12 kDa heat shock protein




### 4.2.10 Score per residue for model 10

- Molecule 1: 12 kDa heat shock protein




### 4.2.11 Score per residue for model 11 (medoid)

- Molecule 1: 12 kDa heat shock protein




### 4.2.12 Score per residue for model 12

- Molecule 1: 12 kDa heat shock protein



### 4.2.13 Score per residue for model 13


- Molecule 1: 12 kDa heat shock protein






### 4.2.14 Score per residue for model 14

- Molecule 1: 12 kDa heat shock protein




### 4.2.15 Score per residue for model 15

- Molecule 1: 12 kDa heat shock protein




### 4.2.16 Score per residue for model 16

- Molecule 1: 12 kDa heat shock protein




### 4.2.17 Score per residue for model 17

- Molecule 1: 12 kDa heat shock protein




### 4.2.18 Score per residue for model 18

- Molecule 1: 12 kDa heat shock protein




### 4.2.19 Score per residue for model 19

- Molecule 1: 12 kDa heat shock protein



### 4.2.20 Score per residue for model 20

- Molecule 1: 12 kDa heat shock protein



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Molmol	structure solution	
CYANA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2l9q_cs.str
Number of chemical shift lists	1
Total number of shifts	1103
Number of shifts mapped to atoms	1103
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	83%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	472	382	458	5±1
All	All	9440	7640	9160	98

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:92:ALA:O	1:A:96:VAL:HG23	0.64	1.93	10	20
1:A:89:LEU:O	1:A:93:VAL:HG23	0.63	1.94	9	19
1:A:36:LYS:O	1:A:40:VAL:HG12	0.57	2.00	20	20
1:A:17:LEU:HD12	1:A:18:LYS:N	0.54	2.17	12	19
1:A:29:GLY:O	1:A:33:ILE:HD12	0.52	2.04	15	8
1:A:53:PHE:O	1:A:56:VAL:HG12	0.51	2.05	20	5
1:A:17:LEU:HB3	1:A:60:ALA:HB1	0.49	1.85	11	1
1:A:17:LEU:HD12	1:A:18:LYS:HD2	0.46	1.86	9	1
1:A:89:LEU:C	1:A:89:LEU:HD13	0.44	2.32	1	1
1:A:56:VAL:HG22	1:A:81:TYR:OH	0.44	2.12	17	1
1:A:96:VAL:O	1:A:100:VAL:HG23	0.43	2.14	16	2
1:A:89:LEU:HD23	1:A:93:VAL:CG2	0.41	2.45	15	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	62/109 (57%)	61±1 (98±1%)	1±1 (2±1%)	0±1 (0±1%)	48	82
All	All	1240/2180 (57%)	1210 (98%)	26 (2%)	4 (0%)	48	82

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	51	GLY	3
1	A	100	VAL	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	46/83 (55%)	30±3 (64±6%)	16±3 (36±6%)	1	9
All	All	920/1660 (55%)	591 (64%)	329 (36%)	1	9

All 31 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	54	GLN	20
1	A	97	SER	20
1	A	94	GLU	20
1	A	9	PHE	20
1	A	91	ASP	18
1	A	80	ASP	17
1	A	38	ASP	17
1	A	25	TYR	17

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	14	SER	16
1	A	39	LYS	15
1	A	11	GLU	15
1	A	18	LYS	13
1	A	90	ASN	12
1	A	99	ARG	12
1	A	35	ASP	12
1	A	27	GLU	10
1	A	59	SER	9
1	A	79	ARG	9
1	A	34	THR	8
1	A	53	PHE	7
1	A	87	SER	7
1	A	58	ASP	7
1	A	43	LYS	6
1	A	61	GLU	6
1	A	81	TYR	5
1	A	86	LYS	4
1	A	82	MET	2
1	A	89	LEU	2
1	A	17	LEU	1
1	A	57	HIS	1
1	A	12	LYS	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 83% for the well-defined parts and 81% for the entire structure.

### 7.1 Chemical shift list 1

File name: 2l9q\_cs.str

Chemical shift list name: *HSP12\_100mMSDS\_110210.bmrB*

#### 7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1103
Number of shifts mapped to atoms	1103
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	107	$-0.51 \pm 0.08$	Should be applied
$^{13}\text{C}_\beta$	95	$-0.01 \pm 0.06$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	102	$-0.35 \pm 0.09$	None needed ( $< 0.5$ ppm)
$^{15}\text{N}$	102	$0.29 \pm 0.23$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 83%, i.e. 599 atoms were assigned a chemical shift out of a possible 722. 8 out of 8 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	306/310 (99%)	123/124 (99%)	122/124 (98%)	61/62 (98%)
Sidechain	293/355 (83%)	183/207 (88%)	110/131 (84%)	0/17 (0%)

*Continued on next page...*



Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	0/57 (0%)	0/30 (0%)	0/26 (0%)	0/1 (0%)
Overall	599/722 (83%)	306/361 (85%)	232/281 (83%)	61/80 (76%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 81%, i.e. 1012 atoms were assigned a chemical shift out of a possible 1257. 10 out of 10 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	520/539 (96%)	209/215 (97%)	209/218 (96%)	102/106 (96%)
Sidechain	492/654 (75%)	312/385 (81%)	180/236 (76%)	0/33 (0%)
Aromatic	0/64 (0%)	0/34 (0%)	0/28 (0%)	0/2 (0%)
Overall	1012/1257 (81%)	521/634 (82%)	389/482 (81%)	102/141 (72%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

