



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 02:42 pm GMT

PDB ID : 4LCD
Title : Structure of an Rsp5xUbxSna3 complex: Mechanism of ubiquitin ligation and lysine prioritization by a HECT E3
Authors : Kamadurai, H.B.; Miller, D.; Schulman, B.A.
Deposited on : 2013-06-21
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1001 (3.12-3.08)
Clashscore	112137	1099 (3.12-3.08)
Ramachandran outliers	110173	1057 (3.12-3.08)
Sidechain outliers	110143	1057 (3.12-3.08)
RSRZ outliers	101464	1006 (3.12-3.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	432	<div> <div>2%</div> <div> <div></div> <div>67%</div> <div>28%</div> <div>• •</div> </div> </div>
1	B	432	<div> <div>3%</div> <div> <div></div> <div>63%</div> <div>30%</div> <div>• •</div> </div> </div>
2	C	24	<div> <div></div> <div> <div>33%</div> <div>67%</div> </div> </div>
2	D	24	<div> <div></div> <div> <div>21%</div> <div>33%</div> <div>•</div> <div>42%</div> </div> </div>
3	E	83	<div> <div>2%</div> <div> <div></div> <div>55%</div> <div>28%</div> <div>7%</div> <div>10%</div> </div> </div>
3	F	83	<div> <div>6%</div> <div> <div></div> <div>47%</div> <div>36%</div> <div>7%</div> <div>10%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7944 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase RSP5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	419	Total	C	N	O	S	0	0	0
			3389	2169	585	624	11			
1	B	413	Total	C	N	O	S	0	0	0
			3289	2110	565	604	10			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	378	GLY	-	EXPRESSION TAG	UNP P39940
A	379	SER	-	EXPRESSION TAG	UNP P39940
A	380	GLY	-	EXPRESSION TAG	UNP P39940
A	381	GLY	-	EXPRESSION TAG	UNP P39940
A	382	SER	-	EXPRESSION TAG	UNP P39940
A	455	LEU	CYS	ENGINEERED MUTATION	UNP P39940
A	517	GLY	CYS	ENGINEERED MUTATION	UNP P39940
A	721	ALA	CYS	ENGINEERED MUTATION	UNP P39940
B	378	GLY	-	EXPRESSION TAG	UNP P39940
B	379	SER	-	EXPRESSION TAG	UNP P39940
B	380	GLY	-	EXPRESSION TAG	UNP P39940
B	381	GLY	-	EXPRESSION TAG	UNP P39940
B	382	SER	-	EXPRESSION TAG	UNP P39940
B	455	LEU	CYS	ENGINEERED MUTATION	UNP P39940
B	517	GLY	CYS	ENGINEERED MUTATION	UNP P39940
B	721	ALA	CYS	ENGINEERED MUTATION	UNP P39940

- Molecule 2 is a protein called Protein SNA3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	8	Total	C	N	O	0	0	0
			51	34	8	9			
2	D	14	Total	C	N	O	0	0	0
			80	51	14	15			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	125	ALA	LYS	ENGINEERED MUTATION	UNP P14359
D	125	ALA	LYS	ENGINEERED MUTATION	UNP P14359

- Molecule 3 is a protein called Ubiquitin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	75	Total	C	N	O	S	0	0	0
			565	353	98	112	2			
3	F	75	Total	C	N	O	S	0	0	0
			570	359	96	113	2			


There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-7	MET	-	EXPRESSION TAG	UNP P0CG48
E	-6	GLY	-	EXPRESSION TAG	UNP P0CG48
E	-5	HIS	-	EXPRESSION TAG	UNP P0CG48
E	-4	HIS	-	EXPRESSION TAG	UNP P0CG48
E	-3	HIS	-	EXPRESSION TAG	UNP P0CG48
E	-2	HIS	-	EXPRESSION TAG	UNP P0CG48
E	-1	HIS	-	EXPRESSION TAG	UNP P0CG48
E	0	HIS	-	EXPRESSION TAG	UNP P0CG48
E	75	CYS	GLY	ENGINEERED MUTATION	UNP P0CG48
F	-7	MET	-	EXPRESSION TAG	UNP P0CG48
F	-6	GLY	-	EXPRESSION TAG	UNP P0CG48
F	-5	HIS	-	EXPRESSION TAG	UNP P0CG48
F	-4	HIS	-	EXPRESSION TAG	UNP P0CG48
F	-3	HIS	-	EXPRESSION TAG	UNP P0CG48
F	-2	HIS	-	EXPRESSION TAG	UNP P0CG48
F	-1	HIS	-	EXPRESSION TAG	UNP P0CG48
F	0	HIS	-	EXPRESSION TAG	UNP P0CG48
F	75	CYS	GLY	ENGINEERED MUTATION	UNP P0CG48


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: E3 ubiquitin-protein ligase RSP5




• Molecule 1: E3 ubiquitin-protein ligase RSP5




• Molecule 2: Protein SNA3

Chain C:  33% 67%



• Molecule 2: Protein SNA3

Chain D:  21% 33% 42%




• Molecule 3: Ubiquitin

Chain E:  2% 55% 28% 7% 10%



• Molecule 3: Ubiquitin

Chain F:  6% 47% 36% 7% 10%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	83.05Å 78.92Å 96.72Å 90.00° 101.67° 90.00°	Depositor
Resolution (Å)	29.40 – 3.10 29.40 – 3.10	Depositor EDS
% Data completeness (in resolution range)	95.0 (29.40-3.10) 91.4 (29.40-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.48 (at 3.11Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.7.3_928)	Depositor
R, R_{free}	0.251 , 0.299 0.250 , 0.300	Depositor DCC
R_{free} test set	1087 reflections (5.10%)	DCC
Wilson B-factor (Å ²)	73.2	Xtriage
Anisotropy	0.488	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 51.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	7944	wwPDB-VP
Average B, all atoms (Å ²)	96.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/3472	0.57	0/4702
1	B	0.47	0/3371	0.59	0/4574
2	C	0.41	0/53	0.76	0/74
2	D	0.28	0/82	0.67	0/114
3	E	0.36	0/571	0.56	0/774
3	F	0.33	0/576	0.56	0/782
All	All	0.45	0/8125	0.58	0/11020

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3389	0	3222	81	0
1	B	3289	0	3067	100	0
2	C	51	0	35	0	0
2	D	80	0	57	9	0
3	E	565	0	551	20	0
3	F	570	0	565	31	0
All	All	7944	0	7497	225	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 225 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:449:ARG:HH11	1:B:449:ARG:CG	1.95	0.78
1:B:599:LEU:HD11	1:B:644:VAL:HG21	1.68	0.75
1:B:408:HIS:HD1	2:D:109:TYR:HH	1.36	0.73
3:F:71:LEU:N	3:F:71:LEU:HD12	2.05	0.71
1:B:449:ARG:HH11	1:B:449:ARG:HG2	1.57	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	415/432 (96%)	390 (94%)	25 (6%)	0	100	100
1	B	407/432 (94%)	366 (90%)	41 (10%)	0	100	100
2	C	6/24 (25%)	2 (33%)	4 (67%)	0	100	100
2	D	12/24 (50%)	6 (50%)	5 (42%)	1 (8%)	1	6
3	E	73/83 (88%)	63 (86%)	10 (14%)	0	100	100
3	F	73/83 (88%)	65 (89%)	8 (11%)	0	100	100
All	All	986/1078 (92%)	892 (90%)	93 (9%)	1 (0%)	55	88

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	111	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	353/386 (92%)	332 (94%)	21 (6%)	23	58
1	B	333/386 (86%)	307 (92%)	26 (8%)	15	48
2	C	3/18 (17%)	3 (100%)	0	100	100
2	D	3/18 (17%)	3 (100%)	0	100	100
3	E	60/76 (79%)	50 (83%)	10 (17%)	2	11
3	F	62/76 (82%)	52 (84%)	10 (16%)	3	12
All	All	814/960 (85%)	747 (92%)	67 (8%)	13	45

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	583	ASP
1	B	723	SER
3	F	48	LYS
1	B	624	VAL
1	B	655	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	668	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	419/432 (96%)	-0.01	9 (2%) 64 43	43, 84, 147, 180	0
1	B	413/432 (95%)	0.11	12 (2%) 52 28	46, 92, 140, 206	0
2	C	8/24 (33%)	0.41	0 100 100	107, 128, 133, 134	0
2	D	14/24 (58%)	0.39	0 100 100	113, 143, 157, 161	0
3	E	75/83 (90%)	0.24	2 (2%) 55 30	84, 124, 183, 204	0
3	F	75/83 (90%)	0.38	5 (6%) 19 7	94, 122, 143, 157	0
All	All	1004/1078 (93%)	0.09	28 (2%) 53 29	43, 96, 149, 206	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	603	ILE	7.5
1	B	589	ALA	5.7
1	A	607	LEU	4.2
3	F	2	GLN	3.8
1	B	406	VAL	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.