



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2017 – 01:09 AM EDT

PDB ID : 3LFT
Title : The Crystal Structure of the ABC domain in complex with L-Trp from *Streptococcus pneumoniae* to 1.35Å
Authors : Stein, A.J.; Mulligan, R.; Abdullah, J.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : unknown
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1024 (1.38-1.34)
Clashscore	112137	1063 (1.38-1.34)
Ramachandran outliers	110173	1048 (1.38-1.34)
Sidechain outliers	110143	1048 (1.38-1.34)
RSRZ outliers	101464	1025 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	<div> <div>9%</div> <div> <div></div> <div>92%</div> <div>8%</div> </div> </div>
1	B	295	<div> <div>30%</div> <div> <div></div> <div>85%</div> <div>10%</div> <div>...</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	TRP	A	1	-	-	-	X
2	TRP	B	2	-	-	-	X

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4868 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	295	Total	C	N	O	Se	0	13	0
			2180	1384	353	436	7			
1	B	288	Total	C	N	O	Se	0	12	0
			2117	1346	339	424	8			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	49	SER	-	expression tag	UNP Q97QX5
A	50	ASN	-	expression tag	UNP Q97QX5
A	51	ALA	-	expression tag	UNP Q97QX5
B	49	SER	-	expression tag	UNP Q97QX5
B	50	ASN	-	expression tag	UNP Q97QX5
B	51	ALA	-	expression tag	UNP Q97QX5

- Molecule 2 is TRYPTOPHAN (three-letter code: TRP) (formula: $C_{11}H_{12}N_2O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			15	11	2	2		
2	B	1	Total	C	N	O	0	0
			15	11	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	347	Total	O	0	0
			347	347		
3	B	194	Total	O	0	0
			194	194		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: uncharacterized protein



- Molecule 1: uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.62Å 68.63Å 137.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.29 – 1.35 36.29 – 1.35	Depositor EDS
% Data completeness (in resolution range)	99.7 (36.29-1.35) 99.6 (36.29-1.35)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.28 (at 1.35Å)	Xtriage
Refinement program	REFMAC refmac_5.5.0102	Depositor
R, R_{free}	0.182 , 0.203 0.216 , 0.230	Depositor DCC
R_{free} test set	5696 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	14.5	Xtriage
Anisotropy	0.162	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 37.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4868	wwPDB-VP
Average B, all atoms (Å ²)	8.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/2243	0.63	1/3046 (0.0%)
1	B	0.75	4/2177 (0.2%)	0.90	8/2960 (0.3%)
All	All	0.62	4/4420 (0.1%)	0.78	9/6006 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	203	LYS	CB-CG	-11.27	1.22	1.52
1	B	294[A]	MSE	N-CA	-10.37	1.25	1.46
1	B	294[B]	MSE	N-CA	-10.37	1.25	1.46
1	B	296	VAL	CB-CG2	-5.45	1.41	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	294[A]	MSE	CA-C-O	-16.95	84.50	120.10
1	B	294[B]	MSE	CA-C-O	-16.95	84.50	120.10
1	B	301	GLY	N-CA-C	11.26	141.25	113.10
1	B	294[A]	MSE	CA-C-N	10.89	141.15	117.20
1	B	294[B]	MSE	CA-C-N	10.89	141.15	117.20
1	B	294[A]	MSE	N-CA-CB	-6.03	99.75	110.60
1	B	294[B]	MSE	N-CA-CB	-6.03	99.75	110.60
1	B	299	LEU	N-CA-C	-5.80	95.34	111.00
1	A	313	SER	C-N-CA	-5.22	108.65	121.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	294[A]	MSE	Mainchain
1	B	294[B]	MSE	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2180	0	2245	23	0
1	B	2117	0	2161	26	1
2	A	15	0	12	0	0
2	B	15	0	9	1	0
3	A	347	0	0	9	0
3	B	194	0	0	0	0
All	All	4868	0	4427	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:314:THR:HG21	1:B:212:THR:OG1	1.29	1.31
1:A:261:LYS:HA	3:A:535:HOH:O	1.59	0.99
1:A:314:THR:HG21	1:B:212:THR:HG1	1.08	0.89
1:B:297:GLN:HG2	1:B:298:VAL:N	1.96	0.79
1:A:314:THR:CG2	1:B:212:THR:OG1	2.22	0.78
1:A:314:THR:CG2	1:B:212:THR:HG1	1.98	0.70
1:A:316:LYS:HE3	3:A:574:HOH:O	1.93	0.68
1:A:73:ASP:OD2	1:A:285:HIS:HE1	1.76	0.68
1:A:330:THR:HG23	3:A:401:HOH:O	1.93	0.67
1:A:221[B]:THR:HG23	3:A:633:HOH:O	1.95	0.66
1:B:297:GLN:O	1:B:300:LYS:HB2	1.95	0.66
1:B:141:ASP:OD2	1:B:197[B]:THR:HG21	1.96	0.66
1:B:157:ASN:HD22	1:B:157:ASN:H	1.44	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:295:ILE:O	1:B:299:LEU:HG	1.99	0.62
1:A:293:LYS:HE3	3:A:571:HOH:O	2.03	0.58
1:A:314:THR:HG23	3:A:589:HOH:O	2.05	0.56
1:B:129[B]:THR:HG22	1:B:131:ASP:H	1.70	0.56
1:B:297:GLN:O	1:B:300:LYS:N	2.29	0.55
1:B:55:VAL:HG22	1:B:115[B]:VAL:HG12	1.90	0.53
1:B:55:VAL:HG22	1:B:115[B]:VAL:CG1	2.40	0.52
1:B:75:LEU:HD12	1:B:296:VAL:HG23	1.91	0.52
1:B:150:ASP:HB3	1:B:153:LYS:O	2.11	0.51
1:B:140:THR:HG1	2:B:2:TRP:N	2.07	0.50
1:B:61:HIS:CE1	1:B:64:LEU:HG	2.47	0.50
1:B:297:GLN:CG	1:B:298:VAL:N	2.55	0.50
1:A:61:HIS:HD2	3:A:9:HOH:O	1.96	0.48
1:A:285:HIS:HD2	3:A:537:HOH:O	1.97	0.48
1:B:143:ILE:HD11	1:B:151:LEU:HG	1.97	0.47
1:B:297:GLN:O	1:B:300:LYS:CB	2.62	0.47
1:A:260[A]:SER:OG	1:A:262[A]:LYS:HG2	2.15	0.46
1:A:61:HIS:HE1	1:A:244:ASN:OD1	1.98	0.46
1:A:184:ILE:HD11	1:A:211:LEU:HD13	1.97	0.46
1:A:96:GLN:HE22	1:A:120:PRO:HB3	1.81	0.45
1:B:289:VAL:O	1:B:293:LYS:HD3	2.17	0.45
1:B:80:TYR:CE2	1:B:296:VAL:HG13	2.53	0.44
1:A:232:THR:HB	1:A:260[A]:SER:OG	2.19	0.42
1:A:260[A]:SER:OG	1:A:262[A]:LYS:HE2	2.18	0.42
1:A:317:SER:OG	1:A:339:ALA:HA	2.19	0.42
1:B:231[A]:MSE:HE2	1:B:238:ILE:HG12	2.02	0.42
1:A:221[B]:THR:CG2	3:A:633:HOH:O	2.60	0.41
1:B:300:LYS:HD3	1:B:300:LYS:N	2.32	0.41
1:A:69[B]:LYS:HA	1:A:72:GLN:HE21	1.85	0.41
1:A:93[A]:GLU:HA	1:A:93[A]:GLU:OE1	2.21	0.41
1:B:119:THR:HA	1:B:139:ILE:HG13	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:223:GLU:OE2	1:B:325[B]:GLN:NE2[3_545]	2.01	0.19

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/295 (104%)	297 (97%)	9 (3%)	0	100	100
1	B	296/295 (100%)	284 (96%)	12 (4%)	0	100	100
All	All	602/590 (102%)	581 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/236 (103%)	242 (99%)	2 (1%)	85	61
1	B	236/236 (100%)	229 (97%)	7 (3%)	46	11
All	All	480/472 (102%)	471 (98%)	9 (2%)	60	25

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81	LYS
1	A	314	THR
1	B	75	LEU
1	B	132	LEU
1	B	157	ASN
1	B	203	LYS
1	B	293	LYS
1	B	300	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	304	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	HIS
1	A	96	GLN
1	A	111	ASN
1	A	146	ASN
1	A	222	ASN
1	A	285	HIS
1	B	157	ASN
1	B	222	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TRP	A	1	-	11,16,16	1.29	1 (9%)	11,22,22	1.43	1 (9%)
2	TRP	B	2	-	11,16,16	1.10	0	11,22,22	1.14	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRP	A	1	-	-	0/3/8/8	0/2/2/2
2	TRP	B	2	-	-	0/3/8/8	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	TRP	CE3-CD2	-2.05	1.38	1.42

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	TRP	CZ3-CE3-CD2	-2.28	117.62	120.88
2	A	1	TRP	CE3-CD2-CE2	3.13	122.32	118.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	TRP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	289/295 (97%)	0.87	28 (9%) 8 12	4, 7, 11, 14	0
1	B	282/295 (95%)	1.64	88 (31%) 0 1	2, 7, 13, 37	0
All	All	571/590 (96%)	1.25	116 (20%) 1 2	2, 7, 11, 37	0

All (116) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	85	VAL	11.0
1	B	80	TYR	9.4
1	B	75	LEU	9.4
1	B	76	ALA	9.3
1	B	296	VAL	8.4
1	B	298	VAL	8.0
1	B	299	LEU	7.7
1	B	155	GLY	7.0
1	B	109	ASN	7.0
1	B	295	ILE	6.6
1	B	87	ILE	6.5
1	B	158	VAL	6.2
1	B	79	GLY	5.5
1	B	71	ILE	5.4
1	B	307	THR	5.3
1	B	149	LYS	5.3
1	B	132	LEU	5.0
1	B	151	LEU	5.0
1	B	156	GLY	5.0
1	B	148	VAL	4.7
1	B	108	ALA	4.7
1	B	305	ALA	4.7
1	B	110	GLY	4.7
1	B	289	VAL	4.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	113	LEU	4.5
1	B	131	ASP	4.5
1	B	135	ILE	4.4
1	B	306	ASP	4.4
1	B	309	VAL	4.3
1	B	300	LYS	4.3
1	A	82	ASP	4.2
1	B	159	THR	4.2
1	B	152	LYS	4.1
1	B	150	ASP	4.1
1	B	77	GLU	4.0
1	B	88	ASP	4.0
1	B	308	PRO	3.9
1	B	133	PRO	3.9
1	A	161	VAL	3.9
1	B	134	VAL	3.9
1	B	112	ASP	3.8
1	B	143	ILE	3.8
1	B	174	ILE	3.8
1	B	292	GLY	3.7
1	B	74	GLY	3.6
1	A	282[A]	ILE	3.5
1	B	160	GLY	3.5
1	B	72	GLN	3.3
1	A	115	VAL	3.3
1	B	154	PRO	3.3
1	B	73	ASP	3.3
1	B	53	ILE	3.2
1	B	147	LEU	3.1
1	B	339	ALA	3.1
1	B	161	VAL	3.1
1	A	135	ILE	3.1
1	B	313	SER	3.1
1	B	78	GLU	3.1
1	B	89	PHE	3.0
1	A	67	ILE	3.0
1	A	309	VAL	3.0
1	B	293	LYS	3.0
1	A	314	THR	3.0
1	B	153	LYS	3.0
1	A	83	ASP	2.9
1	B	311	VAL	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	304	PRO	2.8
1	A	312	PHE	2.8
1	A	287	LEU	2.8
1	A	242	ILE	2.8
1	B	184	ILE	2.8
1	B	290	ALA	2.7
1	B	171	VAL	2.7
1	B	178	THR	2.7
1	A	134	VAL	2.6
1	A	158	VAL	2.6
1	B	105	GLN	2.6
1	B	177	LEU	2.6
1	B	107	VAL	2.6
1	B	173	LEU	2.5
1	B	144	GLY	2.5
1	B	285	HIS	2.5
1	A	330	THR	2.5
1	B	291	THR	2.5
1	B	115[A]	VAL	2.5
1	B	145	ALA	2.5
1	B	340	GLY	2.5
1	B	86	LYS	2.5
1	B	114	VAL	2.5
1	B	297	GLN	2.5
1	A	64	LEU	2.4
1	A	311	VAL	2.4
1	A	81	LYS	2.3
1	B	314	THR	2.3
1	B	302	ALA	2.3
1	B	104	LYS	2.3
1	A	295	ILE	2.3
1	A	329	ILE	2.3
1	B	100	ALA	2.3
1	A	140	THR	2.2
1	B	265	TYR	2.2
1	A	117	ILE	2.2
1	A	139	ILE	2.2
1	B	146	ASN	2.1
1	B	301	GLY	2.1
1	A	71	ILE	2.1
1	B	239	TRP	2.1
1	B	70	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	205	TYR	2.1
1	A	138	ALA	2.1
1	A	151	LEU	2.1
1	A	116	GLY	2.1
1	B	66	LEU	2.1
1	B	264	ILE	2.1
1	B	343	ILE	2.1
1	B	101	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	TRP	B	2	15/15	0.91	0.18	5.01	11,12,13,14	0
2	TRP	A	1	15/15	0.96	0.18	2.16	7,8,8,9	0

6.5 Other polymers [i](#)

There are no such residues in this entry.