



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 06:07 am GMT

PDB ID : 4LF9  
Title : Crystal Structure of 30S ribosomal subunit from *Thermus thermophilus*  
Authors : Demirci, H.; Belardinelli, R.; Carr, J.; Murphy IV, F.; Jögl, G.; Dahlberg, A.E.; Gregory, S.T.  
Deposited on : 2013-06-26  
Resolution : 3.28 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




















Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1006 (3.34-3.22)
Clashscore	112137	1070 (3.34-3.22)
Ramachandran outliers	110173	1050 (3.34-3.22)
Sidechain outliers	110143	1049 (3.34-3.22)
RSRZ outliers	101464	1011 (3.34-3.22)
RNA backbone	2435	1089 (3.76-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1522	<div> <div>71%</div> <div>24%</div> <div>• •</div> </div>
2	B	256	<div> <div>73%</div> <div>19%</div> <div>8%</div> </div>
3	C	239	<div> <div>66%</div> <div>19%</div> <div>•</div> <div>13%</div> </div>
4	D	209	<div> <div>77%</div> <div>19%</div> <div>•</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	162	
6	F	101	
7	G	156	
8	H	138	
9	I	128	
10	J	105	
11	K	129	
12	L	135	
13	M	126	
14	N	61	
15	O	89	
16	P	88	
17	Q	105	
18	R	88	
19	S	93	
20	T	106	
21	U	27	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
22	MG	A	1612	-	-	-	X
22	MG	A	1618	-	-	-	X
22	MG	A	1623	-	-	-	X
22	MG	A	1626	-	-	-	X
22	MG	A	1628	-	-	-	X
22	MG	A	1629	-	-	-	X
22	MG	A	1635	-	-	-	X
22	MG	A	1640	-	-	-	X

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
22	MG	A	1646	-	-	-	X
22	MG	A	1647	-	-	-	X
22	MG	A	1651	-	-	-	X
22	MG	A	1660	-	-	-	X
22	MG	A	1663	-	-	-	X
22	MG	A	1664	-	-	-	X
22	MG	A	1672	-	-	-	X
22	MG	A	1673	-	-	-	X
22	MG	A	1674	-	-	-	X
22	MG	A	1675	-	-	-	X
22	MG	A	1679	-	-	-	X
22	MG	A	1682	-	-	-	X
22	MG	A	1688	-	-	-	X
22	MG	A	1689	-	-	-	X
22	MG	A	1691	-	-	-	X
22	MG	A	1693	-	-	-	X
22	MG	A	1694	-	-	-	X
22	MG	A	1696	-	-	-	X
22	MG	A	1700	-	-	-	X
22	MG	A	1702	-	-	-	X
22	MG	A	1704	-	-	-	X
22	MG	A	1706	-	-	-	X
22	MG	A	1708	-	-	-	X
22	MG	A	1711	-	-	-	X
22	MG	A	1729	-	-	-	X
22	MG	A	1736	-	-	-	X
23	K	A	1712	-	-	-	X

## 2 Entry composition

There are 26 unique types of molecules in this entry. The entry contains 51883 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 16S rRNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1512	Total	C	N	O	P	0	0	0
			32504	14477	6011	10505	1511			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1534	C	A	CONFLICT	GB M26923.1
A	1535	A	C	CONFLICT	GB M26923.1

- Molecule 2 is a protein called ribosomal protein S2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	236	Total	C	N	O	S	0	0	1
			1874	1195	336	338	5			

- Molecule 3 is a protein called ribosomal protein S3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	207	Total	C	N	O	S	0	0	1
			1613	1016	315	281	1			

- Molecule 4 is a protein called ribosomal protein S4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	208	Total	C	N	O	S	0	0	0
			1703	1066	339	291	7			

- Molecule 5 is a protein called ribosomal protein S5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	151	Total	C	N	O	S	0	0	1
			1147	724	218	201	4			

- Molecule 6 is a protein called ribosomal protein S6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	101	Total	C	N	O	S	0	0	0
			843	531	155	154	3			

- Molecule 7 is a protein called ribosomal protein S7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	155	Total	C	N	O	S	0	0	0
			1257	781	252	218	6			

- Molecule 8 is a protein called ribosomal protein S8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	138	Total	C	N	O	S	0	0	0
			1116	705	215	193	3			

- Molecule 9 is a protein called ribosomal protein S9.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
9	I	127	Total	C	N	O	0	0	0
			1010	639	197	174			

- Molecule 10 is a protein called ribosomal protein S10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	99	Total	C	N	O	S	0	0	1
			793	498	157	137	1			

- Molecule 11 is a protein called ribosomal protein S11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	119	Total	C	N	O	S	0	0	0
			885	549	168	165	3			

- Molecule 12 is a protein called ribosomal protein S12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	125	Total	C	N	O	S	0	0	1
			973	612	196	163	2			

- Molecule 13 is a protein called ribosomal protein S13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	118	Total	C	N	O	S	0	0	0
			937	579	193	163	2			

- Molecule 14 is a protein called ribosomal protein S14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	60	Total	C	N	O	S	0	0	0
			492	312	104	72	4			

- Molecule 15 is a protein called ribosomal protein S15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
15	O	88	Total	C	N	O	S	0	0	0
			734	459	147	126	2			

- Molecule 16 is a protein called ribosomal protein S16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
16	P	84	Total	C	N	O	S	0	0	1
			701	443	140	117	1			

- Molecule 17 is a protein called ribosomal protein S17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
17	Q	104	Total	C	N	O	S	0	0	0
			857	547	160	148	2			

- Molecule 18 is a protein called ribosomal protein S18.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
18	R	73	Total	C	N	O	0	0	0
			598	381	118	99			

- Molecule 19 is a protein called ribosomal protein S19.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
19	S	81	Total	C	N	O	S	0	0	1
			648	414	120	112	2			

- Molecule 20 is a protein called ribosomal protein S20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
20	T	99	Total	C	N	O	S	0	0	0
			763	470	162	129	2			

- Molecule 21 is a protein called ribosomal protein THX.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
21	U	25	Total	C	N	O	0	0	1
			209	128	51	30			

- Molecule 22 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
22	P	1	Total	Mg	0	0
			1	1		
22	D	1	Total	Mg	0	0
			1	1		
22	E	1	Total	Mg	0	0
			1	1		
22	B	1	Total	Mg	0	0
			1	1		
22	A	134	Total	Mg	0	0
			134	134		
22	L	1	Total	Mg	0	0
			1	1		
22	S	2	Total	Mg	0	0
			2	2		
22	F	1	Total	Mg	0	0
			1	1		
22	M	1	Total	Mg	0	0
			1	1		

- Molecule 23 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
23	A	8	Total	K	0	0
			8	8		

- Molecule 24 is (2R,3R,4R,5R)-2-((1S,2S,3R,4S,6R)-4,6-DIAMINO-3-((2R,3R,6S)-3-AMINO-6-(AMINOMETHYL)-TETRAHYDRO-2H-PYRAN-2-YLOXY)-2-HYDROXYCYCLOHEXYLOXY)-5-METHYL-4-(METHYLAMINO)-TETRAHYDRO-2H-PYRAN-3,5-DIOL (three-letter code: LLL) (formula: C<sub>19</sub>H<sub>39</sub>N<sub>5</sub>O<sub>7</sub>).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
24	A	1	Total	C	N	O	0	0
			31	19	5	7		


- Molecule 25 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
25	D	1	Total	Zn	0	0
			1	1		
25	N	1	Total	Zn	0	0
			1	1		



- Molecule 26 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
26	A	39	Total	O	0	0
			39	39		
26	S	3	Total	O	0	0
			3	3		








• Molecule 2: ribosomal protein S2




• Molecule 3: ribosomal protein S3




• Molecule 4: ribosomal protein S4




• Molecule 5: ribosomal protein S5




- Molecule 6: ribosomal protein S6




- Molecule 7: ribosomal protein S7




- Molecule 8: ribosomal protein S8




- Molecule 9: ribosomal protein S9




- Molecule 10: ribosomal protein S10




- Molecule 11: ribosomal protein S11



- Molecule 12: ribosomal protein S12




- Molecule 13: ribosomal protein S13





- Molecule 14: ribosomal protein S14





- Molecule 15: ribosomal protein S15



- Molecule 16: ribosomal protein S16




- Molecule 17: ribosomal protein S17



- Molecule 18: ribosomal protein S18







- Molecule 19: ribosomal protein S19



- Molecule 20: ribosomal protein S20



- Molecule 21: ribosomal protein THX



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	402.04Å 402.04Å 176.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.48 – 3.28 34.47 – 3.28	Depositor EDS
% Data completeness (in resolution range)	99.1 (34.48-3.28) 98.8 (34.47-3.28)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.04 (at 3.25Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1119)	Depositor
R, $R_{free}$	0.169 , 0.201 0.170 , 0.203	Depositor DCC
$R_{free}$ test set	10925 reflections (5.05%)	DCC
Wilson B-factor (Å <sup>2</sup> )	115.7	Xtriage
Anisotropy	0.203	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.25 , 91.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	51883	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	117.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MA6, K, 0TD, ZN, LLL, 2MG, 5MC, UR3, 4OC, M2G, 7MG, PSU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/36037	0.82	35/56239 (0.1%)
2	B	0.27	0/1909	0.45	0/2579
3	C	0.26	0/1637	0.45	0/2207
4	D	0.32	0/1733	0.47	0/2318
5	E	0.35	0/1163	0.55	0/1566
6	F	0.25	0/856	0.42	0/1154
7	G	0.27	0/1276	0.45	0/1709
8	H	0.33	0/1136	0.50	0/1527
9	I	0.26	0/1029	0.49	0/1379
10	J	0.27	0/806	0.51	0/1084
11	K	0.31	0/900	0.53	0/1213
12	L	0.31	0/978	0.55	0/1308
13	M	0.25	0/947	0.47	0/1270
14	N	0.29	0/501	0.47	0/664
15	O	0.28	0/745	0.47	0/992
16	P	0.30	0/717	0.50	0/965
17	Q	0.32	0/870	0.54	0/1159
18	R	0.28	0/604	0.46	0/801
19	S	0.23	0/662	0.46	0/892
20	T	0.31	0/765	0.52	0/1007
21	U	0.26	0/213	0.45	0/279
All	All	0.34	0/55484	0.73	35/82312 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
9	I	0	1



There are no bond length outliers.

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	328	C	C2-N1-C1'	8.33	127.96	118.80
1	A	1054	C	C2-N1-C1'	7.77	127.35	118.80
1	A	839	U	C2-N1-C1'	7.48	126.67	117.70
1	A	1054	C	N1-C2-O2	7.41	123.35	118.90
1	A	839	U	N1-C2-O2	7.11	127.78	122.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
9	I	38	GLN	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	32504	0	16434	220	0
2	B	1874	0	1887	25	0
3	C	1613	0	1677	33	0
4	D	1703	0	1763	31	0
5	E	1147	0	1207	22	0
6	F	843	0	857	9	0
7	G	1257	0	1296	16	0
8	H	1116	0	1177	17	0
9	I	1010	0	1037	24	0
10	J	793	0	835	26	0
11	K	885	0	904	14	0
12	L	973	0	1058	11	0
13	M	937	0	995	22	0
14	N	492	0	529	11	0
15	O	734	0	771	13	0
16	P	701	0	720	5	0
17	Q	857	0	928	16	0
18	R	598	0	670	8	0
19	S	648	0	673	14	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
20	T	763	0	861	14	0
21	U	209	0	221	2	0
22	A	134	0	0	0	0
22	B	1	0	0	0	0
22	D	1	0	0	0	0
22	E	1	0	0	0	0
22	F	1	0	0	0	0
22	L	1	0	0	0	0
22	M	1	0	0	0	0
22	P	1	0	0	0	0
22	S	2	0	0	0	0
23	A	8	0	0	0	0
24	A	31	0	39	2	0
25	D	1	0	0	0	0
25	N	1	0	0	0	0
26	A	39	0	0	0	0
26	S	3	0	0	0	0
All	All	51883	0	36539	474	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 474 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
19:S:33:THR:HG22	19:S:35:SER:H	1.42	0.85
1:A:1502:A:H2	1:A:1505:G:H1	1.25	0.85
9:I:10:ARG:HD3	9:I:105:ASP:HB3	1.62	0.82
1:A:235:C:H5'	17:Q:70:ARG:HG2	1.62	0.81
1:A:279:A:OP2	17:Q:95:TYR:OH	1.99	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	234/256 (91%)	218 (93%)	15 (6%)	1 (0%)	38	72
3	C	205/239 (86%)	190 (93%)	15 (7%)	0	100	100
4	D	206/209 (99%)	197 (96%)	8 (4%)	1 (0%)	32	67
5	E	149/162 (92%)	145 (97%)	4 (3%)	0	100	100
6	F	99/101 (98%)	95 (96%)	4 (4%)	0	100	100
7	G	153/156 (98%)	149 (97%)	4 (3%)	0	100	100
8	H	136/138 (99%)	134 (98%)	2 (2%)	0	100	100
9	I	125/128 (98%)	114 (91%)	11 (9%)	0	100	100
10	J	97/105 (92%)	81 (84%)	14 (14%)	2 (2%)	8	38
11	K	117/129 (91%)	106 (91%)	11 (9%)	0	100	100
12	L	122/135 (90%)	112 (92%)	9 (7%)	1 (1%)	22	58
13	M	116/126 (92%)	109 (94%)	7 (6%)	0	100	100
14	N	58/61 (95%)	55 (95%)	3 (5%)	0	100	100
15	O	86/89 (97%)	81 (94%)	5 (6%)	0	100	100
16	P	82/88 (93%)	80 (98%)	2 (2%)	0	100	100
17	Q	102/105 (97%)	97 (95%)	5 (5%)	0	100	100
18	R	71/88 (81%)	66 (93%)	5 (7%)	0	100	100
19	S	79/93 (85%)	70 (89%)	8 (10%)	1 (1%)	14	48
20	T	97/106 (92%)	90 (93%)	7 (7%)	0	100	100
21	U	23/27 (85%)	22 (96%)	1 (4%)	0	100	100
All	All	2357/2541 (93%)	2211 (94%)	140 (6%)	6 (0%)	44	77

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
12	L	28	LYS
19	S	31	ILE
10	J	72	VAL
10	J	34	VAL
2	B	229	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	194/220 (88%)	185 (95%)	9 (5%)	31	66
3	C	160/188 (85%)	148 (92%)	12 (8%)	16	47
4	D	180/181 (99%)	168 (93%)	12 (7%)	19	53
5	E	115/123 (94%)	106 (92%)	9 (8%)	15	46
6	F	90/90 (100%)	88 (98%)	2 (2%)	57	80
7	G	126/127 (99%)	123 (98%)	3 (2%)	54	78
8	H	119/119 (100%)	112 (94%)	7 (6%)	23	58
9	I	98/99 (99%)	92 (94%)	6 (6%)	22	57
10	J	87/92 (95%)	83 (95%)	4 (5%)	31	66
11	K	90/99 (91%)	88 (98%)	2 (2%)	57	80
12	L	103/110 (94%)	100 (97%)	3 (3%)	48	75
13	M	94/101 (93%)	87 (93%)	7 (7%)	16	48
14	N	49/50 (98%)	46 (94%)	3 (6%)	22	57
15	O	79/80 (99%)	73 (92%)	6 (8%)	15	47
16	P	72/74 (97%)	70 (97%)	2 (3%)	49	76
17	Q	96/97 (99%)	89 (93%)	7 (7%)	16	49
18	R	64/77 (83%)	63 (98%)	1 (2%)	68	84
19	S	71/80 (89%)	68 (96%)	3 (4%)	34	68
20	T	76/82 (93%)	69 (91%)	7 (9%)	11	37
21	U	19/22 (86%)	19 (100%)	0	100	100
All	All	1982/2111 (94%)	1877 (95%)	105 (5%)	26	62

5 of 105 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
8	H	33	GLU
9	I	121	ARG
19	S	81	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
8	H	39	LEU
8	H	133	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
9	I	73	GLN
9	I	124	GLN
13	M	106	ASN
9	I	3	GLN
10	J	13	HIS

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	1507/1522 (99%)	210 (13%)	0

5 of 210 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	9	G
1	A	32	A
1	A	39	G
1	A	44	G
1	A	47	C

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

15 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	2MG	A	1207	1	19,26,27	2.09	3 (15%)	20,38,41	2.01	3 (15%)
1	5MC	A	1400	1	15,22,23	0.92	0	17,32,35	0.94	1 (5%)
1	4OC	A	1402	1	16,23,24	1.03	2 (12%)	19,32,35	0.60	0
1	5MC	A	1404	1	15,22,23	1.00	1 (6%)	17,32,35	1.03	2 (11%)
1	5MC	A	1407	1	15,22,23	1.02	0	17,32,35	0.93	0
1	UR3	A	1498	1	14,22,23	0.74	0	16,32,35	1.13	0
1	MA6	A	1518	1	16,26,27	0.62	0	18,38,41	1.09	1 (5%)
1	MA6	A	1519	1	16,26,27	0.89	1 (6%)	18,38,41	1.09	1 (5%)
1	PSU	A	1540	1	16,21,22	1.05	1 (6%)	20,30,33	3.53	6 (30%)
1	PSU	A	1541	1	16,21,22	1.12	3 (18%)	20,30,33	3.51	6 (30%)
1	PSU	A	516	1,22	16,21,22	1.06	1 (6%)	20,30,33	3.58	6 (30%)
1	7MG	A	527	1	20,26,27	2.70	7 (35%)	22,39,42	1.71	6 (27%)
1	M2G	A	966	1	20,27,28	1.81	4 (20%)	21,40,43	2.31	3 (14%)
1	5MC	A	967	1	15,22,23	0.93	0	17,32,35	0.87	0
12	0TD	L	92	12	5,9,10	1.95	1 (20%)	3,11,13	2.51	2 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	2MG	A	1207	1	-	0/5/27/28	0/3/3/3
1	5MC	A	1400	1	-	0/3/25/26	0/2/2/2
1	4OC	A	1402	1	-	0/7/29/30	0/2/2/2
1	5MC	A	1404	1	-	0/3/25/26	0/2/2/2
1	5MC	A	1407	1	-	0/3/25/26	0/2/2/2
1	UR3	A	1498	1	-	0/3/25/26	0/2/2/2
1	MA6	A	1518	1	-	0/7/29/30	0/3/3/3
1	MA6	A	1519	1	-	0/7/29/30	0/3/3/3
1	PSU	A	1540	1	-	0/7/25/26	0/2/2/2
1	PSU	A	1541	1	-	0/7/25/26	0/2/2/2
1	PSU	A	516	1,22	-	0/7/25/26	0/2/2/2
1	7MG	A	527	1	-	0/7/37/38	0/3/3/3
1	M2G	A	966	1	-	0/7/29/30	0/3/3/3
1	5MC	A	967	1	-	0/3/25/26	0/2/2/2
12	0TD	L	92	12	-	0/2/12/14	0/0/0/0

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	527	7MG	C8-N9	-7.68	1.34	1.45
1	A	527	7MG	O5'-C5'	-2.16	1.41	1.44
1	A	1541	PSU	C5-C1'	-2.05	1.50	1.52
1	A	527	7MG	CM7-N7	-2.04	1.42	1.46
1	A	1541	PSU	O4'-C1'	-2.02	1.41	1.44

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	516	PSU	N1-C2-N3	-12.28	119.57	128.40
1	A	1540	PSU	N1-C2-N3	-11.87	119.86	128.40
1	A	1541	PSU	N1-C2-N3	-11.62	120.04	128.40
1	A	966	M2G	C5-C6-N1	-7.75	112.44	123.48
1	A	1207	2MG	C5-C6-N1	-7.14	113.31	123.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1207	2MG	1	0
1	A	1498	UR3	1	0
1	A	1518	MA6	1	0
1	A	1519	MA6	1	0
1	A	527	7MG	1	0
12	L	92	0TD	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 154 ligands modelled in this entry, 153 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
24	LLL	A	1743	-	30,33,33	1.18	2 (6%)	33,49,49	1.25	6 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
24	LLL	A	1743	-	-	0/11/65/65	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
24	A	1743	LLL	O43-C43	-3.27	1.38	1.44
24	A	1743	LLL	O23-C23	-2.33	1.37	1.43

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
24	A	1743	LLL	O62-C13-O53	-2.79	103.27	108.98
24	A	1743	LLL	C41-C51-C61	-2.14	108.58	112.83
24	A	1743	LLL	C11-O51-C51	2.04	115.38	113.09
24	A	1743	LLL	O43-C43-C83	2.10	112.68	108.10
24	A	1743	LLL	C62-C52-C42	2.47	114.17	109.01

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
24	A	1743	LLL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.



## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1498/1522 (98%)	-0.14	22 (1%) 74 70	65, 101, 191, 299	0
2	B	236/256 (92%)	-0.11	5 (2%) 64 60	82, 134, 223, 254	0
3	C	207/239 (86%)	-0.00	9 (4%) 36 33	79, 138, 182, 217	0
4	D	208/209 (99%)	0.00	8 (3%) 41 36	78, 114, 158, 180	0
5	E	151/162 (93%)	-0.18	2 (1%) 77 75	72, 95, 135, 193	0
6	F	101/101 (100%)	-0.27	1 (0%) 82 81	101, 134, 163, 221	0
7	G	155/156 (99%)	-0.12	4 (2%) 56 53	92, 123, 183, 214	0
8	H	138/138 (100%)	-0.22	1 (0%) 87 87	65, 93, 126, 157	0
9	I	127/128 (99%)	0.01	1 (0%) 86 85	99, 145, 180, 203	0
10	J	99/105 (94%)	0.36	4 (4%) 39 35	83, 170, 250, 276	0
11	K	119/129 (92%)	0.04	2 (1%) 70 67	75, 106, 157, 176	0
12	L	124/135 (91%)	-0.04	1 (0%) 86 85	65, 104, 145, 255	0
13	M	118/126 (93%)	0.01	5 (4%) 37 34	96, 130, 166, 233	0
14	N	60/61 (98%)	0.22	3 (5%) 30 26	99, 124, 167, 237	0
15	O	88/89 (98%)	0.05	4 (4%) 34 31	71, 108, 151, 184	0
16	P	84/88 (95%)	-0.11	1 (1%) 79 76	80, 104, 139, 197	0
17	Q	104/105 (99%)	0.35	4 (3%) 41 36	74, 101, 157, 216	0
18	R	73/88 (82%)	0.03	3 (4%) 38 34	84, 115, 180, 227	0
19	S	81/93 (87%)	-0.01	1 (1%) 79 76	65, 152, 211, 246	0
20	T	99/106 (93%)	-0.15	2 (2%) 65 62	78, 105, 162, 249	0
21	U	25/27 (92%)	0.86	5 (20%) 1 1	76, 132, 150, 204	0
All	All	3895/4063 (95%)	-0.06	88 (2%) 61 58	65, 114, 189, 299	0

The worst 5 of 88 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
11	K	128	ALA	15.3
17	Q	104	LYS	15.2
11	K	129	SER	14.6
17	Q	103	GLY	11.8
17	Q	105	ALA	10.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	M2G	A	966	25/26	0.98	0.13	-	91,109,117,121	0
1	PSU	A	1540	20/21	0.86	0.33	-	167,192,205,206	0
1	UR3	A	1498	21/22	0.95	0.21	-	72,79,89,101	0
1	4OC	A	1402	22/23	0.98	0.18	-	80,85,94,96	0
1	7MG	A	527	24/25	0.95	0.18	-	83,90,102,105	0
1	5MC	A	1407	21/22	0.97	0.21	-	86,92,103,105	0
1	5MC	A	967	21/22	0.96	0.12	-	89,96,102,107	0
1	PSU	A	1541	20/21	0.90	0.24	-	171,183,195,197	0
12	0TD	L	92	10/11	0.93	0.44	-	98,106,163,288	0
1	MA6	A	1518	24/25	0.98	0.18	-	77,84,94,100	0
1	5MC	A	1404	21/22	0.97	0.16	-	75,78,103,108	0
1	2MG	A	1207	24/25	0.96	0.14	-	125,128,130,132	0
1	MA6	A	1519	24/25	0.97	0.20	-	72,79,87,91	0
1	5MC	A	1400	21/22	0.96	0.20	-	79,91,106,109	0
1	PSU	A	516	20/21	0.92	0.14	-	118,127,134,137	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
22	MG	A	1674	1/1	0.93	2.14	47.16	91,91,91,91	0
22	MG	A	1612	1/1	0.76	1.36	42.24	80,80,80,80	0
22	MG	A	1691	1/1	0.98	1.24	41.93	74,74,74,74	0
22	MG	A	1618	1/1	0.97	0.81	35.57	58,58,58,58	0
22	MG	A	1736	1/1	0.95	1.03	35.08	116,116,116,116	0
22	MG	A	1682	1/1	0.92	0.63	34.11	64,64,64,64	0
22	MG	A	1623	1/1	0.95	0.64	26.36	68,68,68,68	0
22	MG	A	1675	1/1	0.98	0.70	24.03	58,58,58,58	0
22	MG	A	1711	1/1	0.77	1.37	23.29	106,106,106,106	0
22	MG	A	1702	1/1	0.87	1.33	23.21	95,95,95,95	0
22	MG	A	1664	1/1	0.89	0.78	21.04	69,69,69,69	0
22	MG	A	1660	1/1	0.97	0.66	19.17	72,72,72,72	0
22	MG	A	1640	1/1	0.96	0.68	18.42	73,73,73,73	0
22	MG	A	1663	1/1	0.98	0.45	17.31	83,83,83,83	0
22	MG	A	1700	1/1	0.96	0.47	16.18	70,70,70,70	0
22	MG	A	1628	1/1	0.90	0.38	14.36	58,58,58,58	0
22	MG	A	1635	1/1	0.91	0.35	13.31	66,66,66,66	0
22	MG	A	1651	1/1	0.95	0.42	12.86	89,89,89,89	0
22	MG	A	1688	1/1	0.93	0.33	11.92	56,56,56,56	0
22	MG	A	1672	1/1	0.91	0.48	11.50	67,67,67,67	0
22	MG	A	1729	1/1	0.92	0.42	10.19	79,79,79,79	0
22	MG	A	1694	1/1	0.99	0.80	10.13	76,76,76,76	0
23	K	A	1712	1/1	0.88	0.32	9.06	124,124,124,124	0
22	MG	A	1647	1/1	0.98	0.47	7.96	59,59,59,59	0
22	MG	A	1706	1/1	0.91	0.45	7.27	72,72,72,72	0
22	MG	A	1708	1/1	0.89	0.33	6.72	72,72,72,72	0
22	MG	A	1629	1/1	0.98	0.38	6.15	82,82,82,82	0
22	MG	A	1693	1/1	0.94	0.40	6.12	86,86,86,86	0
22	MG	A	1646	1/1	0.81	0.93	5.69	82,82,82,82	0
22	MG	A	1626	1/1	0.87	0.45	5.04	100,100,100,100	0
22	MG	A	1679	1/1	0.96	0.26	3.72	70,70,70,70	0
22	MG	A	1673	1/1	0.84	0.33	3.41	59,59,59,59	0
22	MG	A	1689	1/1	0.95	0.25	3.28	65,65,65,65	0
22	MG	A	1704	1/1	0.91	0.19	2.71	75,75,75,75	0
22	MG	M	201	1/1	0.96	0.31	1.93	93,93,93,93	0
22	MG	A	1696	1/1	0.96	0.41	1.78	108,108,108,108	0
22	MG	A	1683	1/1	0.97	0.22	1.45	59,59,59,59	0
24	LLL	A	1743	31/31	0.96	0.21	1.23	64,79,103,105	0
25	ZN	N	101	1/1	1.00	0.21	0.98	135,135,135,135	0
22	MG	A	1709	1/1	0.95	0.31	0.54	81,81,81,81	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
22	MG	A	1630	1/1	0.92	0.17	-0.17	86,86,86,86	0
22	MG	A	1643	1/1	0.97	0.19	-0.24	59,59,59,59	0
22	MG	A	1607	1/1	0.97	0.19	-0.36	91,91,91,91	0
22	MG	A	1705	1/1	0.97	0.15	-0.37	55,55,55,55	0
22	MG	D	302	1/1	0.89	0.14	-0.55	89,89,89,89	0
22	MG	A	1701	1/1	0.94	0.14	-0.72	86,86,86,86	0
22	MG	A	1665	1/1	0.96	0.13	-0.81	64,64,64,64	0
22	MG	A	1728	1/1	0.91	0.13	-0.87	75,75,75,75	0
22	MG	A	1732	1/1	0.95	0.12	-1.27	71,71,71,71	0
22	MG	A	1685	1/1	0.97	0.11	-1.43	75,75,75,75	0
22	MG	A	1636	1/1	0.78	1.00	-	78,78,78,78	0
22	MG	A	1638	1/1	0.96	0.59	-	70,70,70,70	0
22	MG	A	1686	1/1	0.99	0.23	-	66,66,66,66	0
23	K	A	1717	1/1	0.74	0.91	-	151,151,151,151	0
22	MG	A	1631	1/1	0.99	0.36	-	10,10,10,10	0
23	K	A	1715	1/1	0.94	0.28	-	150,150,150,150	0
22	MG	A	1680	1/1	0.85	1.27	-	81,81,81,81	0
22	MG	A	1669	1/1	0.76	0.48	-	98,98,98,98	0
22	MG	A	1667	1/1	0.73	0.53	-	72,72,72,72	0
23	K	A	1716	1/1	0.88	0.40	-	158,158,158,158	0
22	MG	A	1737	1/1	0.77	0.29	-	103,103,103,103	0
22	MG	A	1610	1/1	0.79	1.54	-	113,113,113,113	0
22	MG	A	1621	1/1	0.88	0.41	-	97,97,97,97	0
22	MG	A	1722	1/1	0.90	0.33	-	323,323,323,323	0
22	MG	A	1698	1/1	0.98	0.51	-	120,120,120,120	0
22	MG	A	1658	1/1	0.91	0.53	-	89,89,89,89	0
22	MG	A	1727	1/1	0.91	0.21	-	77,77,77,77	0
22	MG	A	1627	1/1	0.98	0.34	-	57,57,57,57	0
22	MG	A	1703	1/1	0.93	0.57	-	93,93,93,93	0
22	MG	A	1624	1/1	0.88	0.53	-	60,60,60,60	0
22	MG	A	1735	1/1	0.91	0.38	-	99,99,99,99	0
22	MG	A	1625	1/1	0.93	0.50	-	82,82,82,82	0
22	MG	A	1605	1/1	0.96	0.50	-	28,28,28,28	0
22	MG	A	1611	1/1	0.69	1.08	-	94,94,94,94	0
22	MG	A	1725	1/1	0.94	0.38	-	322,322,322,322	0
22	MG	A	1652	1/1	0.94	0.58	-	66,66,66,66	0
22	MG	A	1741	1/1	0.41	0.27	-	102,102,102,102	0
22	MG	E	201	1/1	0.90	0.39	-	92,92,92,92	0
22	MG	A	1684	1/1	0.94	0.72	-	64,64,64,64	0
22	MG	A	1632	1/1	0.99	0.40	-	54,54,54,54	0
22	MG	A	1617	1/1	0.93	1.12	-	88,88,88,88	0
22	MG	A	1649	1/1	0.73	0.69	-	86,86,86,86	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
22	MG	A	1609	1/1	0.83	0.84	-	94,94,94,94	0
22	MG	F	201	1/1	0.49	0.78	-	136,136,136,136	0
22	MG	A	1710	1/1	0.98	0.12	-	75,75,75,75	0
22	MG	A	1730	1/1	0.83	0.91	-	75,75,75,75	0
22	MG	A	1656	1/1	0.93	0.66	-	73,73,73,73	0
22	MG	A	1659	1/1	0.96	0.18	-	67,67,67,67	0
22	MG	A	1740	1/1	0.92	0.89	-	88,88,88,88	0
22	MG	A	1692	1/1	0.96	1.48	-	87,87,87,87	0
22	MG	A	1695	1/1	0.98	0.15	-	79,79,79,79	0
22	MG	A	1613	1/1	0.71	0.47	-	105,105,105,105	0
22	MG	A	1687	1/1	0.94	0.37	-	67,67,67,67	0
23	K	A	1718	1/1	0.94	0.21	-	165,165,165,165	0
22	MG	A	1616	1/1	0.94	0.29	-	88,88,88,88	0
22	MG	A	1657	1/1	0.56	0.90	-	99,99,99,99	0
22	MG	A	1614	1/1	0.92	1.45	-	65,65,65,65	0
22	MG	A	1726	1/1	0.55	0.96	-	110,110,110,110	0
22	MG	A	1648	1/1	0.91	0.28	-	88,88,88,88	0
22	MG	A	1668	1/1	0.93	0.83	-	72,72,72,72	0
22	MG	A	1641	1/1	0.86	1.13	-	81,81,81,81	0
22	MG	A	1666	1/1	0.97	0.32	-	52,52,52,52	0
22	MG	A	1642	1/1	0.97	0.57	-	67,67,67,67	0
22	MG	A	1644	1/1	0.90	0.80	-	83,83,83,83	0
22	MG	A	1645	1/1	0.94	0.79	-	90,90,90,90	0
23	K	A	1713	1/1	0.96	0.48	-	123,123,123,123	0
22	MG	A	1601	1/1	0.95	0.17	-	99,99,99,99	0
22	MG	A	1604	1/1	0.86	1.46	-	79,79,79,79	0
22	MG	A	1608	1/1	0.73	0.34	-	68,68,68,68	0
23	K	A	1714	1/1	0.91	0.17	-	152,152,152,152	0
22	MG	S	101	1/1	0.96	0.44	-	282,282,282,282	0
22	MG	A	1723	1/1	0.93	0.12	-	300,300,300,300	0
22	MG	A	1603	1/1	0.96	0.98	-	51,51,51,51	0
22	MG	A	1720	1/1	0.92	0.11	-	167,167,167,167	0
22	MG	A	1677	1/1	0.84	0.73	-	98,98,98,98	0
22	MG	A	1738	1/1	0.39	0.36	-	107,107,107,107	0
22	MG	A	1639	1/1	0.94	1.02	-	88,88,88,88	0
22	MG	A	1742	1/1	0.76	1.00	-	84,84,84,84	0
22	MG	A	1724	1/1	0.91	0.07	-	238,238,238,238	0
22	MG	A	1606	1/1	0.79	1.37	-	110,110,110,110	0
22	MG	A	1634	1/1	0.74	0.70	-	94,94,94,94	0
22	MG	A	1653	1/1	0.95	0.27	-	86,86,86,86	0
22	MG	A	1670	1/1	0.81	0.39	-	91,91,91,91	0
22	MG	A	1681	1/1	0.90	0.39	-	90,90,90,90	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
22	MG	A	1655	1/1	0.91	0.47	-	94,94,94,94	0
22	MG	A	1619	1/1	0.95	0.92	-	61,61,61,61	0
22	MG	A	1734	1/1	0.98	0.26	-	118,118,118,118	0
22	MG	A	1615	1/1	0.77	0.87	-	106,106,106,106	0
22	MG	B	301	1/1	0.93	0.37	-	135,135,135,135	0
25	ZN	D	301	1/1	0.99	0.41	-	99,99,99,99	0
22	MG	A	1671	1/1	0.89	0.66	-	91,91,91,91	0
22	MG	A	1733	1/1	0.77	0.21	-	111,111,111,111	0
22	MG	P	101	1/1	0.77	0.15	-	102,102,102,102	0
22	MG	A	1676	1/1	0.87	2.15	-	77,77,77,77	0
22	MG	A	1661	1/1	0.89	0.62	-	76,76,76,76	0
22	MG	A	1654	1/1	0.74	0.91	-	84,84,84,84	0
22	MG	A	1650	1/1	0.62	0.33	-	101,101,101,101	0
22	MG	A	1721	1/1	0.97	0.04	-	111,111,111,111	0
22	MG	A	1633	1/1	0.91	1.90	-	80,80,80,80	0
22	MG	A	1620	1/1	0.69	0.58	-	110,110,110,110	0
22	MG	A	1707	1/1	0.95	0.56	-	82,82,82,82	0
22	MG	A	1739	1/1	0.85	0.36	-	90,90,90,90	0
23	K	A	1719	1/1	0.91	0.32	-	160,160,160,160	0
22	MG	A	1678	1/1	0.92	0.16	-	89,89,89,89	0
22	MG	A	1637	1/1	0.93	0.63	-	63,63,63,63	0
22	MG	S	102	1/1	0.78	0.17	-	100,100,100,100	0
22	MG	A	1731	1/1	0.99	0.32	-	106,106,106,106	0
22	MG	A	1622	1/1	0.80	0.81	-	66,66,66,66	0
22	MG	A	1602	1/1	0.95	1.14	-	84,84,84,84	0
22	MG	L	201	1/1	0.85	0.08	-	98,98,98,98	0
22	MG	A	1690	1/1	0.96	0.65	-	76,76,76,76	0
22	MG	A	1662	1/1	0.81	0.29	-	105,105,105,105	0
22	MG	A	1699	1/1	0.94	0.11	-	78,78,78,78	0
22	MG	A	1697	1/1	0.97	0.54	-	66,66,66,66	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.