



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 6, 2017 – 08:15 AM EDT

PDB ID : 5LGS
Title : Crystal structure of mouse CARM1 in complex with ligand P2C3u
Authors : Marechal, N.; Troffer-Charlier, N.; Cura, V.; Bonnefond, L.; Cavarelli, J.
Deposited on : 2016-07-08
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029077
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029077


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION






The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	 93% 5%
1	B	361	 93% 5%
1	C	361	 94% 5%
1	D	361	 92% 5%
2	E	9	 89% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain		
2	F	9		89%	11%
2	G	9		89%	11%
2	H	9		89%	11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	DXE	A	502	-	-	-	X
4	DXE	D	502	-	-	-	X
5	PEG	A	503	-	-	-	X
7	EDO	C	502	-	-	-	X

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 23279 atoms, of which 11249 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-arginine methyltransferase CARM1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	344	Total	C	H	N	O	S	0	6	0
			5498	1788	2725	460	510	15			
1	B	343	Total	C	H	N	O	S	0	9	0
			5500	1788	2728	459	508	17			
1	C	343	Total	C	H	N	O	S	0	5	0
			5451	1776	2700	454	507	14			
1	D	343	Total	C	H	N	O	S	0	9	0
			5499	1788	2727	459	508	17			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	GLY	-	expression tag	UNP Q9WVG6
A	128	HIS	-	expression tag	UNP Q9WVG6
A	129	MET	-	expression tag	UNP Q9WVG6
B	127	GLY	-	expression tag	UNP Q9WVG6
B	128	HIS	-	expression tag	UNP Q9WVG6
B	129	MET	-	expression tag	UNP Q9WVG6
C	127	GLY	-	expression tag	UNP Q9WVG6
C	128	HIS	-	expression tag	UNP Q9WVG6
C	129	MET	-	expression tag	UNP Q9WVG6
D	127	GLY	-	expression tag	UNP Q9WVG6
D	128	HIS	-	expression tag	UNP Q9WVG6
D	129	MET	-	expression tag	UNP Q9WVG6

- Molecule 2 is a protein called Polyadenylate-binding protein 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	E	8	Total	C	H	N	O	0	0	0
			116	39	57	11	9			
2	F	9	Total	C	H	N	O	0	0	0
			132	44	66	12	10			

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	G	8	Total	C	H	N	O	0	0	0
			116	39	57	11	9			
2	H	8	Total	C	H	N	O	0	0	0
			116	39	57	11	9			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 1,2-DIMETHOXYETHANE (three-letter code: DXE) (formula: C₄H₁₀O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	16	4	10	2	0	0
4	B	1	16	4	10	2	0	0
4	D	1	16	4	10	2	0	0

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
5	A	1	17	4	10	3	0	0

- Molecule 6 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
6	B	1	31	8	18	5	0	0

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
7	C	1	10	2	6	2	0	0
7	C	1	10	2	6	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	D	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 8 is (2 {R},3 {R},4 {S},5 {R})-2-(6-aminopurin-9-yl)-5-[({E})-prop-1-enyl]oxolane-3,4-diol (three-letter code: QVR) (formula: C₁₂H₁₅N₅O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	E	1	Total	C	H	N	O	0	0
			34	12	14	5	3		
8	F	1	Total	C	H	N	O	0	0
			34	12	14	5	3		
8	G	1	Total	C	H	N	O	0	0
			34	12	14	5	3		
8	H	1	Total	C	H	N	O	0	0
			34	12	14	5	3		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	194	Total	O	0	0
			194	194		
9	B	171	Total	O	0	0
			171	171		
9	C	109	Total	O	0	0
			109	109		
9	D	89	Total	O	0	0
			89	89		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	E	7	Total O 7 7	0	0
9	F	7	Total O 7 7	0	0
9	G	2	Total O 2 2	0	0
9	H	5	Total O 5 5	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone-arginine methyltransferase CARM1




- Molecule 1: Histone-arginine methyltransferase CARM1



- Molecule 1: Histone-arginine methyltransferase CARM1




- Molecule 1: Histone-arginine methyltransferase CARM1




- Molecule 2: Polyadenylate-binding protein 1




- Molecule 2: Polyadenylate-binding protein 1



- Molecule 2: Polyadenylate-binding protein 1



- Molecule 2: Polyadenylate-binding protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	74.99Å 98.60Å 208.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.97 – 2.10 47.97 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.97-2.10) 100.0 (47.97-2.10)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.10Å)	Xtrriage
Refinement program	PHENIX (dev_2386: ???)	Depositor
R, R_{free}	0.186 , 0.215 0.181 , 0.212	Depositor DCC
R_{free} test set	4501 reflections (4.96%)	DCC
Wilson B-factor (Å ²)	37.0	Xtrriage
Anisotropy	0.328	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 46.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	23279	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.95 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.2281e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, PG4, SO4, DXE, PEG, QVR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/2865	0.45	0/3882
1	B	0.25	0/2884	0.44	0/3906
1	C	0.25	0/2827	0.43	0/3832
1	D	0.25	0/2877	0.43	0/3897
2	E	0.37	0/62	0.44	0/85
2	F	0.35	0/70	0.49	0/96
2	G	0.41	0/62	0.43	0/85
2	H	0.45	0/62	0.55	0/85
All	All	0.26	0/11709	0.44	0/15868

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2773	2725	2705	4	0
1	B	2772	2728	2682	4	0
1	C	2751	2700	2684	1	0
1	D	2772	2727	2686	4	0
2	E	59	57	59	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	66	66	66	1	0
2	G	59	57	59	0	0
2	H	59	57	59	0	0
3	A	5	0	0	0	0
4	A	6	10	10	0	0
4	B	6	10	10	0	0
4	D	6	10	10	0	0
5	A	7	10	10	0	0
6	B	13	18	18	0	0
7	C	8	12	12	0	0
7	D	4	6	6	0	0
8	E	20	14	0	0	0
8	F	20	14	0	0	0
8	G	20	14	0	0	0
8	H	20	14	0	0	0
9	A	194	0	0	1	0
9	B	171	0	0	2	0
9	C	109	0	0	0	0
9	D	89	0	0	0	0
9	E	7	0	0	0	0
9	F	7	0	0	0	0
9	G	2	0	0	0	0
9	H	5	0	0	0	0
All	All	12030	11249	11076	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:182:THR:O	9:B:601:HOH:O	2.17	0.58
1:D:169:ARG:NE	1:D:258:GLU:OE2	2.38	0.54
1:D:390:PHE:HE1	1:D:421:CYS:HG	1.60	0.46
1:A:160:GLN:NE2	9:A:610:HOH:O	2.45	0.46
1:D:266:ASN:O	1:D:267:GLU:HB2	2.18	0.43
1:A:266:ASN:O	1:A:267:GLU:HB2	2.18	0.43
1:B:175:ARG:NH2	9:B:606:HOH:O	2.43	0.43
1:C:266:ASN:O	1:C:267:GLU:HB2	2.20	0.42
1:B:295:LEU:HA	1:B:389:ALA:O	2.19	0.42
1:A:280:LEU:HD11	1:A:284:GLY:HA3	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:451:ILE:HB	1:D:468:LEU:HB2	2.02	0.41
1:B:266:ASN:O	1:B:267:GLU:HB2	2.21	0.40
1:A:179:GLN:NE2	1:A:401:MET:SD	2.92	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/361 (96%)	336 (97%)	10 (3%)	0	100	100
1	B	348/361 (96%)	336 (97%)	12 (3%)	0	100	100
1	C	342/361 (95%)	332 (97%)	10 (3%)	0	100	100
1	D	348/361 (96%)	338 (97%)	10 (3%)	0	100	100
2	E	6/9 (67%)	6 (100%)	0	0	100	100
2	F	7/9 (78%)	7 (100%)	0	0	100	100
2	G	6/9 (67%)	6 (100%)	0	0	100	100
2	H	6/9 (67%)	4 (67%)	2 (33%)	0	100	100
All	All	1409/1480 (95%)	1365 (97%)	44 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/313 (97%)	303 (100%)	1 (0%)	94	96
1	B	305/313 (97%)	302 (99%)	3 (1%)	80	85
1	C	300/313 (96%)	300 (100%)	0	100	100
1	D	304/313 (97%)	301 (99%)	3 (1%)	80	85
2	E	6/7 (86%)	6 (100%)	0	100	100
2	F	7/7 (100%)	7 (100%)	0	100	100
2	G	6/7 (86%)	6 (100%)	0	100	100
2	H	6/7 (86%)	6 (100%)	0	100	100
All	All	1238/1280 (97%)	1231 (99%)	7 (1%)	89	92

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	295	LEU
1	B	136	SER
1	B	295	LEU
1	B	328	ARG
1	D	169	ARG
1	D	295	LEU
1	D	308	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	501	-	4,4,4	0.15	0	6,6,6	0.07	0
4	DXE	A	502	-	5,5,5	0.42	0	4,4,4	0.25	0
5	PEG	A	503	-	6,6,6	0.50	0	5,5,5	0.49	0
4	DXE	B	501	-	5,5,5	0.42	0	4,4,4	0.24	0
6	PG4	B	502	-	12,12,12	0.53	0	11,11,11	0.31	0
7	EDO	C	501	-	3,3,3	0.46	0	2,2,2	0.51	0
7	EDO	C	502	-	3,3,3	0.47	0	2,2,2	0.54	0
7	EDO	D	501	-	3,3,3	0.48	0	2,2,2	0.51	0
4	DXE	D	502	-	5,5,5	0.43	0	4,4,4	0.29	0
8	QVR	E	1001	2	19,22,22	0.56	0	17,32,32	0.61	0
8	QVR	F	1001	2	19,22,22	0.57	0	17,32,32	0.59	0
8	QVR	G	101	2	19,22,22	0.56	0	17,32,32	0.58	0
8	QVR	H	101	2	19,22,22	0.57	0	17,32,32	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	501	-	-	0/0/0/0	0/0/0/0
4	DXE	A	502	-	-	0/3/3/3	0/0/0/0
5	PEG	A	503	-	-	0/4/4/4	0/0/0/0
4	DXE	B	501	-	-	0/3/3/3	0/0/0/0
6	PG4	B	502	-	-	0/10/10/10	0/0/0/0
7	EDO	C	501	-	-	0/1/1/1	0/0/0/0
7	EDO	C	502	-	-	0/1/1/1	0/0/0/0
7	EDO	D	501	-	-	0/1/1/1	0/0/0/0
4	DXE	D	502	-	-	0/3/3/3	0/0/0/0
8	QVR	E	1001	2	-	0/2/23/23	0/3/3/3
8	QVR	F	1001	2	-	0/2/23/23	0/3/3/3
8	QVR	G	101	2	-	0/2/23/23	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	QVR	H	101	2	-	0/2/23/23	0/3/3/3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	344/361 (95%)	0.04	2 (0%) 89 91	21, 35, 56, 84	0
1	B	343/361 (95%)	0.07	5 (1%) 74 77	25, 40, 59, 114	0
1	C	343/361 (95%)	0.30	7 (2%) 65 70	34, 51, 75, 90	0
1	D	343/361 (95%)	0.24	9 (2%) 56 62	38, 50, 70, 91	0
2	E	8/9 (88%)	0.67	1 (12%) 4 6	37, 40, 56, 60	0
2	F	9/9 (100%)	1.28	1 (11%) 6 7	38, 45, 73, 93	0
2	G	8/9 (88%)	0.58	1 (12%) 4 6	46, 49, 67, 72	0
2	H	8/9 (88%)	0.99	1 (12%) 4 6	48, 51, 65, 72	0
All	All	1406/1480 (95%)	0.18	27 (1%) 67 71	21, 46, 70, 114	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	-3	PRO	6.1
1	B	135	ARG	5.2
1	D	347	ARG	4.0
2	E	-2	ALA	3.9
2	G	-2	ALA	3.8
1	D	135	ARG	3.7
2	H	-2	ALA	3.6
1	B	347	ARG	3.0
1	C	284	GLY	2.8
1	A	283	SER	2.8
1	B	467	LEU	2.7
1	D	346	ILE	2.6
1	D	248	LEU	2.5
1	D	380	LEU	2.4
1	C	208	ALA	2.4
1	D	350[A]	MET	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	183	ASP	2.4
1	D	378	HIS	2.3
1	C	276	ALA	2.3
1	B	350[A]	MET	2.3
1	C	282	PRO	2.3
1	D	477	TYR	2.2
1	C	209	ARG	2.2
1	A	135	ARG	2.1
1	B	346	ILE	2.0
1	D	421	CYS	2.0
1	C	368	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	EDO	C	502	4/4	0.77	0.29	4.62	47,56,57,60	0
4	DXE	A	502	6/6	0.73	0.34	3.42	43,51,55,55	0
4	DXE	D	502	6/6	0.68	0.28	3.05	48,58,63,63	0
5	PEG	A	503	7/7	0.73	0.20	2.41	40,48,52,53	0
6	PG4	B	502	13/13	0.68	0.34	1.94	74,89,98,99	0
8	QVR	F	1001	20/20	0.97	0.14	0.30	29,33,41,42	0
8	QVR	G	101	20/20	0.95	0.13	-0.08	41,46,56,56	0
3	SO4	A	501	5/5	0.93	0.14	-0.47	89,90,90,91	0
8	QVR	H	101	20/20	0.96	0.11	-0.90	42,45,55,56	0
8	QVR	E	1001	20/20	0.96	0.10	-1.03	28,31,42,44	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
7	EDO	D	501	4/4	0.92	0.13	-	53,64,65,66	0
7	EDO	C	501	4/4	0.85	0.21	-	58,70,74,77	0
4	DXE	B	501	6/6	0.83	0.15	-	55,66,73,73	0

6.5 Other polymers [i](#)

There are no such residues in this entry.