



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 15, 2018 – 07:45 AM EDT

PDB ID : 5LLB  
Title : Structure of Polyphosphate Kinase 2 from *Francisella tularensis* with AMP-PCH2PPP and polyphosphate  
Authors : Roach, P.L.; Parnell, A.E.  
Deposited on : 2016-07-27  
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030736  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	6276 (1.94-1.90)
Clashscore	112137	7025 (1.94-1.90)
Ramachandran outliers	110173	6947 (1.94-1.90)
Sidechain outliers	110143	6948 (1.94-1.90)
RSRZ outliers	101464	6332 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	<div> <div>4%</div> <div>86%</div> <div>11%</div> </div>
1	B	269	<div> <div>4%</div> <div>81%</div> <div>8%</div> <div>11%</div> </div>
2	C	263	<div> <div>2%</div> <div>86%</div> <div>5%</div> <div>9%</div> </div>
2	D	263	<div> <div>3%</div> <div>86%</div> <div>5%</div> <div>9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria.

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	C	301	-	-	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 16918 atoms, of which 7943 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyphosphate kinase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	240	Total	C	H	N	O	S	0	0	0
			3969	1294	1959	351	358	7			
1	B	240	Total	C	H	N	O	S	0	0	0
			3960	1293	1951	351	358	7			

- Molecule 2 is a protein called Polyphosphate kinase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	240	Total	C	H	N	O	S	0	0	0
			3998	1294	1988	351	358	7			
2	D	240	Total	C	H	N	O	S	0	0	0
			3999	1294	1989	351	358	7			

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	1	Total	Cl	0	0
			1	1		
3	D	1	Total	Cl	0	0
			1	1		
3	C	1	Total	Cl	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	A	1	Total	Mg	0	0
			1	1		


*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		

- Molecule 5 is [2 {R},3 {S},4 {R},5 {R}]-5-(6-aminopurin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methoxy-oxidanyl-phosphoryl]oxy-bis(oxidanyl)-5-phosphanyl]methyl-bis(oxidanyl)-5-phosphanyl] phosphono hydrogen phosphate (three-letter code: 6YZ) (formula: C<sub>11</sub>H<sub>24</sub>N<sub>5</sub>O<sub>18</sub>P<sub>5</sub>).







Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total 25	O 19	P 6	0	0
6	B	1	Total 25	O 19	P 6	0	0
6	C	1	Total 25	O 19	P 6	0	0
6	D	1	Total 25	O 19	P 6	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	164	Total O 164 164	0	0
7	B	163	Total O 163 163	0	0
7	C	201	Total O 201 201	0	0
7	D	144	Total O 144 144	0	0



- Molecule 1: Polyphosphate kinase 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.70Å 144.91Å 70.59Å 90.00° 113.04° 90.00°	Depositor
Resolution (Å)	72.46 – 1.92 43.31 – 1.92	Depositor EDS
% Data completeness (in resolution range)	99.2 (72.46-1.92) 93.3 (43.31-1.92)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.51 (at 1.92Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.179 , 0.231 0.177 , 0.230	Depositor DCC
$R_{free}$ test set	3737 reflections (4.85%)	DCC
Wilson B-factor (Å <sup>2</sup> )	26.4	Xtriage
Anisotropy	0.682	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 78.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.045 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	16918	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6YZ, MG, 6YW, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.57	0/2062	0.63	0/2781
1	B	0.56	0/2061	0.64	0/2780
2	C	0.57	0/2062	0.64	0/2781
2	D	0.54	0/2062	0.63	0/2781
All	All	0.56	0/8247	0.64	0/11123

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2010	1959	2000	9	0
1	B	2009	1951	1998	12	0
2	C	2010	1988	2000	7	0
2	D	2010	1989	2000	6	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	39	14	0	1	0
5	B	39	14	0	0	0
5	C	39	14	0	1	0
5	D	39	14	0	0	0
6	A	25	0	0	2	0
6	B	25	0	0	1	0
6	C	25	0	0	0	0
6	D	25	0	0	1	0
7	A	164	0	0	3	0
7	B	163	0	0	1	1
7	C	201	0	0	5	1
7	D	144	0	0	0	0
All	All	8975	7943	7998	36	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:26:ARG:NH2	6:B:304:6YW:O15	1.97	0.97
2:D:228:LYS:NZ	6:D:304:6YW:O18	2.14	0.79
2:C:31:LYS:NZ	7:C:401:HOH:O	2.18	0.76
1:B:172:LYS:NZ	7:B:401:HOH:O	2.21	0.73
5:C:303:6YZ:N31	7:C:403:HOH:O	2.26	0.68
5:A:303:6YZ:P02	7:A:401:HOH:O	2.59	0.60
1:A:228:LYS:NZ	6:A:304:6YW:O25	2.32	0.56
2:D:43:PHE:HE1	2:D:243:VAL:HG21	1.71	0.55
1:A:48:LYS:NZ	1:A:80:ARG:O	2.40	0.54
2:C:150:LYS:NZ	7:C:408:HOH:O	2.41	0.53
1:B:185:GLN:O	1:B:188:LEU:HG	2.09	0.53
1:B:26:ARG:NH2	1:B:30:GLU:OE2	2.42	0.52
1:B:117:ASP:O	1:B:118:ARG:HB2	2.12	0.49
1:B:225:SER:HA	1:B:231:ALA:HB2	1.94	0.49
1:A:83:LYS:NZ	7:A:402:HOH:O	2.45	0.49
2:C:117:ASP:O	2:C:118:ARG:HB2	2.14	0.47
1:A:25:PRO:HG2	1:A:28:VAL:HG12	1.96	0.47
1:B:163:TRP:HB3	1:B:222:ILE:HD13	1.96	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:LYS:HG2	7:A:487:HOH:O	2.16	0.46
1:B:68:GLY:O	1:B:72:ARG:HD3	2.17	0.45
1:B:175:PHE:HD2	1:B:196:LEU:HD21	1.81	0.44
1:A:184:LYS:NZ	6:A:304:6YW:O09	2.37	0.44
2:D:43:PHE:CE1	2:D:243:VAL:HG21	2.51	0.44
2:C:43:PHE:CE1	2:C:243:VAL:HG11	2.53	0.44
2:C:191:ILE:CD1	7:C:559:HOH:O	2.65	0.44
1:A:25:PRO:O	1:A:28:VAL:HG12	2.18	0.43
1:B:93:GLN:O	1:B:97:GLN:HG3	2.18	0.43
2:D:117:ASP:O	2:D:118:ARG:HB2	2.18	0.43
2:C:52:LYS:HE3	2:C:158:MET:HB2	2.00	0.42
1:A:24:ILE:CG1	1:A:25:PRO:HD2	2.50	0.42
1:B:24:ILE:CG1	1:B:25:PRO:HD2	2.50	0.41
2:C:232:ARG:NE	7:C:402:HOH:O	2.25	0.41
1:A:117:ASP:O	1:A:118:ARG:HB2	2.21	0.41
2:D:93:GLN:O	2:D:97:GLN:HG3	2.21	0.41
2:D:82:ALA:O	2:D:83:LYS:HD3	2.21	0.40
1:B:190:PRO:O	1:B:193:LYS:HB2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:448:HOH:O	7:C:457:HOH:O[2_655]	1.98	0.22

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	238/269 (88%)	233 (98%)	5 (2%)	0	100	100
1	B	238/269 (88%)	234 (98%)	4 (2%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	238/263 (90%)	234 (98%)	4 (2%)	0	100	100
2	D	238/263 (90%)	230 (97%)	8 (3%)	0	100	100
All	All	952/1064 (90%)	931 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/244 (88%)	214 (100%)	0	100	100
1	B	214/244 (88%)	210 (98%)	4 (2%)	62	56
2	C	214/238 (90%)	210 (98%)	4 (2%)	62	56
2	D	214/238 (90%)	209 (98%)	5 (2%)	56	48
All	All	856/964 (89%)	843 (98%)	13 (2%)	70	66

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	26	ARG
1	B	188	LEU
1	B	191	ILE
1	B	195	SER
2	C	79	PRO
2	C	135	GLU
2	C	166	VAL
2	C	191	ILE
2	D	95	ARG
2	D	188	LEU
2	D	197	ASP
2	D	246	ASP
2	D	247	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	76	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	6YZ	A	303	4	29,41,41	4.40	10 (34%)	27,66,66	1.57	2 (7%)
6	6YW	A	304	-	16,24,24	0.76	1 (6%)	12,41,41	0.46	0
5	6YZ	B	303	4	29,41,41	4.31	10 (34%)	27,66,66	1.65	4 (14%)
6	6YW	B	304	-	16,24,24	0.88	1 (6%)	12,41,41	0.35	0
5	6YZ	C	303	4	29,41,41	3.91	10 (34%)	27,66,66	1.62	2 (7%)
6	6YW	C	304	-	16,24,24	0.92	1 (6%)	12,41,41	0.72	0
5	6YZ	D	303	4	29,41,41	4.30	11 (37%)	27,66,66	1.66	2 (7%)
6	6YW	D	304	-	16,24,24	0.89	1 (6%)	12,41,41	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	6YZ	A	303	4	-	0/22/50/50	0/3/3/3
6	6YW	A	304	-	-	0/30/30/30	0/0/0/0
5	6YZ	B	303	4	-	0/22/50/50	0/3/3/3
6	6YW	B	304	-	-	0/30/30/30	0/0/0/0
5	6YZ	C	303	4	-	0/22/50/50	0/3/3/3
6	6YW	C	304	-	-	0/30/30/30	0/0/0/0
5	6YZ	D	303	4	-	0/22/50/50	0/3/3/3
6	6YW	D	304	-	-	0/30/30/30	0/0/0/0

All (45) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	303	6YZ	C32-C21	-13.49	1.32	1.53
5	D	303	6YZ	C32-C21	-13.40	1.32	1.53
5	A	303	6YZ	C32-C21	-12.98	1.33	1.53
5	C	303	6YZ	C32-C21	-12.41	1.33	1.53
5	D	303	6YZ	O20-C19	-7.40	1.28	1.45
5	A	303	6YZ	O20-C19	-7.33	1.28	1.45
5	B	303	6YZ	O20-C19	-7.13	1.28	1.45
5	C	303	6YZ	O20-C19	-6.85	1.29	1.45
5	A	303	6YZ	O35-C34	-3.49	1.34	1.43
5	B	303	6YZ	O35-C34	-3.05	1.35	1.43
6	C	304	6YW	P02-O04	-2.95	1.55	1.60
5	D	303	6YZ	O35-C34	-2.76	1.36	1.43
6	B	304	6YW	P02-O04	-2.46	1.56	1.60
5	C	303	6YZ	O35-C34	-2.36	1.37	1.43
6	D	304	6YW	P17-O16	-2.28	1.56	1.60
6	A	304	6YW	P02-O04	-2.12	1.56	1.60
5	D	303	6YZ	C28-N27	2.02	1.35	1.32
5	C	303	6YZ	C28-N27	2.33	1.36	1.32
5	C	303	6YZ	C30-N31	2.68	1.45	1.34
5	D	303	6YZ	O33-C32	2.69	1.49	1.43
5	B	303	6YZ	C30-N31	2.80	1.45	1.34
5	A	303	6YZ	C30-N31	2.84	1.45	1.34
5	A	303	6YZ	P04-O03	2.88	1.64	1.60
5	A	303	6YZ	O33-C32	2.92	1.49	1.43
5	D	303	6YZ	C30-N31	2.93	1.46	1.34
5	C	303	6YZ	O33-C32	3.13	1.50	1.43
5	B	303	6YZ	O33-C32	3.17	1.50	1.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	303	6YZ	P04-O03	3.20	1.65	1.60
5	D	303	6YZ	C26-N27	3.22	1.40	1.35
5	C	303	6YZ	C26-N27	3.30	1.40	1.35
5	B	303	6YZ	C26-N27	3.38	1.40	1.35
5	A	303	6YZ	C26-N27	3.69	1.41	1.35
5	A	303	6YZ	P12-O14	3.96	1.61	1.55
5	C	303	6YZ	P12-O14	4.06	1.61	1.55
5	D	303	6YZ	P12-O14	4.16	1.61	1.55
5	B	303	6YZ	P12-O14	4.50	1.62	1.55
5	D	303	6YZ	P09-O08	4.52	1.62	1.55
5	C	303	6YZ	P09-O08	4.71	1.62	1.55
5	B	303	6YZ	P04-O03	4.76	1.67	1.60
5	B	303	6YZ	P09-O08	4.85	1.62	1.55
5	A	303	6YZ	P09-O08	6.50	1.65	1.55
5	C	303	6YZ	O20-C21	12.51	1.58	1.41
5	B	303	6YZ	O20-C21	13.63	1.60	1.41
5	D	303	6YZ	O20-C21	14.27	1.61	1.41
5	A	303	6YZ	O20-C21	14.63	1.61	1.41

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	303	6YZ	N27-C28-N29	-7.38	122.43	128.86
5	D	303	6YZ	N27-C28-N29	-7.19	122.60	128.86
5	A	303	6YZ	N27-C28-N29	-6.59	123.12	128.86
5	B	303	6YZ	N27-C28-N29	-6.46	123.23	128.86
5	D	303	6YZ	C26-C25-N24	-3.37	106.16	109.41
5	B	303	6YZ	C19-O20-C21	-2.77	106.82	109.77
5	B	303	6YZ	C18-C19-C34	-2.43	106.03	115.29
5	C	303	6YZ	C18-C19-C34	-2.22	106.84	115.29
5	B	303	6YZ	C26-C25-N24	-2.17	107.32	109.41
5	A	303	6YZ	C18-C19-C34	-2.09	107.32	115.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	303	6YZ	1	0
6	A	304	6YW	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	304	6YW	1	0
5	C	303	6YZ	1	0
6	D	304	6YW	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	240/269 (89%)	0.32	12 (5%) 30 33	37, 49, 73, 89	0
1	B	240/269 (89%)	0.30	11 (4%) 33 37	38, 50, 72, 123	0
2	C	240/263 (91%)	0.09	4 (1%) 70 74	37, 48, 66, 82	0
2	D	240/263 (91%)	0.41	9 (3%) 41 45	38, 53, 77, 90	0
All	All	960/1064 (90%)	0.28	36 (3%) 41 45	37, 49, 73, 123	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	188	LEU	5.5
1	B	191	ILE	5.0
1	B	188	LEU	4.6
2	D	24	ILE	4.2
2	D	166	VAL	4.0
1	A	28	VAL	3.5
1	A	190	PRO	3.5
2	D	191	ILE	3.3
1	A	191	ILE	3.3
1	B	25	PRO	3.3
1	B	193	LYS	3.2
1	A	196	LEU	3.1
1	A	24	ILE	2.6
2	D	188	LEU	2.6
1	A	192	ASP	2.5
1	B	28	VAL	2.4
2	C	27	ASN	2.4
1	A	251	GLU	2.4
1	A	27	ASN	2.4
1	B	196	LEU	2.3
1	A	132	PHE	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	D	175	PHE	2.3
1	A	263	THR	2.2
1	B	192	ASP	2.2
2	D	246	ASP	2.2
2	C	191	ILE	2.2
1	B	186	TRP	2.2
1	B	139	PHE	2.1
1	B	24	ILE	2.1
2	D	263	THR	2.1
2	C	244	ASP	2.1
1	A	181	HIS	2.1
1	B	32	GLN	2.1
2	C	24	ILE	2.1
2	D	210	PHE	2.0
2	D	211	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	CL	C	301	1/1	0.97	0.15	2.82	50,50,50,50	0
6	6YW	D	304	25/25	0.68	0.18	1.65	83,115,139,147	0
6	6YW	C	304	25/25	0.73	0.17	1.62	69,92,118,150	0
4	MG	A	302	1/1	0.92	0.26	1.33	62,62,62,62	0
6	6YW	B	304	25/25	0.73	0.17	1.01	68,96,112,122	0
3	CL	A	301	1/1	0.91	0.13	0.85	57,57,57,57	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	MG	C	302	1/1	0.98	0.15	0.66	45,45,45,45	0
6	6YW	A	304	25/25	0.67	0.17	0.50	75,106,151,161	0
4	MG	B	302	1/1	0.94	0.18	-0.02	73,73,73,73	0
5	6YZ	B	303	39/39	0.89	0.15	-0.17	49,67,106,123	0
5	6YZ	C	303	39/39	0.95	0.11	-0.37	40,52,62,64	0
5	6YZ	D	303	39/39	0.92	0.12	-0.55	53,63,75,79	0
5	6YZ	A	303	39/39	0.92	0.12	-0.96	50,64,79,88	0
4	MG	D	302	1/1	0.93	0.10	-1.43	58,58,58,58	0
3	CL	D	301	1/1	0.96	0.08	-1.63	72,72,72,72	0
3	CL	B	301	1/1	0.94	0.10	-1.92	62,62,62,62	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.