



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2017 – 05:27 PM EDT

PDB ID : 3LMG  
Title : Crystal structure of the ERBB3 kinase domain in complex with AMP-PNP  
Authors : Shi, F.; Lemmon, M.A.  
Deposited on : unknown  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	<div> <div>4%</div> <div> <div></div> <div>55%</div> <div>21%</div> <div>•</div> <div>22%</div> </div> </div>
1	B	344	<div> <div>60%</div> <div>17%</div> <div>•</div> <div>22%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4288 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Receptor tyrosine-protein kinase erbB-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	0	0	0
			2108	1363	363	369	13			
1	B	268	Total	C	N	O	S	0	0	0
			2109	1364	360	372	13			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	658	MET	-	EXPRESSION TAG	UNP P21860
A	659	HIS	-	EXPRESSION TAG	UNP P21860
A	660	HIS	-	EXPRESSION TAG	UNP P21860
A	661	HIS	-	EXPRESSION TAG	UNP P21860
A	662	HIS	-	EXPRESSION TAG	UNP P21860
A	663	HIS	-	EXPRESSION TAG	UNP P21860
A	664	HIS	-	EXPRESSION TAG	UNP P21860
B	658	MET	-	EXPRESSION TAG	UNP P21860
B	659	HIS	-	EXPRESSION TAG	UNP P21860
B	660	HIS	-	EXPRESSION TAG	UNP P21860
B	661	HIS	-	EXPRESSION TAG	UNP P21860
B	662	HIS	-	EXPRESSION TAG	UNP P21860
B	663	HIS	-	EXPRESSION TAG	UNP P21860
B	664	HIS	-	EXPRESSION TAG	UNP P21860

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter

code: ANP) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>6</sub>O<sub>12</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 31	C 10	N 6	O 12	P 3	0	0
3	B	1	Total 31	C 10	N 6	O 12	P 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total O 2 2	0	0
4	B	5	Total O 5 5	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Receptor tyrosine-protein kinase erbB-3



- Molecule 1: Receptor tyrosine-protein kinase erbB-3



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	193.35Å 47.98Å 82.22Å 90.00° 108.11° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 41.37 – 2.80	Depositor EDS
% Data completeness (in resolution range)	89.1 (50.00-2.80) 89.2 (41.37-2.80)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.03 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, $R_{free}$	0.252 , 0.285 0.243 , 0.277	Depositor DCC
$R_{free}$ test set	1612 reflections (11.19%)	DCC
Wilson B-factor (Å <sup>2</sup> )	62.7	Xtriage
Anisotropy	0.580	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 75.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	4288	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/2158	0.52	0/2929
1	B	0.33	0/2158	0.52	0/2927
All	All	0.32	0/4316	0.52	0/5856

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2108	0	2136	75	0
1	B	2109	0	2133	43	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	31	0	13	1	0
3	B	31	0	13	1	0
4	A	2	0	0	0	0
4	B	5	0	0	0	0
All	All	4288	0	4295	116	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (116) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:685:PHE:CE2	1:A:757:GLY:HA3	1.66	1.28
1:B:680:VAL:N	1:B:681:LEU:HA	1.72	1.03
1:A:685:PHE:HE2	1:A:757:GLY:HA3	0.95	1.01
1:A:685:PHE:HE2	1:A:757:GLY:CA	1.73	1.01
1:A:741:MET:HE2	1:A:758:LEU:HD21	1.43	1.01
1:A:741:MET:CE	1:A:758:LEU:HD21	2.02	0.90
1:A:758:LEU:HD11	1:A:764:LEU:HD11	1.52	0.89
1:A:751:HIS:CE1	1:A:798:GLN:HG2	2.07	0.89
1:A:861:LEU:HD13	1:A:905:LEU:HD11	1.60	0.82
1:A:870:THR:HB	1:A:872:GLN:OE1	1.84	0.78
1:B:737:VAL:HA	1:B:741:MET:CE	2.14	0.76
1:B:737:VAL:HA	1:B:741:MET:HE2	1.69	0.73
1:A:815:ASN:O	1:A:820:ASN:ND2	2.22	0.73
1:A:811:MET:HE1	1:A:835:GLY:HA3	1.71	0.73
1:A:683:ARG:HD3	1:A:685:PHE:CZ	2.24	0.72
1:B:781:ARG:NH1	1:B:889:GLY:O	2.23	0.72
1:B:881:THR:O	1:B:885:LEU:HD13	1.92	0.69
1:A:751:HIS:HE1	1:A:798:GLN:HG2	1.58	0.69
1:A:790:GLN:HE21	1:A:794:ASN:HD21	1.41	0.68
1:A:903:ASP:O	1:A:907:LYS:HG2	1.93	0.68
1:B:803:MET:HE1	1:B:831:VAL:HG11	1.76	0.66
1:A:683:ARG:HD3	1:A:685:PHE:CE1	2.31	0.65
1:A:861:LEU:HG	1:A:865:HIS:CD2	2.33	0.64
1:B:735:GLN:O	1:B:736:ALA:HB3	1.97	0.63
1:A:735:GLN:O	1:A:736:ALA:HB3	1.99	0.63
1:A:711:PRO:HG2	1:A:716:ILE:HD11	1.80	0.62
1:A:711:PRO:HG2	1:A:716:ILE:CD1	2.29	0.62
1:A:880:VAL:O	1:A:884:GLU:HG3	2.00	0.61
1:B:684:ILE:HD11	1:B:737:VAL:HG21	1.83	0.60
1:B:737:VAL:HA	1:B:741:MET:HE1	1.85	0.59
1:A:699:GLY:HA3	3:A:301:ANP:O3G	2.02	0.59
1:A:811:MET:CE	1:A:835:GLY:HA3	2.32	0.59
1:A:741:MET:HE3	1:A:764:LEU:HD11	1.85	0.58
1:A:790:GLN:NE2	1:A:794:ASN:HD21	2.01	0.57
1:B:823:LEU:HD23	1:B:829:VAL:HG12	1.87	0.57
1:B:734:PHE:CD1	1:B:764:LEU:HB2	2.41	0.56
1:B:921:VAL:HG22	1:B:949:MET:CE	2.36	0.56
1:A:689:GLU:HB3	1:A:709:TRP:HE1	1.71	0.56
1:A:685:PHE:CE2	1:A:757:GLY:CA	2.57	0.55
1:A:751:HIS:CE1	1:A:798:GLN:CG	2.85	0.55

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:803:MET:CE	1:B:831:VAL:HG11	2.37	0.54
1:A:734:PHE:CD1	1:A:764:LEU:HB2	2.42	0.54
1:B:883:TRP:HD1	1:B:925:MET:CE	2.21	0.54
1:B:680:VAL:N	1:B:681:LEU:CA	2.56	0.54
1:B:734:PHE:CG	1:B:764:LEU:HB2	2.42	0.54
1:A:861:LEU:O	1:A:865:HIS:HD2	1.92	0.53
1:A:703:THR:HG21	1:B:705:HIS:HE1	1.73	0.53
1:A:807:GLU:HA	1:A:871:HIS:CE1	2.44	0.53
1:B:953:PRO:HG2	1:B:954:PRO:HD3	1.91	0.53
1:A:741:MET:CE	1:A:764:LEU:HD11	2.39	0.52
1:A:741:MET:HE3	1:A:758:LEU:HD21	1.90	0.52
1:B:910:ARG:HH22	1:B:931:ILE:HD11	1.74	0.52
1:A:735:GLN:O	1:A:736:ALA:CB	2.59	0.51
1:A:913:GLN:HG2	1:A:922:TYR:CE2	2.46	0.51
1:B:708:VAL:CG1	1:B:717:LYS:HG3	2.41	0.51
1:A:776:LEU:CD2	1:A:885:LEU:HD11	2.41	0.51
1:B:755:LEU:HD11	1:B:766:LEU:HD22	1.94	0.50
1:A:689:GLU:HB3	1:A:709:TRP:NE1	2.26	0.50
1:A:759:CYS:HB3	1:A:765:GLN:HB2	1.94	0.49
1:A:861:LEU:O	1:A:865:HIS:CD2	2.64	0.49
1:A:751:HIS:O	1:A:830:GLN:HA	2.12	0.49
1:A:685:PHE:CD2	1:A:757:GLY:HA3	2.36	0.49
1:B:939:PHE:HA	1:B:942:LEU:HD12	1.94	0.49
1:B:883:TRP:HD1	1:B:925:MET:HE1	1.77	0.48
1:A:953:PRO:HG2	1:A:954:PRO:HD3	1.95	0.48
1:A:693:LEU:HD21	1:A:708:VAL:HG23	1.95	0.48
1:A:902:PRO:O	1:A:906:GLU:HB2	2.14	0.48
1:B:880:VAL:O	1:B:884:GLU:HG3	2.14	0.48
1:A:813:HIS:CD2	1:A:834:PHE:HB3	2.49	0.47
1:A:776:LEU:HD13	1:A:829:VAL:HG11	1.97	0.47
1:A:693:LEU:CD2	1:A:708:VAL:HG23	2.45	0.47
1:B:922:TYR:O	1:B:926:VAL:HG23	2.15	0.46
1:A:814:ARG:HG3	1:A:869:TYR:CD2	2.50	0.46
1:B:747:LEU:HB3	1:B:805:TYR:HE2	1.80	0.46
1:B:921:VAL:HG22	1:B:949:MET:HE3	1.97	0.46
1:A:695:VAL:HG21	1:B:695:VAL:HG21	1.97	0.46
1:A:813:HIS:CE1	1:A:815:ASN:C	2.89	0.46
1:B:881:THR:O	1:B:885:LEU:CD1	2.63	0.46
1:B:755:LEU:HA	1:B:768:THR:HG22	1.97	0.45
1:A:797:VAL:HG22	1:A:946:PHE:HB3	1.98	0.45
1:B:836:VAL:HA	1:B:839:LEU:HD23	1.98	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:857:LYS:HB2	1:B:858:TRP:CZ3	2.52	0.45
1:B:701:PHE:HE2	1:B:815:ASN:HD22	1.65	0.45
1:A:922:TYR:O	1:A:926:VAL:HG23	2.17	0.45
1:B:860:ALA:HA	1:B:876:TRP:CD2	2.52	0.44
1:A:758:LEU:HD11	1:A:764:LEU:CD1	2.34	0.44
1:B:759:CYS:HB3	1:B:765:GLN:HB2	2.00	0.44
1:A:813:HIS:CD2	1:A:834:PHE:HA	2.53	0.44
1:A:711:PRO:CG	1:A:716:ILE:HD11	2.47	0.44
1:A:727:ASP:OD2	1:A:732:GLN:HA	2.17	0.44
1:B:921:VAL:HG22	1:B:949:MET:HE1	1.99	0.44
1:B:949:MET:HG2	1:B:956:TYR:CE2	2.53	0.43
1:B:735:GLN:O	1:B:736:ALA:CB	2.61	0.43
1:A:860:ALA:HA	1:A:876:TRP:CD2	2.53	0.43
1:A:807:GLU:HG3	1:A:871:HIS:ND1	2.32	0.43
1:A:872:GLN:CD	1:A:872:GLN:H	2.19	0.43
1:B:699:GLY:HA3	3:B:301:ANP:O3G	2.18	0.43
1:B:910:ARG:NH1	1:B:929:TRP:O	2.52	0.43
1:A:825:SER:HB3	1:A:828:GLN:HB2	2.01	0.42
1:A:855:PRO:O	1:A:859:MET:HG3	2.20	0.42
1:A:771:LEU:HD22	1:A:824:LYS:HA	2.01	0.42
1:A:689:GLU:HB3	1:A:709:TRP:CD1	2.55	0.42
1:A:727:ASP:HB2	1:A:732:GLN:O	2.19	0.42
1:B:953:PRO:CG	1:B:954:PRO:HD3	2.49	0.42
1:A:881:THR:O	1:A:885:LEU:HD22	2.20	0.42
1:A:711:PRO:HG2	1:A:716:ILE:HD12	2.00	0.42
1:A:861:LEU:HD13	1:A:905:LEU:CD1	2.39	0.42
1:A:727:ASP:O	1:A:728:LYS:HB2	2.19	0.42
1:A:882:VAL:O	1:A:886:MET:HG2	2.19	0.41
1:A:775:SER:HA	1:A:822:LEU:HD23	2.03	0.41
1:A:683:ARG:CD	1:A:685:PHE:CZ	3.02	0.41
1:A:872:GLN:O	1:A:875:VAL:HB	2.21	0.41
1:B:932:ASP:HB3	1:B:935:ILE:HD12	2.02	0.41
1:B:913:GLN:O	1:B:913:GLN:HG3	2.20	0.40
1:A:698:SER:HB3	1:A:703:THR:HG23	2.02	0.40
1:A:813:HIS:HD2	1:A:834:PHE:HB3	1.86	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/344 (76%)	254 (97%)	7 (3%)	1 (0%)	38	72
1	B	260/344 (76%)	252 (97%)	7 (3%)	1 (0%)	38	72
All	All	522/688 (76%)	506 (97%)	14 (3%)	2 (0%)	38	72

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	736	ALA
1	B	736	ALA

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/297 (76%)	219 (96%)	8 (4%)	41	75
1	B	228/297 (77%)	223 (98%)	5 (2%)	57	87
All	All	455/594 (77%)	442 (97%)	13 (3%)	48	81

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	681	LEU
1	A	693	LEU
1	A	694	LYS
1	A	716	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	721	CYS
1	A	885	LEU
1	A	903	ASP
1	A	920	ASP
1	B	721	CYS
1	B	807	GLU
1	B	839	LEU
1	B	905	LEU
1	B	923	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	705	HIS
1	A	751	HIS
1	A	765	GLN
1	A	794	ASN
1	A	865	HIS
1	B	705	HIS
1	B	783	HIS
1	B	790	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	ANP	A	301	2	29,33,33	2.03	5 (17%)	28,52,52	2.02	7 (25%)
3	ANP	B	301	2	29,33,33	1.99	6 (20%)	28,52,52	1.96	6 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	A	301	2	-	0/13/38/38	0/3/3/3
3	ANP	B	301	2	-	0/13/38/38	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	301	ANP	PB-O3A	2.06	1.61	1.59
3	B	301	ANP	C5-C4	3.06	1.47	1.40
3	A	301	ANP	C5-C4	3.18	1.47	1.40
3	B	301	ANP	PB-N3B	4.17	1.74	1.63
3	B	301	ANP	PG-N3B	4.25	1.74	1.63
3	A	301	ANP	PB-N3B	4.32	1.74	1.63
3	A	301	ANP	PG-N3B	4.43	1.75	1.63
3	B	301	ANP	PG-O1G	4.56	1.51	1.46
3	B	301	ANP	PB-O1B	4.58	1.51	1.46
3	A	301	ANP	PG-O1G	4.61	1.51	1.46
3	A	301	ANP	PB-O1B	4.78	1.51	1.46

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	301	ANP	N3-C2-N1	-6.42	123.27	128.86
3	B	301	ANP	N3-C2-N1	-6.19	123.47	128.86
3	A	301	ANP	O1G-PG-N3B	-4.48	105.09	111.79
3	B	301	ANP	O1G-PG-N3B	-3.88	105.98	111.79

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	301	ANP	O1B-PB-N3B	-3.08	107.19	111.79
3	A	301	ANP	PA-O3A-PB	-2.76	122.65	132.38
3	A	301	ANP	C4-C5-N7	-2.53	106.96	109.41
3	B	301	ANP	C4-C5-N7	-2.48	107.01	109.41
3	A	301	ANP	O1B-PB-N3B	-2.42	108.17	111.79
3	B	301	ANP	O3G-PG-O2G	2.09	113.55	107.69
3	A	301	ANP	O3G-PG-O2G	2.24	113.97	107.69
3	A	301	ANP	O2B-PB-O1B	3.92	118.03	109.87
3	B	301	ANP	O2B-PB-O1B	4.14	118.47	109.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	ANP	1	0
3	B	301	ANP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	268/344 (77%)	0.42	14 (5%) 28 19	24, 33, 42, 44	0
1	B	268/344 (77%)	0.16	1 (0%) 92 90	20, 27, 33, 42	0
All	All	536/688 (77%)	0.29	15 (2%) 53 43	20, 30, 41, 44	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	957	LEU	4.9
1	A	866	PHE	3.1
1	A	797	VAL	3.0
1	A	930	MET	2.7
1	A	786	ALA	2.6
1	A	823	LEU	2.4
1	A	791	LEU	2.3
1	A	843	ASP	2.3
1	A	935	ILE	2.2
1	A	959	ILE	2.2
1	A	885	LEU	2.2
1	A	709	TRP	2.1
1	B	896	LEU	2.0
1	A	910	ARG	2.0
1	A	951	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ANP	B	301	31/31	0.97	0.17	-0.45	38,38,40,40	0
3	ANP	A	301	31/31	0.96	0.17	-0.90	44,45,48,48	0
2	MG	B	202	1/1	0.91	0.28	-	57,57,57,57	0
2	MG	A	202	1/1	0.87	0.32	-	49,49,49,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.