



wwPDB NMR Structure Validation Summary Report ⓘ

Feb 13, 2017 – 12:18 am GMT

PDB ID : 2LP4
Title : Solution structure of P1-CheY/P2 complex in bacterial chemotaxis
Authors : Dahlquist, F.; Mo, G.; Zhou, H.; Kamamura, T.
Deposited on : 2012-01-31

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	trunk28760
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 6%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	225	
2	Y	128	

2 Ensemble composition and analysis

This entry contains 25 models. Model 3 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:7-A:132 (126)	0.04	7
2	A:160-A:225, Y:2-Y:129 (194)	0.00	3

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 6, 10, 11, 12, 14, 21, 24
2	2, 5, 8, 17, 18, 19, 20, 22
3	3, 7, 9, 16, 23
4	4, 13, 15, 25

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5367 atoms, of which 2669 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Chemotaxis protein CheA.

Mol	Chain	Residues	Atoms						Trace
1	A	225	Total	C	H	N	O	S	0
			3439	1087	1695	284	365	8	

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	29	ASP	VAL	SEE REMARK 999	UNP P07363
A	31	VAL	GLN	SEE REMARK 999	UNP P07363
A	34	SER	ALA	SEE REMARK 999	UNP P07363
A	60	THR	SER	SEE REMARK 999	UNP P07363
A	61	ILE	VAL	SEE REMARK 999	UNP P07363
A	107	ASN	GLN	SEE REMARK 999	UNP P07363
A	109	GLU	GLN	SEE REMARK 999	UNP P07363
A	117	GLU	ASP	SEE REMARK 999	UNP P07363
A	121	ASN	GLN	SEE REMARK 999	UNP P07363

- Molecule 2 is a protein called Chemotaxis protein CheY.


Mol	Chain	Residues	Atoms						Trace
2	Y	128	Total	C	H	N	O	S	0
			1928	612	974	154	182	6	

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Chemotaxis protein CheA




- Molecule 2: Chemotaxis protein CheY



4.2 Residue scores for the representative (medoid) model from the NMR ensemble


The representative model is number 3. Colouring as in section 4.1 above.

- Molecule 1: Chemotaxis protein CheA





● Molecule 2: Chemotaxis protein CheY



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 200 calculated structures, 25 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
XPLOR-NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2lp4_cs.str
Number of chemical shift lists	1
Total number of shifts	216
Number of shifts mapped to atoms	216
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	6%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.94±0.00	8±0/1517 (0.5±0.0%)	1.11±0.00	17±0/2054 (0.8±0.0%)
2	Y	1.37±0.00	8±0/966 (0.8±0.0%)	1.58±0.00	15±0/1302 (1.2±0.0%)
All	All	1.12	400/62075 (0.6%)	1.31	800/83900 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.3±0.7
2	Y	0.0±0.0	0.8±0.4
All	All	0	54

5 of 17 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	178	GLU	CD-OE1	6.55	1.32	1.25	1	25
2	Y	67	GLU	CD-OE1	6.46	1.32	1.25	1	25
1	A	23	MET	CG-SD	6.24	1.97	1.81	8	25
2	Y	37	GLU	CD-OE1	6.16	1.32	1.25	12	15
2	Y	37	GLU	CD-OE2	6.14	1.32	1.25	22	10

5 of 35 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	Y	18	ARG	NE-CZ-NH1	8.12	124.36	120.30	15	16
2	Y	18	ARG	NE-CZ-NH2	8.00	124.30	120.30	4	9
2	Y	38	ASP	CB-CG-OD1	7.65	125.18	118.30	1	25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	Y	3	ASP	CB-CG-OD1	-7.31	111.72	118.30	1	25
2	Y	57	ASP	CB-CG-OD1	-6.71	112.26	118.30	8	13

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
2	Y	18	ARG	Sidechain	21
1	A	45	ARG	Sidechain	20
1	A	7	ASP	Peptide	13

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1497	1452	1448	62±9
2	Y	954	974	957	96±9
All	All	61275	60650	60121	3701

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

5 of 779 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:7:ASP:HB3	1:A:8:PHE:CD1	1.59	1.26	6	3
1:A:7:ASP:CB	1:A:8:PHE:CE1	1.51	1.90	6	2
1:A:7:ASP:HB3	1:A:8:PHE:CE1	1.45	1.44	6	3
2:Y:17:MET:O	2:Y:21:VAL:HG22	1.26	1.25	12	2
2:Y:17:MET:O	2:Y:21:VAL:CG2	1.23	1.85	14	3

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	191/225 (85%)	183±1 (96±0%)	7±1 (4±0%)	1±1 (0±0%)	42	80
2	Y	126/128 (98%)	119±0 (94±0%)	6±0 (5±0%)	1±0 (1±0%)	27	73
All	All	7925/8825 (90%)	7554 (95%)	325 (4%)	46 (1%)	33	77

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	Y	79	SER	25
1	A	7	ASP	10
1	A	160	ARG	7
1	A	132	GLY	4

6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/193 (83%)	143±2 (89±1%)	17±2 (11±1%)	12	55
2	Y	95/102 (93%)	79±2 (83±2%)	16±2 (17±2%)	5	41
All	All	6375/7375 (86%)	5526 (87%)	849 (13%)	8	49

5 of 60 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	203	ILE	25
2	Y	60	MET	25
2	Y	78	MET	25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
2	Y	119	LYS	25
1	A	131	LYS	25

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 6% for the well-defined parts and 5% for the entire structure.

7.1 Chemical shift list 1

File name: 2lp4_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	216
Number of shifts mapped to atoms	216
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	0	—	None (insufficient data)
$^{13}\text{C}_\beta$	0	—	None (insufficient data)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	108	0.15 ± 0.36	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 6%, i.e. 216 atoms were assigned a chemical shift out of a possible 3830. 0 out of 58 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	216/1586 (14%)	108/633 (17%)	0/640 (0%)	108/313 (35%)
Sidechain	0/2020 (0%)	0/1163 (0%)	0/782 (0%)	0/75 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	0/224 (0%)	0/122 (0%)	0/97 (0%)	0/5 (0%)
Overall	216/3830 (6%)	108/1918 (6%)	0/1519 (0%)	108/393 (27%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain Y:

