



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 15, 2018 – 06:10 PM EST

PDB ID : 5LQD  
Title : Trehalose-6-phosphate synthase, GDP-glucose-dependent OtsA  
Authors : Miah, F.; Asencion Diez, M.D.; Stevenson, C.E.M.; Lawson, D.M.; Iglesias, A.A.; Bornemann, S.  
Deposited on : 2016-08-16  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030736  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030736


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	465	<div> <div>3%</div> <div>92%</div> <div>7%</div> </div>
1	B	465	<div> <div>4%</div> <div>89%</div> <div>7%</div> </div>
1	C	465	<div> <div>3%</div> <div>93%</div> <div>7%</div> </div>
1	D	465	<div> <div>4%</div> <div>93%</div> <div>7%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria.

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	B	1001	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14466 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha,alpha-trehalose-phosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	446	Total	C	N	O	S	0	3	0
			3474	2182	636	648	8			
1	B	432	Total	C	N	O	S	0	1	0
			3354	2112	615	619	8			
1	C	446	Total	C	N	O	S	0	0	0
			3446	2168	625	645	8			
1	D	447	Total	C	N	O	S	0	3	0
			3476	2185	634	649	8			

There are 56 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MET	-	initiating methionine	UNP F2RH01
A	-11	HIS	-	expression tag	UNP F2RH01
A	-10	HIS	-	expression tag	UNP F2RH01
A	-9	HIS	-	expression tag	UNP F2RH01
A	-8	HIS	-	expression tag	UNP F2RH01
A	-7	HIS	-	expression tag	UNP F2RH01
A	-6	HIS	-	expression tag	UNP F2RH01
A	-5	GLU	-	expression tag	UNP F2RH01
A	-4	ASN	-	expression tag	UNP F2RH01
A	-3	LEU	-	expression tag	UNP F2RH01
A	-2	TYR	-	expression tag	UNP F2RH01
A	-1	PHE	-	expression tag	UNP F2RH01
A	0	GLN	-	expression tag	UNP F2RH01
A	1	GLY	-	expression tag	UNP F2RH01
B	-12	MET	-	initiating methionine	UNP F2RH01
B	-11	HIS	-	expression tag	UNP F2RH01
B	-10	HIS	-	expression tag	UNP F2RH01
B	-9	HIS	-	expression tag	UNP F2RH01
B	-8	HIS	-	expression tag	UNP F2RH01
B	-7	HIS	-	expression tag	UNP F2RH01
B	-6	HIS	-	expression tag	UNP F2RH01

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	GLU	-	expression tag	UNP F2RH01
B	-4	ASN	-	expression tag	UNP F2RH01
B	-3	LEU	-	expression tag	UNP F2RH01
B	-2	TYR	-	expression tag	UNP F2RH01
B	-1	PHE	-	expression tag	UNP F2RH01
B	0	GLN	-	expression tag	UNP F2RH01
B	1	GLY	-	expression tag	UNP F2RH01
C	-12	MET	-	initiating methionine	UNP F2RH01
C	-11	HIS	-	expression tag	UNP F2RH01
C	-10	HIS	-	expression tag	UNP F2RH01
C	-9	HIS	-	expression tag	UNP F2RH01
C	-8	HIS	-	expression tag	UNP F2RH01
C	-7	HIS	-	expression tag	UNP F2RH01
C	-6	HIS	-	expression tag	UNP F2RH01
C	-5	GLU	-	expression tag	UNP F2RH01
C	-4	ASN	-	expression tag	UNP F2RH01
C	-3	LEU	-	expression tag	UNP F2RH01
C	-2	TYR	-	expression tag	UNP F2RH01
C	-1	PHE	-	expression tag	UNP F2RH01
C	0	GLN	-	expression tag	UNP F2RH01
C	1	GLY	-	expression tag	UNP F2RH01
D	-12	MET	-	initiating methionine	UNP F2RH01
D	-11	HIS	-	expression tag	UNP F2RH01
D	-10	HIS	-	expression tag	UNP F2RH01
D	-9	HIS	-	expression tag	UNP F2RH01
D	-8	HIS	-	expression tag	UNP F2RH01
D	-7	HIS	-	expression tag	UNP F2RH01
D	-6	HIS	-	expression tag	UNP F2RH01
D	-5	GLU	-	expression tag	UNP F2RH01
D	-4	ASN	-	expression tag	UNP F2RH01
D	-3	LEU	-	expression tag	UNP F2RH01
D	-2	TYR	-	expression tag	UNP F2RH01
D	-1	PHE	-	expression tag	UNP F2RH01
D	0	GLN	-	expression tag	UNP F2RH01
D	1	GLY	-	expression tag	UNP F2RH01

- Molecule 2 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
2	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
2	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
2	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	169	Total O 169 169	0	0
4	B	151	Total O 151 151	0	0
4	C	163	Total O 163 163	0	0
4	D	169	Total O 169 169	0	0







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	41.43 Å 168.35 Å 133.92 Å 90.00° 97.19° 90.00°	Depositor
Resolution (Å)	32.64 – 1.95 32.64 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.6 (32.64-1.95) 99.6 (32.64-1.95)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.65 (at 1.95 Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.197 , 0.238 0.205 , 0.246	Depositor DCC
$R_{free}$ test set	6621 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.1	Xtriage
Anisotropy	0.573	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 37.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.047 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	14466	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.62	0/3555	0.79	6/4844 (0.1%)
1	B	0.62	0/3434	0.80	3/4681 (0.1%)
1	C	0.61	0/3524	0.82	4/4803 (0.1%)
1	D	0.63	0/3563	0.83	6/4857 (0.1%)
All	All	0.62	0/14076	0.81	19/19185 (0.1%)

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	361	ASP	CB-CG-OD1	8.43	125.89	118.30
1	D	263	ARG	NE-CZ-NH1	7.70	124.15	120.30
1	D	113	ARG	NE-CZ-NH2	-7.38	116.61	120.30
1	A	272	ARG	NE-CZ-NH1	7.15	123.87	120.30
1	D	263	ARG	NE-CZ-NH2	-6.28	117.16	120.30
1	A	272	ARG	NE-CZ-NH2	-6.07	117.27	120.30
1	B	122	ARG	NE-CZ-NH1	6.04	123.32	120.30
1	B	272	ARG	NE-CZ-NH1	6.00	123.30	120.30
1	D	113	ARG	NE-CZ-NH1	5.91	123.25	120.30
1	D	272	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	A	188	ARG	NE-CZ-NH1	5.56	123.08	120.30
1	C	290	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	A	278	ARG	NE-CZ-NH1	5.35	122.98	120.30
1	A	69	ARG	NE-CZ-NH1	5.34	122.97	120.30
1	D	221	ARG	NE-CZ-NH1	5.34	122.97	120.30
1	B	118	ARG	NE-CZ-NH2	-5.28	117.66	120.30
1	A	308	ARG	NE-CZ-NH1	5.16	122.88	120.30
1	C	361	ASP	CB-CG-OD2	-5.12	113.69	118.30
1	C	301	ARG	NE-CZ-NH1	5.08	122.84	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3474	0	3405	5	0
1	B	3354	0	3276	5	0
1	C	3446	0	3371	7	0
1	D	3476	0	3415	4	0
2	A	12	0	13	0	0
2	B	12	0	13	0	0
2	C	12	0	13	0	0
2	D	12	0	13	0	0
3	A	4	0	6	0	0
3	B	4	0	6	0	0
3	C	4	0	6	0	0
3	D	4	0	6	0	0
4	A	169	0	0	0	0
4	B	151	0	0	0	0
4	C	163	0	0	0	0
4	D	169	0	0	0	0
All	All	14466	0	13543	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:56:ARG:NH1	1:C:72:ASP:OD1	2.32	0.63
1:A:229:ALA:HB2	1:A:434:THR:HG21	1.86	0.58
1:B:368:LYS:HB3	1:B:433:ALA:HB1	1.85	0.57
1:A:13:VAL:HG21	1:A:65:GLU:HG3	1.88	0.56
1:C:39:VAL:O	1:C:39:VAL:CG1	2.54	0.55
1:C:39:VAL:O	1:C:39:VAL:HG12	2.07	0.54
1:D:368:LYS:HB3	1:D:433:ALA:HB1	1.95	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:174:TYR:CZ	1:D:177:MET:HE1	2.49	0.47
1:C:36:LEU:HD23	1:C:442:PHE:CD2	2.49	0.47
1:C:361:ASP:OD2	1:C:364:ASN:HB2	2.16	0.45
1:B:13:VAL:HG21	1:B:65:GLU:HG3	1.98	0.45
1:B:229:ALA:HB2	1:B:434:THR:HG21	1.98	0.44
1:C:96:HIS:HA	1:C:174:TYR:CE2	2.53	0.43
1:A:4:VAL:O	1:A:44:SER:HA	2.19	0.43
1:A:361:ASP:OD2	1:A:364:ASN:HB2	2.19	0.42
1:D:230:GLU:N	1:D:230:GLU:OE1	2.52	0.42
1:A:368:LYS:HB3	1:A:433:ALA:HB1	2.00	0.42
1:C:215:GLN:O	1:C:215:GLN:HG2	2.19	0.42
1:D:215[A]:GLN:HB2	1:D:215[A]:GLN:HE21	1.68	0.41
1:B:160:ARG:HD2	1:B:449:LEU:O	2.21	0.41
1:B:84:GLY:O	1:B:88:SER:HB2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	445/465 (96%)	433 (97%)	12 (3%)	0	100	100
1	B	427/465 (92%)	411 (96%)	16 (4%)	0	100	100
1	C	442/465 (95%)	430 (97%)	10 (2%)	2 (0%)	32	19
1	D	446/465 (96%)	431 (97%)	13 (3%)	2 (0%)	38	25
All	All	1760/1860 (95%)	1705 (97%)	51 (3%)	4 (0%)	51	41

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	142	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	142	ASP
1	C	215	GLN
1	D	30	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	352/366 (96%)	350 (99%)	2 (1%)	89	88
1	B	336/366 (92%)	330 (98%)	6 (2%)	64	57
1	C	348/366 (95%)	345 (99%)	3 (1%)	82	81
1	D	353/366 (96%)	347 (98%)	6 (2%)	66	60
All	All	1389/1464 (95%)	1372 (99%)	17 (1%)	75	72

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	262	ASP
1	B	17	ARG
1	B	27	ARG
1	B	142	ASP
1	B	262	ASP
1	B	329	ASP
1	B	361	ASP
1	C	53	GLU
1	C	262	ASP
1	C	361	ASP
1	D	69	ARG
1	D	215[A]	GLN
1	D	215[B]	GLN
1	D	262	ASP
1	D	329	ASP
1	D	338	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	141	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	MES	A	1000	-	12,12,12	0.65	0	14,16,16	0.42	0
3	EDO	A	1001	-	3,3,3	0.64	0	2,2,2	0.22	0
2	MES	B	1000	-	12,12,12	0.65	0	14,16,16	0.48	0
3	EDO	B	1001	-	3,3,3	0.70	0	2,2,2	0.21	0
2	MES	C	1000	-	12,12,12	0.65	0	14,16,16	0.48	0
3	EDO	C	1001	-	3,3,3	0.62	0	2,2,2	0.22	0
2	MES	D	1000	-	12,12,12	0.72	0	14,16,16	0.52	0
3	EDO	D	1001	-	3,3,3	0.49	0	2,2,2	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MES	A	1000	-	-	0/6/14/14	0/1/1/1
3	EDO	A	1001	-	-	0/1/1/1	0/0/0/0
2	MES	B	1000	-	-	0/6/14/14	0/1/1/1
3	EDO	B	1001	-	-	0/1/1/1	0/0/0/0
2	MES	C	1000	-	-	0/6/14/14	0/1/1/1
3	EDO	C	1001	-	-	0/1/1/1	0/0/0/0
2	MES	D	1000	-	-	0/6/14/14	0/1/1/1
3	EDO	D	1001	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	446/465 (95%)	0.21	15 (3%)	46 56	22, 31, 51, 62	0
1	B	432/465 (92%)	0.30	20 (4%)	33 44	21, 35, 65, 90	0
1	C	446/465 (95%)	0.18	13 (2%)	52 62	20, 33, 56, 68	0
1	D	447/465 (96%)	0.22	19 (4%)	36 47	20, 31, 53, 67	0
All	All	1771/1860 (95%)	0.23	67 (3%)	41 51	20, 32, 56, 90	0

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	1	GLY	5.6
1	B	328	ALA	5.5
1	D	63	ILE	4.7
1	A	66	PRO	4.3
1	D	64	GLY	4.0
1	B	16	VAL	3.9
1	C	420	PRO	3.9
1	A	64	GLY	3.8
1	D	15	TYR	3.8
1	D	24	ASP	3.6
1	D	66	PRO	3.5
1	B	216	PRO	3.5
1	B	42	GLN	3.5
1	A	24	ASP	3.4
1	C	29	GLY	3.4
1	D	60	ARG	3.4
1	A	63	ILE	3.3
1	B	247	ARG	3.3
1	B	41	SER	3.2
1	A	25	ALA	3.1
1	B	239	PRO	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	60	ARG	2.9
1	B	213	GLY	2.9
1	A	1	GLY	2.9
1	D	28	GLY	2.8
1	D	42	GLN	2.8
1	D	23	LEU	2.7
1	A	23	LEU	2.7
1	D	61	ARG	2.7
1	A	62	GLY	2.7
1	B	250	GLU	2.6
1	C	1	GLY	2.6
1	C	140	VAL	2.6
1	B	28	GLY	2.5
1	D	62	GLY	2.5
1	B	43	ASP	2.5
1	A	26	ARG	2.4
1	C	76	ASP	2.4
1	B	420	PRO	2.4
1	B	61	ARG	2.3
1	A	61	ARG	2.3
1	B	329	ASP	2.3
1	D	328	ALA	2.2
1	B	1	GLY	2.2
1	C	214	GLU	2.2
1	D	167	THR	2.2
1	D	26	ARG	2.2
1	D	143	TYR	2.2
1	C	333	VAL	2.2
1	C	356	VAL	2.2
1	D	16	VAL	2.2
1	C	328	ALA	2.1
1	A	42	GLN	2.1
1	B	240	GLN	2.1
1	A	164	PHE	2.1
1	A	16	VAL	2.1
1	D	243	GLU	2.1
1	C	17	ARG	2.1
1	C	60	ARG	2.1
1	C	424	ASP	2.1
1	A	146	ALA	2.0
1	A	328	ALA	2.0
1	B	423	ALA	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	213	GLY	2.0
1	B	359	VAL	2.0
1	B	440	ARG	2.0
1	D	25	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	B	1001	4/4	0.91	0.14	2.19	29,31,32,37	0
3	EDO	D	1001	4/4	0.93	0.11	-0.11	23,26,27,28	0
2	MES	B	1000	12/12	0.95	0.11	-0.15	26,33,38,39	0
3	EDO	A	1001	4/4	0.94	0.10	-0.36	31,32,34,35	0
2	MES	D	1000	12/12	0.96	0.10	-0.63	26,29,34,36	0
2	MES	A	1000	12/12	0.96	0.09	-0.84	28,33,36,41	0
2	MES	C	1000	12/12	0.97	0.09	-1.01	27,31,39,39	0
3	EDO	C	1001	4/4	0.95	0.08	-1.52	23,27,29,33	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.