



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 26, 2017 – 02:53 PM EDT

PDB ID : 3LSK
Title : Pyranose 2-oxidase T169S acetate complex
Authors : Tan, T.C.; Spadiut, O.; Divne, C.
Deposited on : unknown
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	623	<div> <div>2%</div> <div>79% 13% 8%</div> </div>
1	B	623	<div> <div>%</div> <div>75% 15% 8%</div> </div>
1	C	623	<div> <div>3%</div> <div>71% 19% 8%</div> </div>
1	D	623	<div> <div>4%</div> <div>76% 14% 8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	901	-	-	X	X
3	ACT	B	901	-	-	X	X
3	ACT	C	901	-	-	X	X
3	ACT	D	901	-	-	X	-
4	12P	A	903	-	-	-	X
4	12P	A	904	-	-	-	X
4	12P	B	902	-	-	-	X
4	12P	C	905	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20361 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pyranose 2-oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	576	Total	C	N	O	S	0	0	0
			4541	2867	777	872	25			
1	B	576	Total	C	N	O	S	0	0	0
			4541	2867	777	872	25			
1	C	575	Total	C	N	O	S	0	0	0
			4532	2862	776	870	24			
1	D	574	Total	C	N	O	S	0	0	0
			4525	2858	775	868	24			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	169	SER	THR	ENGINEERED	UNP Q7ZA32
B	169	SER	THR	ENGINEERED	UNP Q7ZA32
C	169	SER	THR	ENGINEERED	UNP Q7ZA32
D	169	SER	THR	ENGINEERED	UNP Q7ZA32

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	B	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	C	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	D	1	Total 53	C 27	N 9	O 15	P 2	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $\text{C}_2\text{H}_3\text{O}_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

- Molecule 4 is DODECAETHYLENE GLYCOL (three-letter code: 12P) (formula: $C_{24}H_{50}O_{13}$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 16 11 5	0	0
4	A	1	Total C O 16 11 5	0	0
4	B	1	Total C O 16 10 6	0	0
4	C	1	Total C O 14 10 4	0	0
4	D	1	Total C O 12 8 4	0	0
4	D	1	Total C O 11 7 4	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	518	Total 518	O 518	0	0
5	B	534	Total 534	O 534	0	0
5	C	457	Total 457	O 457	0	0
5	D	400	Total 400	O 400	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Pyranose 2-oxidase




• Molecule 1: Pyranose 2-oxidase



• Molecule 1: Pyranose 2-oxidase



• Molecule 1: Pyranose 2-oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	100.26Å 102.45Å 136.72Å 90.00° 91.13° 90.00°	Depositor
Resolution (Å)	30.00 – 1.95 29.29 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.8 (30.00-1.95) 98.8 (29.29-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.16	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, R_{free}	0.196 , 0.256 0.207 , 0.261	Depositor DCC
R_{free} test set	2008 reflections (1.02%)	DCC
Wilson B-factor (Å ²)	13.3	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.059 for -k,-h,-l 0.048 for k,h,-l 0.039 for h,-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	20361	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 12P, FAD, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.19	11/4657 (0.2%)	1.01	11/6331 (0.2%)
1	B	1.18	10/4657 (0.2%)	1.02	13/6331 (0.2%)
1	C	1.02	7/4648 (0.2%)	0.97	10/6320 (0.2%)
1	D	0.99	5/4641 (0.1%)	0.91	3/6310 (0.0%)
All	All	1.10	33/18603 (0.2%)	0.98	37/25292 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	3
1	D	0	2
All	All	0	6

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	166	THR	C-N	13.15	1.64	1.34
1	B	206	PHE	CE1-CZ	-7.30	1.23	1.37
1	B	316	TYR	CD2-CE2	-7.16	1.28	1.39
1	C	322	ALA	CA-CB	-6.94	1.37	1.52
1	A	429	GLU	CB-CG	6.89	1.65	1.52

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	81	ASP	CB-CG-OD1	-10.00	109.30	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	167	HIS	O-C-N	-8.50	109.10	122.70
1	A	147	ASP	CB-CG-OD1	7.44	124.99	118.30
1	C	470	ASP	CB-CG-OD1	7.38	124.94	118.30
1	B	307	ASP	CB-CG-OD1	7.14	124.73	118.30

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	449	ILE	Peptide
1	C	166	THR	Mainchain
1	C	167	HIS	Mainchain
1	C	457	GLY	Peptide
1	D	166	THR	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4541	0	4386	56	0
1	B	4541	0	4386	77	0
1	C	4532	0	4380	96	0
1	D	4525	0	4373	93	0
2	A	53	0	27	1	0
2	B	53	0	26	0	0
2	C	53	0	29	14	0
2	D	53	0	26	1	0
3	A	4	0	3	3	0
3	B	4	0	3	2	0
3	C	4	0	3	2	0
3	D	4	0	3	2	0
4	A	32	0	36	0	0
4	B	16	0	21	0	0
4	C	14	0	16	1	0
4	D	23	0	26	9	0
5	A	518	0	0	9	0
5	B	534	0	0	21	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	457	0	0	12	0
5	D	400	0	0	19	0
All	All	20361	0	17744	321	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 321 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:167:HIS:HE2	2:C:801:FAD:C8M	1.00	1.58
1:C:167:HIS:NE2	2:C:801:FAD:HM82	1.04	1.35
1:B:312:LYS:HG3	5:B:2396:HOH:O	1.41	1.19
1:A:385:THR:O	1:A:388:GLU:HG2	1.38	1.18
1:C:299:HIS:CE1	1:C:310:GLU:HG2	1.78	1.17

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	574/623 (92%)	551 (96%)	22 (4%)	1 (0%)	51 41
1	B	574/623 (92%)	553 (96%)	20 (4%)	1 (0%)	51 41
1	C	573/623 (92%)	545 (95%)	26 (4%)	2 (0%)	44 33
1	D	572/623 (92%)	546 (96%)	25 (4%)	1 (0%)	51 41
All	All	2293/2492 (92%)	2195 (96%)	93 (4%)	5 (0%)	51 41

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	82	SER
1	B	82	SER
1	C	458	ALA
1	C	459	VAL
1	D	82	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	504/542 (93%)	489 (97%)	15 (3%)	46	34
1	B	504/542 (93%)	486 (96%)	18 (4%)	40	26
1	C	503/542 (93%)	479 (95%)	24 (5%)	30	14
1	D	502/542 (93%)	489 (97%)	13 (3%)	51	40
All	All	2013/2168 (93%)	1943 (96%)	70 (4%)	41	27

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	460	GLN
1	C	185	LYS
1	D	400	SER
1	B	593	ASN
1	C	91	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	344	ASN
1	C	324	HIS
1	D	461	GLN
1	C	132	GLN
1	C	299	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	801	1	51,58,58	1.73	11 (21%)	54,89,89	2.56	17 (31%)
3	ACT	A	901	-	1,3,3	2.62	1 (100%)	0,3,3	0.00	-
4	12P	A	903	-	15,15,36	0.69	0	14,14,35	0.51	0
4	12P	A	904	-	15,15,36	0.71	0	14,14,35	0.43	0
2	FAD	B	801	1	51,58,58	1.57	9 (17%)	54,89,89	4.04	14 (25%)
3	ACT	B	901	-	1,3,3	2.05	1 (100%)	0,3,3	0.00	-
4	12P	B	902	-	15,15,36	0.48	0	14,14,35	0.43	0
2	FAD	C	801	1	51,58,58	1.41	6 (11%)	54,89,89	2.90	19 (35%)
3	ACT	C	901	-	1,3,3	3.05	1 (100%)	0,3,3	0.00	-
4	12P	C	905	-	13,13,36	0.77	0	12,12,35	0.31	0
2	FAD	D	801	1	51,58,58	1.51	8 (15%)	54,89,89	2.71	19 (35%)
3	ACT	D	901	-	1,3,3	1.78	0	0,3,3	0.00	-
4	12P	D	906	-	11,11,36	0.53	0	10,10,35	0.28	0
4	12P	D	907	-	10,10,36	0.63	0	9,9,35	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	801	1	-	0/28/50/50	0/6/6/6
3	ACT	A	901	-	-	0/0/0/0	0/0/0/0
4	12P	A	903	-	-	0/13/13/34	0/0/0/0
4	12P	A	904	-	-	0/13/13/34	0/0/0/0
2	FAD	B	801	1	-	0/28/50/50	0/6/6/6
3	ACT	B	901	-	-	0/0/0/0	0/0/0/0
4	12P	B	902	-	-	0/13/13/34	0/0/0/0
2	FAD	C	801	1	-	0/28/50/50	0/6/6/6
3	ACT	C	901	-	-	0/0/0/0	0/0/0/0
4	12P	C	905	-	-	0/11/11/34	0/0/0/0
2	FAD	D	801	1	-	0/28/50/50	0/6/6/6
3	ACT	D	901	-	-	0/0/0/0	0/0/0/0
4	12P	D	906	-	-	0/9/9/34	0/0/0/0
4	12P	D	907	-	-	0/8/8/34	0/0/0/0

The worst 5 of 37 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	FAD	C4A-N3A	-4.18	1.29	1.35
2	A	801	FAD	C2B-C1B	-4.09	1.47	1.53
2	A	801	FAD	O4B-C4B	-3.83	1.36	1.45
2	D	801	FAD	C2B-C1B	-3.47	1.48	1.53
2	B	801	FAD	PA-O1A	-2.93	1.39	1.50

The worst 5 of 69 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	FAD	N3A-C2A-N1A	-12.58	117.91	128.86
2	B	801	FAD	C4X-C4-N3	-12.13	106.20	123.48
2	A	801	FAD	N3A-C2A-N1A	-9.51	120.58	128.86
2	C	801	FAD	C4-C4X-C10	-8.84	112.81	119.96
2	D	801	FAD	N3A-C2A-N1A	-8.73	121.26	128.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 34 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	FAD	1	0
3	A	901	ACT	3	0
3	B	901	ACT	2	0
2	C	801	FAD	14	0
3	C	901	ACT	2	0
4	C	905	12P	1	0
2	D	801	FAD	1	0
3	D	901	ACT	2	0
4	D	907	12P	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	166:THR	C	167:HIS	N	1.64

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	576/623 (92%)	-0.02	11 (1%) 67 76	7, 13, 31, 43	0
1	B	576/623 (92%)	-0.05	9 (1%) 72 80	7, 13, 27, 48	0
1	C	575/623 (92%)	0.16	19 (3%) 47 58	9, 18, 36, 55	0
1	D	574/623 (92%)	0.28	25 (4%) 35 45	10, 22, 41, 51	0
All	All	2301/2492 (92%)	0.09	64 (2%) 53 64	7, 17, 35, 55	0

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	459	VAL	7.6
1	D	459	VAL	6.4
1	D	458	ALA	6.3
1	C	458	ALA	5.9
1	B	459	VAL	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	ACT	C	901	4/4	0.88	0.22	5.66	30,31,31,31	0
4	12P	B	902	16/37	0.93	0.13	2.94	22,24,39,42	0
3	ACT	A	901	4/4	0.95	0.15	2.67	29,31,32,33	0
3	ACT	B	901	4/4	0.93	0.17	2.64	27,27,27,29	0
4	12P	C	905	14/37	0.72	0.26	2.38	35,42,44,45	0
4	12P	A	903	16/37	0.91	0.13	2.29	21,26,38,38	0
4	12P	A	904	16/37	0.90	0.12	2.07	23,26,39,40	0
4	12P	D	907	11/37	0.75	0.26	1.52	45,47,48,49	0
3	ACT	D	901	4/4	0.91	0.14	0.98	26,27,28,28	0
4	12P	D	906	12/37	0.89	0.13	0.74	23,27,34,34	0
2	FAD	A	801	53/53	0.97	0.11	0.11	4,10,13,16	0
2	FAD	B	801	53/53	0.97	0.11	-0.50	2,9,14,18	0
2	FAD	D	801	53/53	0.95	0.11	-0.55	12,16,19,21	0
2	FAD	C	801	53/53	0.97	0.10	-0.75	7,14,18,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.