



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 1, 2017 – 02:43 PM EDT

PDB ID : 5LSG  
Title : PPARgamma complex with the betulinic acid  
Authors : Pochetti, G.; Montanari, R.; Capelli, D.; Loiodice, F.; Laghezza, A.; Calleri, E.; Paiardini, A.  
Deposited on : unknown  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	304	<div> <div>9%</div> <div>67%</div> <div>16%</div> <div>16%</div> </div>
1	B	304	<div> <div>16%</div> <div>68%</div> <div>16%</div> <div>14%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	QZQ	A	501	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4293 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisome proliferator-activated receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	15	0	0
			2047	1321	334	382	10			
1	B	260	Total	C	N	O	S	0	0	0
			2072	1339	338	385	10			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	174	MET	-	initiating methionine	UNP P37231
A	175	GLY	-	expression tag	UNP P37231
A	176	SER	-	expression tag	UNP P37231
A	177	SER	-	expression tag	UNP P37231
A	178	HIS	-	expression tag	UNP P37231
A	179	HIS	-	expression tag	UNP P37231
A	180	HIS	-	expression tag	UNP P37231
A	181	HIS	-	expression tag	UNP P37231
A	182	HIS	-	expression tag	UNP P37231
A	183	HIS	-	expression tag	UNP P37231
A	184	SER	-	expression tag	UNP P37231
A	185	SER	-	expression tag	UNP P37231
A	186	GLY	-	expression tag	UNP P37231
A	187	LEU	-	expression tag	UNP P37231
A	188	VAL	-	expression tag	UNP P37231
A	189	PRO	-	expression tag	UNP P37231
A	190	ARG	-	expression tag	UNP P37231
A	191	GLY	-	expression tag	UNP P37231
A	192	SER	-	expression tag	UNP P37231
A	193	HIS	-	expression tag	UNP P37231
A	194	MET	-	expression tag	UNP P37231
B	174	MET	-	initiating methionine	UNP P37231
B	175	GLY	-	expression tag	UNP P37231
B	176	SER	-	expression tag	UNP P37231
B	177	SER	-	expression tag	UNP P37231

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	178	HIS	-	expression tag	UNP P37231
B	179	HIS	-	expression tag	UNP P37231
B	180	HIS	-	expression tag	UNP P37231
B	181	HIS	-	expression tag	UNP P37231
B	182	HIS	-	expression tag	UNP P37231
B	183	HIS	-	expression tag	UNP P37231
B	184	SER	-	expression tag	UNP P37231
B	185	SER	-	expression tag	UNP P37231
B	186	GLY	-	expression tag	UNP P37231
B	187	LEU	-	expression tag	UNP P37231
B	188	VAL	-	expression tag	UNP P37231
B	189	PRO	-	expression tag	UNP P37231
B	190	ARG	-	expression tag	UNP P37231
B	191	GLY	-	expression tag	UNP P37231
B	192	SER	-	expression tag	UNP P37231
B	193	HIS	-	expression tag	UNP P37231
B	194	MET	-	expression tag	UNP P37231

- Molecule 2 is Betulinic Acid (three-letter code: QZQ) (formula:  $C_{30}H_{46}O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			33	30	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	87	Total 87	O 87	0	0
3	B	54	Total 54	O 54	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.86 Å 60.85 Å 118.03 Å 90.00° 102.48° 90.00°	Depositor
Resolution (Å)	45.68 – 2.00 45.68 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.5 (45.68-2.00) 98.5 (45.68-2.00)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.57 (at 2.00 Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, $R_{free}$	0.227 , 0.264 0.227 , 0.265	Depositor DCC
$R_{free}$ test set	2151 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.8	Xtriage
Anisotropy	0.507	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 56.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4293	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: QZQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.44	0/2080	0.58	0/2801
1	B	0.47	2/2107 (0.1%)	0.63	0/2841
All	All	0.46	2/4187 (0.0%)	0.60	0/5642

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	354	LYS	C-N	6.53	1.49	1.34
1	B	359	PRO	N-CD	5.48	1.55	1.47

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	357	ARG	Peptide
1	B	343	GLU	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2047	0	2111	43	0
1	B	2072	0	2121	33	0
2	A	33	0	0	4	0
3	A	87	0	0	6	0
3	B	54	0	0	5	0
All	All	4293	0	4232	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:243:ASP:OD2	3:A:601:HOH:O	1.80	0.99
1:A:240:LYS:HG2	1:A:241:THR:HG23	1.48	0.95
1:A:325:ILE:HG23	1:A:388:ILE:HD12	1.66	0.78
1:A:349:THR:HG22	1:A:352:PHE:H	1.49	0.77
1:A:298:GLU:OE2	3:A:602:HOH:O	2.04	0.74
1:A:238:THR:O	1:A:240:LYS:N	2.20	0.73
1:B:325:ILE:HD12	1:B:388:ILE:HG23	1.69	0.72
1:B:342:SER:OG	1:B:343:GLU:HG2	1.90	0.70
1:A:207:GLU:OE1	3:A:603:HOH:O	2.10	0.69
1:A:319:LYS:HZ3	1:A:474:LYS:HG2	1.58	0.69
1:B:313:ASP:OD2	3:B:502:HOH:O	2.10	0.68
1:A:357:ARG:HD3	1:A:358:LYS:H	1.59	0.68
1:B:240:LYS:O	1:B:242:THR:N	2.27	0.67
1:A:411:ASP:OD1	3:A:604:HOH:O	2.12	0.67
1:B:471:GLU:OE2	3:B:503:HOH:O	2.13	0.65
1:A:357:ARG:HD3	1:A:358:LYS:N	2.14	0.62
1:A:358:LYS:HD3	1:A:358:LYS:N	2.15	0.61
1:B:407:GLU:OE1	3:B:504:HOH:O	2.16	0.61
1:B:334:MET:CE	1:B:368:PHE:HA	2.33	0.58
1:B:327:TYR:CZ	1:B:367:LYS:HE3	2.39	0.57
1:A:357:ARG:HH21	1:A:358:LYS:HE2	1.69	0.57
1:A:341:ILE:HG22	2:A:501:QZQ:C05	2.34	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:360:PHE:C	1:B:362:ASP:H	2.08	0.57
1:B:276:GLU:HA	1:B:280:ARG:HH21	1.70	0.57
1:A:430:GLN:HG3	1:A:433:ALA:HB3	1.87	0.56
1:B:334:MET:HE1	1:B:368:PHE:HA	1.88	0.56
1:B:430:GLN:HG3	1:B:433:ALA:HB3	1.87	0.55
1:B:342:SER:O	1:B:344:GLY:N	2.39	0.53
1:B:380:ASP:OD1	1:B:382:SER:OG	2.22	0.53
1:A:437:GLN:O	1:A:440:THR:HG22	2.09	0.52
1:B:461:THR:O	1:B:463:MET:N	2.39	0.52
1:A:319:LYS:NZ	1:A:474:LYS:HG2	2.24	0.52
1:A:370:PHE:HB2	1:A:445:ILE:HD11	1.92	0.52
1:B:343:GLU:CB	1:B:345:GLN:H	2.22	0.51
1:A:404:LYS:NZ	1:A:408:ASP:OD2	2.34	0.51
1:A:288:ARG:NH1	1:A:291:GLU:HG2	2.26	0.51
1:B:258:GLY:O	1:B:262:ILE:N	2.41	0.51
1:A:240:LYS:HG2	1:A:241:THR:CG2	2.32	0.51
1:A:396:ASP:OD2	3:A:605:HOH:O	2.19	0.50
1:B:276:GLU:N	3:B:508:HOH:O	2.45	0.50
1:B:359:PRO:HB2	1:B:452:LEU:CD1	2.42	0.50
1:B:327:TYR:CE1	1:B:367:LYS:HE3	2.48	0.49
1:A:290:VAL:O	1:A:294:GLN:HG3	2.12	0.49
1:B:343:GLU:HB3	1:B:345:GLN:H	1.78	0.48
1:A:286:GLN:HG2	2:A:501:QZQ:C37	2.43	0.48
1:B:443:ARG:O	1:B:447:THR:HG23	2.14	0.48
1:A:325:ILE:HD11	1:A:392:ILE:CG1	2.45	0.47
1:A:350:ARG:NH2	1:A:365:GLU:OE2	2.38	0.47
1:A:357:ARG:NH2	1:A:358:LYS:HE2	2.29	0.47
1:A:244:LYS:N	3:A:609:HOH:O	2.33	0.46
1:B:402:ASN:O	1:B:405:PRO:HD2	2.16	0.46
1:A:433:ALA:O	1:A:437:GLN:HG3	2.16	0.46
1:A:358:LYS:H	1:A:358:LYS:HD3	1.79	0.46
1:B:247:PHE:N	1:B:345:GLN:O	2.47	0.45
1:A:323:HIS:HD2	2:A:501:QZQ:O16	2.00	0.44
1:B:360:PHE:C	1:B:362:ASP:N	2.71	0.44
1:A:207:GLU:HG3	1:A:209:ALA:H	1.84	0.43
1:A:325:ILE:HD11	1:A:392:ILE:HG13	2.00	0.43
2:A:501:QZQ:O1	2:A:501:QZQ:C20	2.66	0.43
1:B:343:GLU:HG3	1:B:345:GLN:HB2	2.00	0.43
1:A:212:ARG:HD2	1:A:212:ARG:HA	1.85	0.43
1:B:226:PHE:O	3:B:505:HOH:O	2.21	0.43
1:B:343:GLU:CB	1:B:345:GLN:HB2	2.49	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:226:PHE:HA	1:A:227:PRO:HD3	1.92	0.42
1:B:230:LYS:NZ	1:B:379:LEU:O	2.40	0.42
1:A:290:VAL:HG21	1:A:466:HIS:CD2	2.55	0.42
1:A:363:PHE:CZ	1:A:452:LEU:HG	2.54	0.42
1:B:359:PRO:O	1:B:362:ASP:HB2	2.20	0.42
1:A:252:MET:O	1:A:255:LEU:HB3	2.20	0.41
1:B:290:VAL:HG21	1:B:473:TYR:CD1	2.55	0.41
1:B:363:PHE:HB3	1:B:452:LEU:HD21	2.01	0.41
1:A:357:ARG:CD	1:A:358:LYS:H	2.31	0.41
1:B:334:MET:HE2	1:B:334:MET:HB3	1.79	0.41
1:A:370:PHE:CB	1:A:445:ILE:HD11	2.51	0.41
1:A:323:HIS:CE1	1:A:472:ILE:HG21	2.57	0.40
1:A:452:LEU:O	1:A:456:ILE:HD13	2.21	0.40
1:A:214:LEU:HD23	1:A:416:ALA:HB2	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/304 (83%)	244 (97%)	7 (3%)	1 (0%)	38	33
1	B	256/304 (84%)	240 (94%)	6 (2%)	10 (4%)	3	1
All	All	508/608 (84%)	484 (95%)	13 (3%)	11 (2%)	8	3

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	237	LEU
1	B	241	THR
1	B	242	THR
1	B	343	GLU

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	357	ARG
1	A	239	GLY
1	B	277	VAL
1	B	261	LYS
1	B	263	LYS
1	B	264	PHE
1	B	227	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/273 (84%)	227 (99%)	2 (1%)	82	87
1	B	230/273 (84%)	225 (98%)	5 (2%)	57	60
All	All	459/546 (84%)	452 (98%)	7 (2%)	70	74

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	245	SER
1	A	256	MET
1	B	254	SER
1	B	268	THR
1	B	341	ILE
1	B	357	ARG
1	B	358	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	451	GLN
1	B	466	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	QZQ	A	501	-	34,37,37	2.29	3 (8%)	54,64,64	3.23	24 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QZQ	A	501	-	-	0/4/99/99	0/5/5/5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	QZQ	C12-C11	-3.03	1.45	1.50
2	A	501	QZQ	C15-C20	-2.25	1.49	1.53
2	A	501	QZQ	C26-C25	11.79	1.51	1.32

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	QZQ	C12-C21-C20	-7.66	102.38	112.25
2	A	501	QZQ	C24-C26-C25	-6.34	114.66	122.79
2	A	501	QZQ	C02-C12-C11	-4.62	102.47	112.80
2	A	501	QZQ	C02-C07-C01	-4.36	99.35	106.20
2	A	501	QZQ	C07-C01-C21	-4.07	98.94	103.39
2	A	501	QZQ	C15-C13-C14	-3.87	103.51	112.44
2	A	501	QZQ	C34-C19-C3	-3.46	101.70	107.90
2	A	501	QZQ	C05-C23-C18	-3.18	101.94	111.62
2	A	501	QZQ	C06-C23-C05	-3.07	103.02	107.72
2	A	501	QZQ	C2-C12-C21	-3.06	102.20	109.55
2	A	501	QZQ	C21-C01-C04	-2.54	113.14	117.17
2	A	501	QZQ	C02-C12-C21	-2.17	98.69	101.18
2	A	501	QZQ	C15-C20-C19	2.20	113.60	110.97
2	A	501	QZQ	C13-C15-C20	2.36	115.75	111.86
2	A	501	QZQ	C19-C22-C14	2.41	111.75	108.35
2	A	501	QZQ	C15-C20-C21	2.46	117.81	113.63
2	A	501	QZQ	C19-C20-C21	3.97	116.25	111.38
2	A	501	QZQ	C23-C18-C17	4.01	121.38	114.80
2	A	501	QZQ	C22-C14-C17	4.45	122.75	114.93
2	A	501	QZQ	C12-C21-C01	4.60	111.56	105.23
2	A	501	QZQ	C3-C19-C22	5.51	116.18	110.77
2	A	501	QZQ	C27-C28-C18	5.52	122.03	111.08
2	A	501	QZQ	C28-C27-C22	6.00	123.22	112.80
2	A	501	QZQ	C01-C21-C20	10.71	131.62	120.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	QZQ	4	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	256/304 (84%)	0.64	27 (10%)	7 7	33, 51, 78, 104	3 (1%)
1	B	260/304 (85%)	1.12	50 (19%)	1 1	35, 55, 97, 120	0
All	All	516/608 (84%)	0.88	77 (14%)	3 3	33, 53, 91, 120	3 (0%)

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	465	LEU	13.8
1	B	363	PHE	9.6
1	B	238	THR	7.5
1	B	463	MET	7.4
1	B	464	SER	7.4
1	B	462	ASP	7.2
1	A	262	ILE	7.0
1	A	239	GLY	6.4
1	B	242	THR	6.4
1	B	360	PHE	5.9
1	B	241	THR	5.6
1	B	466	HIS	5.4
1	B	240	LYS	5.3
1	B	243	ASP	5.1
1	B	209	ALA	4.6
1	B	277	VAL	4.6
1	B	454	GLN	4.5
1	B	459	THR	4.3
1	A	243	ASP	4.2
1	B	354	LYS	4.1
1	B	207	GLU	4.0
1	A	256	MET	3.9
1	B	267	ILE	3.8
1	B	244	LYS	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	455	VAL	3.7
1	B	460	GLU	3.5
1	B	458	LYS	3.5
1	B	260	ASP	3.3
1	B	461	THR	3.3
1	A	263	LYS	3.2
1	A	359	PRO	3.2
1	B	256	MET	3.1
1	B	276	GLU	3.1
1	A	241	THR	3.1
1	B	352	PHE	3.0
1	A	252	MET	3.0
1	B	342	SER	2.9
1	A	282	PHE	2.8
1	B	427	GLU	2.8
1	B	211	LEU	2.8
1	A	280	ARG	2.8
1	B	456	ILE	2.8
1	B	451	GLN	2.7
1	A	362	ASP	2.6
1	A	363	PHE	2.6
1	A	261	LYS	2.6
1	B	322	VAL	2.6
1	A	357	ARG	2.5
1	B	359	PRO	2.5
1	A	299	TYR	2.5
1	B	358	LYS	2.4
1	B	326	ILE	2.4
1	B	262	ILE	2.4
1	B	391	ILE	2.4
1	A	277	VAL	2.3
1	A	361	GLY	2.3
1	B	252	MET	2.3
1	B	343	GLU	2.3
1	A	306	PHE	2.2
1	A	465	LEU	2.2
1	B	345	GLN	2.2
1	A	358	LYS	2.2
1	B	445	ILE	2.1
1	A	463	MET	2.1
1	A	260	ASP	2.1
1	B	328	THR	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	240	LYS	2.1
1	A	279	ILE	2.1
1	B	469	LEU	2.1
1	B	239	GLY	2.1
1	A	360	PHE	2.1
1	B	353	LEU	2.1
1	A	473	TYR	2.0
1	B	452	LEU	2.0
1	A	385	ALA	2.0
1	B	357	ARG	2.0
1	B	318	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	QZQ	A	501	33/33	0.55	0.71	11.25	59,64,68,69	33

## 6.5 Other polymers [i](#)

There are no such residues in this entry.