



# wwPDB NMR Structure Validation Summary Report ⓘ

Feb 13, 2017 – 02:10 am GMT

PDB ID : 2LUJ  
Title : Solution structure of a parallel-stranded oligoisoguanine DNA pentaplex formed by d(T(iG)4T) in the presence of Cs ions  
Authors : Kang, M.; Heuberger, B.; Chaput, J.C.; Switzer, C.; Feigon, J.  
Deposited on : 2012-06-14

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	trunk28760
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 69%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	6	 17% 50% 33%
1	B	6	 17% 50% 33%
1	C	6	 17% 50% 33%
1	D	6	 17% 50% 33%
1	E	6	 17% 50% 33%

## 2 Ensemble composition and analysis ⓘ

This entry contains 16 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

### 3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 975 atoms, of which 350 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called DNA (5'-D(\*TP\*(IGU)P\*(IGU)P\*(IGU)P\*(IGU)P\*T)-3').


Mol	Chain	Residues	Atoms						Trace
1	A	6	Total 195	C 60	H 70	N 24	O 36	P 5	0
1	B	6	Total 195	C 60	H 70	N 24	O 36	P 5	0
1	C	6	Total 195	C 60	H 70	N 24	O 36	P 5	0
1	D	6	Total 195	C 60	H 70	N 24	O 36	P 5	0
1	E	6	Total 195	C 60	H 70	N 24	O 36	P 5	0

## 4 Residue-property plots [i](#)


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: DNA (5'-D(\*TP\*(IGU)P\*(IGU)P\*(IGU)P\*(IGU)P\*T)-3')




- Molecule 1: DNA (5'-D(\*TP\*(IGU)P\*(IGU)P\*(IGU)P\*(IGU)P\*T)-3')




- Molecule 1: DNA (5'-D(\*TP\*(IGU)P\*(IGU)P\*(IGU)P\*(IGU)P\*T)-3')



- Molecule 1: DNA (5'-D(\*TP\*(IGU)P\*(IGU)P\*(IGU)P\*(IGU)P\*T)-3')




- Molecule 1: DNA (5'-D(\*TP\*(IGU)P\*(IGU)P\*(IGU)P\*(IGU)P\*T)-3')



## 4.2 Residue scores for the representative (author defined) model from the NMR ensemble


The representative model is number 1. Colouring as in section 4.1 above.

- Molecule 1: DNA (5'-D(\*TP\*(IGU)P\*(IGU)P\*(IGU)P\*(IGU)P\*T)-3')

Chain A: 


T1 G2 G3 G4 G5 T6

- Molecule 1: DNA (5'-D(\*TP\*(IGU)P\*(IGU)P\*(IGU)P\*(IGU)P\*T)-3')

Chain B: 


T7 G8 G9 G10 G11 T12

- Molecule 1: DNA (5'-D(\*TP\*(IGU)P\*(IGU)P\*(IGU)P\*(IGU)P\*T)-3')

Chain C: 


T13 G14 G15 G16 G17 T18

- Molecule 1: DNA (5'-D(\*TP\*(IGU)P\*(IGU)P\*(IGU)P\*(IGU)P\*T)-3')

Chain D: 

T19 G20 G21 G22 G23 T24

- Molecule 1: DNA (5'-D(\*TP\*(IGU)P\*(IGU)P\*(IGU)P\*(IGU)P\*T)-3')

Chain E: 

T25 G26 G27 G28 G29 T30

## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 16 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2luj_cs.str
Number of chemical shift lists	1
Total number of shifts	456
Number of shifts mapped to atoms	155
Number of unparsed shifts	0
Number of shifts with mapping errors	301
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	69%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IGU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.39±0.01	0±0/39 (0.0±0.0%)	2.60±0.02	6±1/56 (10.8±1.0%)
1	B	1.41±0.02	0±0/39 (0.0±0.0%)	2.61±0.01	6±0/56 (10.8±0.4%)
1	C	1.44±0.05	0±0/39 (0.0±0.0%)	2.62±0.03	6±0/56 (11.4±0.9%)
1	D	1.39±0.02	0±0/39 (0.0±0.0%)	2.60±0.01	6±0/56 (11.5±0.9%)
1	E	1.40±0.02	0±0/39 (0.0±0.0%)	2.59±0.02	6±0/56 (10.6±0.8%)
All	All	1.41	0/3120 (0.0%)	2.61	494/4480 (11.0%)

There are no bond-length outliers.

5 of 35 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	E	25	DT	C6-C5-C7	-6.84	118.79	122.90	8	16
1	C	13	DT	C6-C5-C7	-6.79	118.82	122.90	11	16
1	A	1	DT	C6-C5-C7	-6.79	118.82	122.90	4	16
1	B	7	DT	C6-C5-C7	-6.73	118.86	122.90	1	16
1	D	19	DT	C6-C5-C7	-6.69	118.89	122.90	12	16

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.



Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	125	70	70	20±1
1	B	125	70	70	17±2
1	C	125	70	70	16±3
1	D	125	70	70	21±2
1	E	125	70	70	20±2
All	All	10000	5600	5600	1021

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 65.

5 of 113 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:6:DT:O4	1:B:12:DT:H1'	1.00	1.56	11	16
1:A:6:DT:H1'	1:E:30:DT:O4	0.98	1.57	7	16
1:D:24:DT:O4	1:E:30:DT:H1'	0.98	1.59	1	16
1:C:18:DT:O4	1:D:24:DT:H1'	0.97	1.59	7	16
1:B:12:DT:O4	1:C:18:DT:H1'	0.97	1.59	6	8

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

20 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard

deviations the observed value is removed from the expected value. A bond length with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	IGU	A	2	1	16,24,25	1.11±0.01	0±0 (0±0%)
1	IGU	A	3	1	16,24,25	1.08±0.01	0±0 (0±0%)
1	IGU	A	4	1	16,24,25	1.09±0.01	0±0 (0±0%)
1	IGU	A	5	1	16,24,25	1.06±0.01	0±0 (0±0%)
1	IGU	B	10	1	16,24,25	1.09±0.01	0±0 (0±0%)
1	IGU	B	11	1	16,24,25	1.07±0.01	0±0 (0±0%)
1	IGU	B	8	1	16,24,25	1.11±0.01	0±0 (0±0%)
1	IGU	B	9	1	16,24,25	1.08±0.01	0±0 (0±0%)
1	IGU	C	14	1	16,24,25	1.10±0.01	0±0 (0±0%)
1	IGU	C	15	1	16,24,25	1.09±0.01	0±0 (0±0%)
1	IGU	C	16	1	16,24,25	1.09±0.01	0±0 (0±0%)
1	IGU	C	17	1	16,24,25	1.08±0.02	0±0 (0±0%)
1	IGU	D	20	1	16,24,25	1.11±0.01	0±0 (0±0%)
1	IGU	D	21	1	16,24,25	1.08±0.01	0±0 (0±0%)
1	IGU	D	22	1	16,24,25	1.09±0.02	0±0 (0±0%)
1	IGU	D	23	1	16,24,25	1.04±0.02	0±0 (0±0%)
1	IGU	E	26	1	16,24,25	1.11±0.01	0±0 (0±0%)
1	IGU	E	27	1	16,24,25	1.08±0.01	0±0 (0±0%)
1	IGU	E	28	1	16,24,25	1.08±0.01	0±0 (0±0%)
1	IGU	E	29	1	16,24,25	1.05±0.02	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	IGU	A	2	1	16,35,38	1.11±0.01	0±0 (0±0%)
1	IGU	A	3	1	16,35,38	1.14±0.01	0±0 (0±0%)
1	IGU	A	4	1	16,35,38	1.16±0.01	0±0 (0±0%)
1	IGU	A	5	1	16,35,38	1.40±0.02	0±0 (0±0%)
1	IGU	B	10	1	16,35,38	1.18±0.01	0±0 (0±0%)
1	IGU	B	11	1	16,35,38	1.39±0.02	0±0 (0±0%)
1	IGU	B	8	1	16,35,38	1.10±0.01	0±0 (0±0%)
1	IGU	B	9	1	16,35,38	1.13±0.01	0±0 (0±0%)
1	IGU	C	14	1	16,35,38	1.09±0.02	0±0 (0±0%)

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	IGU	C	15	1	16,35,38	1.14±0.01	0±0 (0±0%)
1	IGU	C	16	1	16,35,38	1.17±0.02	0±0 (0±0%)
1	IGU	C	17	1	16,35,38	1.47±0.08	0±0 (0±0%)
1	IGU	D	20	1	16,35,38	1.11±0.01	0±0 (0±0%)
1	IGU	D	21	1	16,35,38	1.14±0.02	0±0 (0±0%)
1	IGU	D	22	1	16,35,38	1.17±0.01	0±0 (0±0%)
1	IGU	D	23	1	16,35,38	1.39±0.02	0±0 (0±0%)
1	IGU	E	26	1	16,35,38	1.11±0.02	0±0 (0±0%)
1	IGU	E	27	1	16,35,38	1.14±0.01	0±0 (0±0%)
1	IGU	E	28	1	16,35,38	1.17±0.01	0±0 (0±0%)
1	IGU	E	29	1	16,35,38	1.40±0.01	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	IGU	A	2	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	A	3	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	A	4	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	A	5	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	B	10	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	B	11	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	B	8	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	B	9	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	C	14	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	C	15	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	C	16	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	C	17	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	D	20	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	D	21	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	D	22	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	D	23	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	E	26	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	E	27	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	E	28	1	-	0±0,3,21,22	0±0,3,3,3
1	IGU	E	29	1	-	0±0,3,21,22	0±0,3,3,3

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 69% for the well-defined parts and 69% for the entire structure.

### 7.1 Chemical shift list 1

File name: 2luj\_cs.str

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	456
Number of shifts mapped to atoms	155
Number of unparsed shifts	0
Number of shifts with mapping errors	301
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Residue not found in structure. First 5 (of 301) occurrences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
D	22	0UH	H8	8.02	-1.0	1
C	17	0UH	H8	8.09	-1.0	1
D	21	0UH	H4'	4.55	-1.0	1
C	15	0UH	H4'	4.55	-1.0	1
D	20	0UH	H8	8.21	-1.0	1

#### 7.1.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 69%, i.e. 125 atoms were assigned a chemical shift out of a possible 180. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Sidechain	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	125/180 (69%)	75/100 (75%)	50/70 (71%)	0/10 (0%)

### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

### 7.1.5 Random Coil Index (RCI) plots [i](#)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned\_chem\_shift\_list\_1). RCI is only applicable to proteins.