



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 7, 2017 – 02:36 AM EST

PDB ID : 5LWF  
Title : Structure of a single domain camelid antibody fragment cAb-G10S in complex with the BlaP beta-lactamase from *Bacillus licheniformis*  
Authors : Vettore, N.; Kerff, F.; Pain, C.; Herman, R.; Sauvage, E.; Preumont, S.; Charlier, P.; Dumoulin, M.  
Deposited on : unknown  
Resolution : 2.56 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3689 (2.60-2.52)
Clashscore	112137	4096 (2.60-2.52)
Ramachandran outliers	110173	4037 (2.60-2.52)
Sidechain outliers	110143	4037 (2.60-2.52)
RSRZ outliers	101464	3700 (2.60-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	273	<div> <div>3%</div> <div>85%</div> <div>9%</div> <div>5%</div> </div>
1	B	273	<div> <div>16%</div> <div>86%</div> <div>7%</div> <div>5%</div> </div>
2	C	131	<div> <div>5%</div> <div>75%</div> <div>17%</div> <div>7%</div> </div>
2	D	131	<div> <div>18%</div> <div>76%</div> <div>12%</div> <div>10%</div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5927 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	259	Total	C	N	O	S	0	0	0
			2019	1266	349	401	3			
1	B	258	Total	C	N	O	S	0	0	0
			2014	1263	348	400	3			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	197A	PRO	-	insertion	UNP P00808
A	197B	GLY	-	insertion	UNP P00808
A	296	GLY	-	expression tag	UNP P00808
A	297	PRO	-	expression tag	UNP P00808
A	298	HIS	-	expression tag	UNP P00808
A	299	HIS	-	expression tag	UNP P00808
A	300	HIS	-	expression tag	UNP P00808
A	301	HIS	-	expression tag	UNP P00808
A	302	HIS	-	expression tag	UNP P00808
B	197A	PRO	-	insertion	UNP P00808
B	197B	GLY	-	insertion	UNP P00808
B	296	GLY	-	expression tag	UNP P00808
B	297	PRO	-	expression tag	UNP P00808
B	298	HIS	-	expression tag	UNP P00808
B	299	HIS	-	expression tag	UNP P00808
B	300	HIS	-	expression tag	UNP P00808
B	301	HIS	-	expression tag	UNP P00808
B	302	HIS	-	expression tag	UNP P00808

- Molecule 2 is a protein called Camelid heavy-chain antibody variable fragment cAb-G10S.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	122	Total	C	N	O	S	0	0	0
			948	599	163	182	4			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	118	Total	C	N	O	S	0	0	0
			925	584	159	178	4			

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-lactamase




- Molecule 1: Beta-lactamase



- Molecule 2: Camelid heavy-chain antibody variable fragment cAb-G10S



- Molecule 2: Camelid heavy-chain antibody variable fragment cAb-G10S



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.28Å 121.69Å 137.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.50 – 2.56 45.50 – 2.56	Depositor EDS
% Data completeness (in resolution range)	93.9 (40.50-2.56) 94.4 (45.50-2.56)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.34 (at 2.58Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.215 , 0.242 0.239 , 0.274	Depositor DCC
$R_{free}$ test set	1230 reflections (4.98%)	DCC
Wilson B-factor (Å <sup>2</sup> )	34.2	Xtriage
Anisotropy	0.535	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 41.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	5927	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/2050	0.64	1/2777 (0.0%)
1	B	0.36	0/2045	0.64	0/2770
2	C	0.35	0/972	0.61	0/1317
2	D	0.37	0/948	0.60	0/1283
All	All	0.36	0/6015	0.63	1/8147 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	197	ASP	CA-CB-CG	-5.38	101.56	113.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2019	0	2028	12	0
1	B	2014	0	2026	8	0
2	C	948	0	900	9	0
2	D	925	0	873	7	0
3	A	8	0	6	0	0
3	C	8	0	6	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	4	0	3	0	0
4	C	1	0	0	0	0
All	All	5927	0	5842	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 36 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:THR:HG21	1:A:133:ALA:HB3	1.50	0.94
1:A:44:LEU:HD21	1:A:278:LEU:HD11	1.58	0.83
1:B:109:THR:HG21	1:B:133:ALA:HB3	1.70	0.73
2:D:71:PHE:HD1	2:D:72:ALA:H	1.38	0.69
1:A:106:ASN:HB3	1:A:109:THR:CG2	2.27	0.63

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/273 (94%)	250 (97%)	5 (2%)	2 (1%)	22	41
1	B	256/273 (94%)	251 (98%)	4 (2%)	1 (0%)	38	59
2	C	120/131 (92%)	115 (96%)	5 (4%)	0	100	100
2	D	114/131 (87%)	106 (93%)	7 (6%)	1 (1%)	20	37
All	All	747/808 (92%)	722 (97%)	21 (3%)	4 (0%)	32	54

All (4) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
2	D	72	ALA
1	A	197(B)	GLY
1	B	103	VAL
1	A	103	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/230 (94%)	210 (97%)	7 (3%)	44	68
1	B	217/230 (94%)	205 (94%)	12 (6%)	25	45
2	C	96/105 (91%)	88 (92%)	8 (8%)	13	23
2	D	94/105 (90%)	87 (93%)	7 (7%)	16	29
All	All	624/670 (93%)	590 (95%)	34 (5%)	26	45

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	233	ASP
2	C	14	LEU
2	D	75	ARG
1	B	268	ASP
1	A	261	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ACT	A	401	-	1,3,3	4.77	1 (100%)	0,3,3	0.00	-
3	ACT	A	402	-	1,3,3	4.84	1 (100%)	0,3,3	0.00	-
3	ACT	C	201	-	1,3,3	4.92	1 (100%)	0,3,3	0.00	-
3	ACT	C	202	-	1,3,3	4.83	1 (100%)	0,3,3	0.00	-
3	ACT	D	201	-	1,3,3	4.80	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACT	A	401	-	-	0/0/0/0	0/0/0/0
3	ACT	A	402	-	-	0/0/0/0	0/0/0/0
3	ACT	C	201	-	-	0/0/0/0	0/0/0/0
3	ACT	C	202	-	-	0/0/0/0	0/0/0/0
3	ACT	D	201	-	-	0/0/0/0	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	ACT	CH3-C	4.77	1.54	1.48
3	D	201	ACT	CH3-C	4.80	1.54	1.48
3	C	202	ACT	CH3-C	4.83	1.55	1.48
3	A	402	ACT	CH3-C	4.84	1.55	1.48
3	C	201	ACT	CH3-C	4.92	1.55	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	259/273 (94%)	0.36	8 (3%) 49 46	21, 44, 78, 106	0
1	B	258/273 (94%)	0.91	44 (17%) 2 1	19, 61, 124, 152	0
2	C	122/131 (93%)	0.31	6 (4%) 30 27	16, 44, 70, 81	0
2	D	118/131 (90%)	1.26	24 (20%) 1 1	22, 72, 106, 116	0
All	All	757/808 (93%)	0.68	82 (10%) 6 5	16, 50, 106, 152	0

The worst 5 of 82 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	256	GLY	11.2
2	D	78	ALA	7.6
1	B	291	ASN	6.1
1	B	229	TRP	5.9
2	D	79	LYS	5.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ACT	C	201	4/4	0.92	0.24	1.83	33,34,35,35	0
3	ACT	A	401	4/4	0.89	0.24	1.34	55,56,56,56	0
3	ACT	A	402	4/4	0.94	0.15	-0.42	33,33,34,34	0
3	ACT	D	201	4/4	0.97	0.16	-0.78	40,40,40,41	0
3	ACT	C	202	4/4	0.69	0.41	-	49,50,50,52	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.