



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 08:34 am GMT

PDB ID : 3LX4
Title : Stepwise [FeFe]-hydrogenase H-cluster assembly revealed in the structure of HydA(deltaEFG)
Authors : Mulder, D.W.; Boyd, E.S.; Sarma, R.; Lange, R.K.; Endrizzi, J.A.; Broderick, J.B.; Peters, J.W.
Deposited on : 2010-02-24
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	9293 (2.00-1.96)
Clashscore	112137	10621 (2.00-1.96)
Ramachandran outliers	110173	10502 (2.00-1.96)
Sidechain outliers	110143	10501 (2.00-1.96)
RSRZ outliers	101464	9395 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	457	 2% 81% 8% 11%
1	B	457	 2% 81% 8% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACT	A	460	-	-	X	X
4	ACT	B	460	-	-	X	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7005 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fe-hydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	407	3102	1954	538	585	25	7	3	0
1	B	407	3102	1954	538	585	25	7	3	0

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP Q9FYU1
A	2	GLY	-	EXPRESSION TAG	UNP Q9FYU1
A	3	SER	-	EXPRESSION TAG	UNP Q9FYU1
A	4	SER	-	EXPRESSION TAG	UNP Q9FYU1
A	5	HIS	-	EXPRESSION TAG	UNP Q9FYU1
A	6	HIS	-	EXPRESSION TAG	UNP Q9FYU1
A	7	HIS	-	EXPRESSION TAG	UNP Q9FYU1
A	8	HIS	-	EXPRESSION TAG	UNP Q9FYU1
A	9	HIS	-	EXPRESSION TAG	UNP Q9FYU1
A	10	HIS	-	EXPRESSION TAG	UNP Q9FYU1
A	11	SER	-	EXPRESSION TAG	UNP Q9FYU1
A	12	GLN	-	EXPRESSION TAG	UNP Q9FYU1
A	13	ASP	-	EXPRESSION TAG	UNP Q9FYU1
A	14	PRO	-	EXPRESSION TAG	UNP Q9FYU1
A	15	ASN	-	EXPRESSION TAG	UNP Q9FYU1
A	16	SER	-	EXPRESSION TAG	UNP Q9FYU1
B	1	MET	-	EXPRESSION TAG	UNP Q9FYU1
B	2	GLY	-	EXPRESSION TAG	UNP Q9FYU1
B	3	SER	-	EXPRESSION TAG	UNP Q9FYU1
B	4	SER	-	EXPRESSION TAG	UNP Q9FYU1
B	5	HIS	-	EXPRESSION TAG	UNP Q9FYU1
B	6	HIS	-	EXPRESSION TAG	UNP Q9FYU1
B	7	HIS	-	EXPRESSION TAG	UNP Q9FYU1
B	8	HIS	-	EXPRESSION TAG	UNP Q9FYU1
B	9	HIS	-	EXPRESSION TAG	UNP Q9FYU1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	10	HIS	-	EXPRESSION TAG	UNP Q9FYU1
B	11	SER	-	EXPRESSION TAG	UNP Q9FYU1
B	12	GLN	-	EXPRESSION TAG	UNP Q9FYU1
B	13	ASP	-	EXPRESSION TAG	UNP Q9FYU1
B	14	PRO	-	EXPRESSION TAG	UNP Q9FYU1
B	15	ASN	-	EXPRESSION TAG	UNP Q9FYU1
B	16	SER	-	EXPRESSION TAG	UNP Q9FYU1

- Molecule 2 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Fe S 8 4 4	0	0
2	B	1	Total Fe S 8 4 4	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Cl 1 1	0	0
3	A	1	Total Cl 1 1	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	364	Total O 364 364	0	0
5	B	411	Total O 411 411	0	0

4 Data and refinement statistics i

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	70.89Å 70.89Å 155.43Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 1.97 19.92 – 1.97	Depositor EDS
% Data completeness (in resolution range)	97.8 (20.00-1.97) 97.8 (19.92-1.97)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.87 (at 1.96Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.170 , 0.216 0.171 , 0.219	Depositor DCC
R_{free} test set	3060 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	22.8	Xtrriage
Anisotropy	0.088	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.006 for -h,-k,l 0.479 for h,-h-k,-l 0.008 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7005	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, CL, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/3173	0.47	0/4289
1	B	0.32	0/3173	0.48	1/4289 (0.0%)
All	All	0.32	0/6346	0.48	1/8578 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	137	LEU	CA-CB-CG	5.34	127.59	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3102	0	3114	31	0
1	B	3102	0	3114	27	0
2	A	8	0	0	1	0
2	B	8	0	0	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	4	0	3	4	0
4	B	4	0	3	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	364	0	0	5	1
5	B	411	0	0	3	1
All	All	7005	0	6234	60	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:458:SF4:S3	4:B:460:ACT:H1	2.26	0.74
1:B:169:LYS:HE2	1:B:242:VAL:HG13	1.70	0.73
1:A:114:GLU:HA	1:A:309:ARG:HD2	1.73	0.71
1:A:406:LEU:HD22	1:A:406:LEU:H	1.56	0.71
1:B:138:GLU:OE1	1:B:194:ARG:NH2	2.24	0.70
1:B:416:HIS:HD2	5:B:527:HOH:O	1.75	0.69
1:B:138:GLU:CD	1:B:194:ARG:HH22	1.98	0.66
1:B:114:GLU:HA	1:B:309:ARG:HD2	1.79	0.64
1:A:416:HIS:HD2	5:A:590:HOH:O	1.80	0.63
1:A:249:LEU:O	1:A:413:ARG:HD3	2.00	0.62
1:B:129[C]:CYS:SG	4:B:460:ACT:C	2.86	0.62
1:A:129[C]:CYS:SG	4:A:460:ACT:C	2.87	0.62
1:A:138:GLU:OE1	1:A:194:ARG:NH2	2.30	0.61
1:A:432:PRO:O	1:A:433:LEU:HB2	2.02	0.58
1:B:249:LEU:O	1:B:413:ARG:HD3	2.03	0.58
1:A:169:LYS:HE2	1:A:242:VAL:HG13	1.85	0.57
1:A:138:GLU:CD	1:A:194:ARG:HH22	2.08	0.57
1:B:339:ARG:NH1	5:B:563:HOH:O	2.23	0.56
1:B:406:LEU:HD22	1:B:406:LEU:H	1.70	0.56
1:B:265:TYR:HA	1:B:294:MET:CE	2.37	0.55
1:A:331:ALA:HB3	1:A:343:THR:HG23	1.89	0.53
1:B:335:SER:OG	1:B:337:ASP:OD1	2.25	0.53
1:B:265:TYR:HA	1:B:294:MET:HE3	1.90	0.53
1:A:351:GLY:HA2	1:A:375:MET:O	2.09	0.52
1:A:284:MET:SD	1:A:361:LYS:HD2	2.49	0.52
1:B:129[C]:CYS:HG	4:B:460:ACT:C	2.23	0.52
2:A:458:SF4:S2	4:A:460:ACT:H2	2.50	0.52
1:A:129[C]:CYS:HG	4:A:460:ACT:C	2.24	0.51
1:A:339:ARG:NH1	5:A:663:HOH:O	2.24	0.50
1:B:53:PRO:HG3	1:B:56[B]:ARG:NH2	2.28	0.49
1:A:397:GLN:NE2	5:A:489:HOH:O	2.45	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:LEU:HD22	1:A:148:VAL:HG21	1.94	0.48
1:B:284:MET:SD	1:B:361:LYS:HD2	2.53	0.48
1:A:375:MET:SD	4:A:460:ACT:H1	2.55	0.47
1:B:166:LEU:HD11	1:B:170:LYS:HE3	1.97	0.47
1:B:179:MET:HG2	1:B:208:LEU:HD13	1.96	0.47
1:B:121:GLU:N	5:B:1100:HOH:O	2.48	0.46
1:B:129[A]:CYS:SG	4:B:460:ACT:H2	2.55	0.45
1:A:409:LYS:HE2	5:A:598:HOH:O	2.15	0.45
1:A:278:LEU:HB3	1:A:288:LYS:HB3	1.99	0.45
1:A:137:LEU:CD2	1:A:148:VAL:HG21	2.47	0.45
1:A:129[C]:CYS:SG	1:A:154:PRO:HD2	2.58	0.43
1:A:275:ARG:NH2	5:A:927:HOH:O	2.40	0.43
1:B:107:HIS:O	1:B:111:GLU:HG2	2.18	0.43
1:B:53:PRO:HG3	1:B:56[B]:ARG:HH22	1.84	0.43
1:A:265:TYR:HA	1:A:294:MET:CE	2.49	0.42
1:A:406:LEU:HD22	1:A:406:LEU:N	2.31	0.42
1:A:265:TYR:HA	1:A:294:MET:HE1	2.01	0.42
1:B:96:LEU:HD21	1:B:424:LEU:HD23	2.01	0.42
1:B:327:TRP:C	1:B:329:GLY:H	2.23	0.42
1:B:56[A]:ARG:HD3	1:B:72:PRO:HG3	2.01	0.42
1:A:56[A]:ARG:HD3	1:A:72:PRO:HG3	2.01	0.42
1:B:179:MET:HG2	1:B:208:LEU:CD1	2.49	0.42
1:A:166:LEU:HD11	1:A:170:LYS:HE3	2.02	0.41
1:A:107:HIS:O	1:A:111:GLU:HG2	2.21	0.41
1:A:406:LEU:H	1:A:406:LEU:CD2	2.29	0.41
1:B:129[C]:CYS:SG	1:B:154:PRO:HD2	2.60	0.41
1:B:352:LEU:O	1:B:356:LYS:HG3	2.21	0.41
1:A:350:ASN:HA	1:A:351:GLY:HA2	1.96	0.40
1:A:258:GLU:OE2	1:A:275:ARG:HD2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:853:HOH:O	5:B:939:HOH:O[3_665]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/457 (88%)	396 (98%)	7 (2%)	0	100	100
1	B	403/457 (88%)	398 (99%)	4 (1%)	1 (0%)	51	46
All	All	806/914 (88%)	794 (98%)	11 (1%)	1 (0%)	55	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	328	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	330/363 (91%)	330 (100%)	0	100	100
1	B	330/363 (91%)	327 (99%)	3 (1%)	82	85
All	All	660/726 (91%)	657 (100%)	3 (0%)	91	93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	137	LEU
1	B	177	MET
1	B	217	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	155	GLN
1	A	416	HIS
1	B	155	GLN
1	B	416	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SF4	A	458	1	0,12,12	0.00	-	0,24,24	0.00	-
4	ACT	A	460	-	1,3,3	0.89	0	0,3,3	0.00	-
2	SF4	B	458	1	0,12,12	0.00	-	0,24,24	0.00	-
4	ACT	B	460	-	1,3,3	0.95	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SF4	A	458	1	-	0/0/48/48	0/6/5/5
4	ACT	A	460	-	-	0/0/0/0	0/0/0/0
2	SF4	B	458	1	-	0/0/48/48	0/6/5/5
4	ACT	B	460	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	458	SF4	1	0
4	A	460	ACT	4	0
2	B	458	SF4	1	0
4	B	460	ACT	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/457 (89%)	-0.36	9 (2%) 62 65	12, 22, 47, 88	1 (0%)
1	B	407/457 (89%)	-0.37	7 (1%) 70 72	12, 22, 46, 88	1 (0%)
All	All	814/914 (89%)	-0.36	16 (1%) 65 68	12, 22, 47, 88	2 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	311	ALA	4.6
1	A	406	LEU	4.2
1	A	311	ALA	3.9
1	A	323	GLY	3.7
1	A	308	HIS	3.6
1	B	327	TRP	3.6
1	A	326	ALA	3.1
1	B	326	ALA	3.0
1	B	406	LEU	2.9
1	A	327	TRP	2.8
1	A	310	ALA	2.6
1	B	323	GLY	2.5
1	A	204	THR	2.3
1	A	309	ARG	2.2
1	B	309	ARG	2.2
1	B	307	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	ACT	A	460	4/4	0.74	0.15	3.37	39,45,45,49	0
4	ACT	B	460	4/4	0.75	0.17	2.88	33,46,51,51	0
2	SF4	B	458	8/8	0.99	0.06	-1.30	19,20,21,24	0
3	CL	B	459	1/1	1.00	0.04	-1.37	16,16,16,16	0
2	SF4	A	458	8/8	0.99	0.05	-1.50	18,20,21,24	0
3	CL	A	459	1/1	1.00	0.04	-2.86	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.