



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 30, 2017 – 07:48 AM EDT

PDB ID : 3LX6
Title : Crystal structure of putative dna cytosine methylase from shigella flexneri 2a str. 2457T
Authors : Ramagopal, U.A.; Toro, R.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : unknown
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	410	<div> <div>10%</div> <div> <div></div> <div>78%</div> <div>11%</div> <div>•</div> <div>10%</div> </div> </div>
1	B	410	<div> <div>19%</div> <div> <div></div> <div>71%</div> <div>16%</div> <div>•</div> <div>12%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5835 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytosine-specific methyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	368	Total	C	N	O	S	Se	0	0	0
			2964	1889	531	532	6	6			
1	B	362	Total	C	N	O	S	Se	0	0	0
			2842	1800	514	516	6	6			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	MSE	-	EXPRESSION TAG	UNP E3Y0J2
A	62	SER	-	EXPRESSION TAG	UNP E3Y0J2
A	463	GLU	-	EXPRESSION TAG	UNP E3Y0J2
A	464	GLY	-	EXPRESSION TAG	UNP E3Y0J2
A	465	HIS	-	EXPRESSION TAG	UNP E3Y0J2
A	466	HIS	-	EXPRESSION TAG	UNP E3Y0J2
A	467	HIS	-	EXPRESSION TAG	UNP E3Y0J2
A	468	HIS	-	EXPRESSION TAG	UNP E3Y0J2
A	469	HIS	-	EXPRESSION TAG	UNP E3Y0J2
A	470	HIS	-	EXPRESSION TAG	UNP E3Y0J2
B	61	MSE	-	EXPRESSION TAG	UNP E3Y0J2
B	62	SER	-	EXPRESSION TAG	UNP E3Y0J2
B	463	GLU	-	EXPRESSION TAG	UNP E3Y0J2
B	464	GLY	-	EXPRESSION TAG	UNP E3Y0J2
B	465	HIS	-	EXPRESSION TAG	UNP E3Y0J2
B	466	HIS	-	EXPRESSION TAG	UNP E3Y0J2
B	467	HIS	-	EXPRESSION TAG	UNP E3Y0J2
B	468	HIS	-	EXPRESSION TAG	UNP E3Y0J2
B	469	HIS	-	EXPRESSION TAG	UNP E3Y0J2
B	470	HIS	-	EXPRESSION TAG	UNP E3Y0J2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	24	Total 24	O 24	0	0
2	B	5	Total 5	O 5	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cytosine-specific methyltransferase



• Molecule 1: Cytosine-specific methyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	122.96Å 83.81Å 113.24Å 90.00° 118.51° 90.00°	Depositor
Resolution (Å)	31.26 – 2.29 31.26 – 2.29	Depositor EDS
% Data completeness (in resolution range)	96.5 (31.26-2.29) 96.5 (31.26-2.29)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.24 (at 2.29Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.243 , 0.288 0.246 , 0.291	Depositor DCC
R_{free} test set	2214 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	43.5	Xtriage
Anisotropy	0.355	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 28.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	5835	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	1/3034 (0.0%)	0.77	1/4100 (0.0%)
1	B	0.62	0/2903	0.68	1/3919 (0.0%)
All	All	0.70	1/5937 (0.0%)	0.73	2/8019 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	110	CYS	CB-SG	-5.04	1.73	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	91	ASP	CB-CG-OD1	-5.14	113.67	118.30
1	B	211	ARG	NE-CZ-NH1	5.01	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2964	0	2911	26	0
1	B	2842	0	2733	50	0
2	A	24	0	0	0	0
2	B	5	0	0	0	0
All	All	5835	0	5644	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:178:GLN:HE22	1:B:205:LEU:H	0.96	0.91
1:A:205:LEU:HD21	1:B:306:GLN:NE2	1.91	0.86
1:B:415:ALA:HB3	1:B:418:GLU:HG3	1.59	0.84
1:B:90:ILE:HD11	1:B:171:LEU:HD12	1.65	0.77
1:B:274:GLN:HB2	1:B:410:LEU:HD11	1.68	0.75
1:B:180:PHE:CD1	1:B:201:THR:HA	2.22	0.74
1:B:178:GLN:HE22	1:B:205:LEU:N	1.80	0.74
1:A:310:LEU:CD2	1:A:351:VAL:HG11	2.22	0.70
1:B:178:GLN:NE2	1:B:205:LEU:H	1.82	0.67
1:A:243:GLN:O	1:A:247:GLU:HG3	1.99	0.62
1:A:310:LEU:HD22	1:A:351:VAL:HG11	1.79	0.62
1:A:82:HIS:HD2	1:A:85:TYR:OH	1.82	0.61
1:B:300:SER:O	1:B:303:PHE:HB2	2.01	0.61
1:A:428:THR:HG21	1:B:298:ASP:OD2	2.00	0.61
1:B:371:GLY:HA3	1:B:401:ARG:HD3	1.83	0.60
1:B:310:LEU:CD2	1:B:351:VAL:HG11	2.32	0.60
1:B:276:ARG:HD3	1:B:439:VAL:CG2	2.32	0.60
1:A:220:MSE:HE1	1:A:454:ILE:HA	1.85	0.58
1:B:146:LEU:CD1	1:B:152:VAL:HG11	2.34	0.58
1:B:445:ALA:HA	1:B:448:LYS:HE2	1.86	0.57
1:B:263:PRO:HB3	1:B:297:ARG:HG3	1.87	0.57
1:B:311:ALA:HB2	1:B:353:PRO:O	2.05	0.56
1:B:220:MSE:HE1	1:B:454:ILE:HA	1.87	0.56
1:B:90:ILE:HD11	1:B:171:LEU:CD1	2.35	0.55
1:A:408:ALA:HB2	1:A:434:PHE:CZ	2.42	0.54
1:B:376:ILE:HB	1:B:400:ARG:HG2	1.90	0.54
1:B:99:ILE:HG22	1:B:172:LEU:HD13	1.90	0.54
1:A:276:ARG:HD3	1:A:439:VAL:HG23	1.91	0.53
1:B:363:LEU:HD11	1:B:407:CYS:SG	2.48	0.53
1:B:276:ARG:HD3	1:B:439:VAL:HG23	1.91	0.53
1:B:276:ARG:NE	1:B:411:MSE:HE1	2.25	0.52
1:A:93:PHE:CD1	1:A:116:TRP:HZ3	2.29	0.51
1:A:93:PHE:CZ	1:A:205:LEU:HD22	2.46	0.50
1:A:136:HIS:NE2	1:A:164:HIS:O	2.38	0.50
1:B:146:LEU:HD13	1:B:152:VAL:HG11	1.94	0.49
1:A:161:ILE:HD11	1:A:211:ARG:HG2	1.94	0.49
1:B:87:PHE:HE2	1:B:106:ILE:HG13	1.77	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:367:TYR:OH	1:B:401:ARG:HG2	2.13	0.48
1:A:93:PHE:HD1	1:A:116:TRP:HZ3	1.61	0.48
1:B:119:HIS:N	1:B:119:HIS:CD2	2.82	0.48
1:A:93:PHE:CE1	1:A:205:LEU:HD22	2.49	0.48
1:B:180:PHE:HD1	1:B:201:THR:HA	1.74	0.48
1:B:145:THR:O	1:B:211:ARG:NH2	2.39	0.47
1:B:90:ILE:HD13	1:B:168:HIS:CE1	2.50	0.47
1:A:364:SER:CB	1:A:366:ARG:HE	2.28	0.46
1:B:89:PHE:CZ	1:B:110:CYS:HB2	2.50	0.46
1:B:92:LEU:HB2	1:B:173:ALA:HB2	1.98	0.46
1:B:310:LEU:HD22	1:B:351:VAL:HG11	1.98	0.46
1:A:175:PHE:HE1	1:A:223:LEU:HB3	1.83	0.44
1:B:173:ALA:O	1:B:223:LEU:HA	2.17	0.44
1:B:425:VAL:HB	1:B:429:GLN:HB2	1.99	0.44
1:B:432:ARG:O	1:B:436:ASN:HB2	2.17	0.44
1:A:142:ARG:HD3	1:B:306:GLN:OE1	2.18	0.44
1:A:85:TYR:CG	1:A:109:GLN:HB2	2.54	0.43
1:B:146:LEU:HD12	1:B:152:VAL:HG11	1.99	0.43
1:B:170:VAL:HA	1:B:220:MSE:O	2.19	0.43
1:B:367:TYR:CD1	1:B:371:GLY:HA2	2.54	0.43
1:B:113:THR:HG21	1:B:124:TYR:CZ	2.54	0.42
1:B:274:GLN:HB2	1:B:410:LEU:CD1	2.44	0.42
1:B:158:ALA:HA	1:B:161:ILE:HD12	2.02	0.42
1:A:118:LYS:HE3	1:A:139:GLU:OE2	2.20	0.41
1:A:276:ARG:HD3	1:A:439:VAL:CG2	2.49	0.41
1:A:294:PHE:C	1:A:294:PHE:CD2	2.93	0.41
1:B:92:LEU:HB2	1:B:173:ALA:CB	2.50	0.41
1:A:125:LYS:HE2	1:A:137:PHE:CE2	2.56	0.41
1:B:310:LEU:HD23	1:B:351:VAL:HG11	2.02	0.41
1:A:269:LYS:HA	1:A:272:LEU:O	2.21	0.41
1:B:274:GLN:HB3	1:B:274:GLN:HE21	1.67	0.40
1:B:364:SER:HB2	1:B:366:ARG:HH21	1.86	0.40
1:B:99:ILE:HG13	1:B:438:VAL:HG21	2.03	0.40
1:A:288:LEU:HB3	1:A:290:LEU:HD12	2.03	0.40
1:B:146:LEU:HD12	1:B:157:ALA:HA	2.03	0.40
1:A:401:ARG:HH21	1:A:401:ARG:HD3	1.75	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	362/410 (88%)	353 (98%)	8 (2%)	1 (0%)	44	55
1	B	352/410 (86%)	334 (95%)	16 (4%)	2 (1%)	28	34
All	All	714/820 (87%)	687 (96%)	24 (3%)	3 (0%)	38	47

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	300	SER
1	B	370	ASP
1	A	316	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/340 (92%)	299 (96%)	14 (4%)	32	44
1	B	290/340 (85%)	278 (96%)	12 (4%)	35	48
All	All	603/680 (89%)	577 (96%)	26 (4%)	33	45

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	LYS
1	A	148	HIS
1	A	149	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	167	GLU
1	A	199	CYS
1	A	211	ARG
1	A	217	ARG
1	A	225	ASN
1	A	228	ASN
1	A	289	ASN
1	A	290	LEU
1	A	366	ARG
1	A	373	GLU
1	A	448	LYS
1	B	116	TRP
1	B	148	HIS
1	B	178	GLN
1	B	211	ARG
1	B	224	GLU
1	B	278	ARG
1	B	334	ARG
1	B	366	ARG
1	B	390	ASP
1	B	422	ARG
1	B	436	ASN
1	B	453	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	82	HIS
1	A	228	ASN
1	A	258	ASN
1	A	354	ASN
1	A	436	ASN
1	B	82	HIS
1	B	178	GLN
1	B	228	ASN
1	B	258	ASN
1	B	357	GLN
1	B	433	GLN
1	B	456	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	362/410 (88%)	0.74	41 (11%) 6 8	19, 30, 43, 77	0
1	B	356/410 (86%)	1.13	77 (21%) 1 1	19, 33, 48, 80	0
All	All	718/820 (87%)	0.93	118 (16%) 2 3	19, 32, 46, 80	0

All (118) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	199	CYS	7.1
1	B	331	TYR	7.0
1	B	335	TYR	6.6
1	B	182	LEU	5.7
1	B	316	PRO	5.5
1	B	332	LEU	5.3
1	B	319	GLU	5.3
1	B	383	ALA	4.8
1	B	394	ASN	4.6
1	B	421	PHE	4.6
1	A	171	LEU	4.6
1	B	390	ASP	4.5
1	B	83	PRO	4.5
1	B	389	PHE	4.4
1	A	92	LEU	4.3
1	A	222	VAL	4.3
1	B	130	CYS	4.2
1	B	336	ALA	4.2
1	A	172	LEU	4.1
1	A	170	VAL	4.1
1	B	240	ILE	4.0
1	B	393	LEU	3.9
1	A	112	PHE	3.9
1	A	173	ALA	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	165	ILE	3.9
1	B	385	GLY	3.8
1	A	221	PHE	3.7
1	A	90	ILE	3.7
1	B	150	GLU	3.6
1	B	203	GLY	3.5
1	A	213	ILE	3.4
1	A	209	VAL	3.4
1	B	397	HIS	3.4
1	A	166	PRO	3.4
1	B	82	HIS	3.4
1	B	395	GLN	3.3
1	B	388	ASP	3.2
1	A	111	VAL	3.1
1	B	84	HIS	3.1
1	B	207	PHE	3.1
1	A	110	CYS	3.1
1	A	387	LYS	3.1
1	B	415	ALA	3.1
1	B	289	ASN	3.0
1	A	91	ASP	3.0
1	B	202	GLN	3.0
1	B	163	GLN	3.0
1	B	151	GLY	3.0
1	A	74	LEU	3.0
1	B	126	ALA	3.0
1	A	174	GLY	3.0
1	A	223	LEU	3.0
1	A	168	HIS	2.9
1	B	77	LYS	2.9
1	B	363	LEU	2.8
1	B	119	HIS	2.8
1	B	322	TYR	2.8
1	B	221	PHE	2.8
1	B	314	LEU	2.7
1	B	376	ILE	2.7
1	B	386	GLU	2.7
1	B	183	ALA	2.7
1	A	338	LYS	2.7
1	B	287	ASP	2.6
1	A	120	ALA	2.6
1	B	387	LYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	96	ILE	2.6
1	A	89	PHE	2.6
1	B	114	SER	2.6
1	B	338	LYS	2.6
1	B	81	HIS	2.6
1	A	114	SER	2.6
1	B	384	THR	2.6
1	B	122	ARG	2.5
1	B	261	ASP	2.5
1	A	430	ALA	2.5
1	B	225	ASN	2.5
1	B	92	LEU	2.5
1	B	236	LYS	2.5
1	A	282	VAL	2.5
1	B	365	ALA	2.5
1	B	352	TYR	2.5
1	B	201	THR	2.5
1	B	160	HIS	2.5
1	B	396	GLN	2.5
1	B	320	ALA	2.4
1	A	99	ILE	2.4
1	A	103	PHE	2.4
1	B	337	LYS	2.4
1	A	435	GLY	2.4
1	B	205	LEU	2.4
1	B	78	PRO	2.4
1	A	402	LEU	2.3
1	B	256	GLU	2.3
1	B	372	ALA	2.3
1	A	210	VAL	2.3
1	B	85	TYR	2.3
1	B	318	VAL	2.3
1	A	97	GLY	2.3
1	B	229	LEU	2.3
1	B	136	HIS	2.3
1	B	334	ARG	2.3
1	A	443	PHE	2.2
1	A	124	TYR	2.2
1	B	326	PRO	2.2
1	B	233	ASP	2.1
1	B	377	ASP	2.1
1	A	212	ILE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	235	GLY	2.1
1	B	109	GLN	2.1
1	B	175	PHE	2.1
1	A	363	LEU	2.1
1	B	366	ARG	2.1
1	A	438	VAL	2.1
1	B	93	PHE	2.1
1	A	167	GLU	2.0
1	A	95	GLY	2.0
1	B	211	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.