



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 5, 2018 – 11:05 AM EST

PDB ID : 5LZH
Title : Cholera toxin classical B-pentamer in complex with inhibitor PC262
Authors : Heggelund, J.E.; Martinsen, T.; Krenzel, U.
Deposited on : 2016-09-29
Resolution : 1.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1062 (1.16-1.08)
Clashscore	112137	1107 (1.16-1.08)
Ramachandran outliers	110173	1065 (1.16-1.08)
Sidechain outliers	110143	1062 (1.16-1.08)
RSRZ outliers	101464	1067 (1.16-1.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	103	<div> <div>90%</div> <div>10%</div> </div>
1	B	103	<div> <div>83%</div> <div>16%</div> <div>.</div> </div>
1	D	103	<div> <div>%</div> <div>80%</div> <div>18%</div> <div>.</div> </div>
2	C	103	<div> <div>%</div> <div>92%</div> <div>8%</div> </div>
2	E	103	<div> <div>2%</div> <div>87%</div> <div>12%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	CSX	A	9	-	-	X	-
3	7BN	D	201	-	-	-	X
4	MRD	B	202	-	-	-	X
6	TRS	D	202	-	-	-	X
7	PEG	E	202	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 5093 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cholera enterotoxin B subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	103	Total	C	N	O	S	0	7	0
			871	546	153	164	8			
1	B	103	Total	C	N	O	S	0	4	0
			841	528	145	162	6			
1	D	103	Total	C	N	O	S	0	6	0
			852	538	147	160	7			

- Molecule 2 is a protein called Cholera enterotoxin B subunit.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	103	Total	C	N	O	S	0	3	0
			835	524	146	160	5			
2	E	103	Total	C	N	O	S	0	5	0
			845	532	147	160	6			

- Molecule 3 is (2 {R},4 {S},5 {R},6 {R})-5-acetamido-2-[4-[3-[2-[(2 {S},3 {R},4 {R},5 {R},6 {R})-6-(hydroxymethyl)-3,4,5-tris(oxidanyl)oxan-2-yl]ethylamino]-3-oxidanylidene-propyl]-1,2,3-triazol-1-yl]-4-oxidanyl-6-[(1 {R},2 {R})-1,2,3-tris(oxidanyl)propyl]oxane-2-carboxylic acid (three-letter code: 7BN) (formula: C₂₄H₃₉N₅O₁₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			43	24	5	14		
3	B	1	Total	C	N	O	0	0
			43	24	5	14		
3	C	1	Total	C	N	O	0	0
			43	24	5	14		
3	D	1	Total	C	N	O	0	0
			43	24	5	14		
3	E	1	Total	C	N	O	0	0
			43	24	5	14		

- Molecule 4 is (4R)-2-METHYLPENTANE-2,4-DIOL (three-letter code: MRD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	2	Total	Ca	0	0
			2	2		
5	E	1	Total	Ca	0	0
			1	1		

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	D	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	E	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	127	Total 127	O 127	0	0
8	B	112	Total 112	O 112	0	0
8	C	125	Total 125	O 125	0	0
8	D	119	Total 119	O 119	0	0
8	E	125	Total 125	O 125	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cholera enterotoxin B subunit

Chain A: 





- Molecule 1: Cholera enterotoxin B subunit

Chain B: 




- Molecule 1: Cholera enterotoxin B subunit

Chain D: 




- Molecule 2: Cholera enterotoxin B subunit

Chain C: 



- Molecule 2: Cholera enterotoxin B subunit

Chain E: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	101.74 Å 66.06 Å 76.39 Å 90.00° 105.67° 90.00°	Depositor
Resolution (Å)	73.55 – 1.13 48.98 – 1.13	Depositor EDS
% Data completeness (in resolution range)	73.8 (73.55-1.13) 73.8 (48.98-1.13)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 1.12 Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.197 , 0.229 0.207 , 0.234	Depositor DCC
R_{free} test set	6819 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	10.8	Xtriage
Anisotropy	1.007	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5093	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MRD, CSX, 7BN, TRS, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.30	1/876 (0.1%)	1.11	1/1176 (0.1%)
1	B	1.40	5/851 (0.6%)	1.17	3/1144 (0.3%)
1	D	1.38	5/864 (0.6%)	1.23	7/1160 (0.6%)
2	C	1.34	3/855 (0.4%)	1.19	4/1154 (0.3%)
2	E	1.39	4/869 (0.5%)	1.25	7/1173 (0.6%)
All	All	1.36	18/4315 (0.4%)	1.19	22/5807 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	D	0	1
All	All	0	2

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	83	GLU	CD-OE2	8.11	1.34	1.25
1	D	55	SER	CB-OG	-7.95	1.31	1.42
1	B	27	TYR	CE1-CZ	-7.35	1.28	1.38
1	D	29	GLU	CB-CG	-7.16	1.38	1.52
1	A	83	GLU	CD-OE1	6.67	1.32	1.25
2	C	30	SER	CA-CB	6.14	1.62	1.52
1	D	11	GLU	CD-OE1	5.93	1.32	1.25
1	D	27	TYR	CE1-CZ	-5.89	1.30	1.38
2	C	29	GLU	CD-OE2	-5.84	1.19	1.25
1	D	79	GLU	CD-OE2	-5.79	1.19	1.25
1	B	29[A]	GLU	CD-OE2	5.73	1.31	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	29[B]	GLU	CD-OE2	5.73	1.31	1.25
2	E	29	GLU	CB-CG	-5.44	1.41	1.52
2	E	100	SER	CA-CB	5.32	1.60	1.52
1	B	30	SER	CA-CB	5.22	1.60	1.52
2	E	26	SER	CB-OG	-5.09	1.35	1.42
1	B	70	ASP	CB-CG	5.09	1.62	1.51
2	C	60	SER	CB-OG	5.08	1.48	1.42

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	70	ASP	CB-CG-OD2	-8.85	110.33	118.30
1	B	73	ARG	NE-CZ-NH1	7.83	124.22	120.30
1	B	35	ARG	NE-CZ-NH1	7.69	124.14	120.30
2	E	67	ARG	NE-CZ-NH1	7.29	123.95	120.30
1	D	22	ASP	CB-CG-OD1	7.07	124.67	118.30
1	A	22	ASP	CB-CG-OD1	7.00	124.60	118.30
2	E	22	ASP	CB-CG-OD1	6.32	123.99	118.30
2	C	73	ARG	NE-CZ-NH1	6.29	123.44	120.30
2	C	29	GLU	OE1-CD-OE2	6.28	130.83	123.30
1	D	22	ASP	CB-CG-OD2	-6.15	112.77	118.30
2	E	81[A]	LYS	CD-CE-NZ	5.96	125.41	111.70
2	E	81[B]	LYS	CD-CE-NZ	5.96	125.41	111.70
2	E	35	ARG	NE-CZ-NH1	5.95	123.28	120.30
1	D	29	GLU	OE1-CD-OE2	5.79	130.25	123.30
1	D	35	ARG	NE-CZ-NH1	5.60	123.10	120.30
2	C	29	GLU	N-CA-CB	-5.59	100.54	110.60
2	C	35	ARG	NE-CZ-NH1	5.45	123.02	120.30
1	B	59	ASP	CB-CG-OD2	-5.44	113.40	118.30
2	E	70	ASP	CB-CG-OD2	-5.44	113.41	118.30
1	D	12	TYR	CB-CG-CD2	5.41	124.25	121.00
2	E	27	TYR	CG-CD1-CE1	5.23	125.48	121.30
1	D	103	ASN	CB-CA-C	5.04	120.49	110.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	49	GLN	Peptide
1	D	29	GLU	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	871	0	879	14	0
1	B	841	0	852	9	0
1	D	852	0	880	14	0
2	C	835	0	842	4	0
2	E	845	0	857	4	0
3	A	43	0	0	0	0
3	B	43	0	0	0	0
3	C	43	0	0	0	0
3	D	43	0	0	1	0
3	E	43	0	0	0	0
4	B	8	0	14	0	0
5	C	2	0	0	0	0
5	E	1	0	0	0	1
6	D	8	0	12	0	0
7	E	7	0	10	0	0
8	A	127	0	0	0	0
8	B	112	0	0	2	0
8	C	125	0	0	3	0
8	D	119	0	0	0	1
8	E	125	0	0	1	0
All	All	5093	0	4346	40	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:C:350:HOH:O	1:D:92[A]:THR:HG22	1.70	0.92
1:A:9:CSX:OD	1:A:15[A]:THR:HB	1.74	0.88
1:D:67[B]:ARG:HG3	1:D:67[B]:ARG:HH21	1.47	0.80
1:A:9:CSX:OD	1:A:15[B]:THR:CB	2.30	0.80
1:A:9:CSX:OD	1:A:15[B]:THR:HB	1.84	0.77
1:D:67[B]:ARG:NH2	1:D:67[B]:ARG:HG3	1.99	0.76
1:D:37[B]:MET:CE	1:D:39:ILE:HD11	2.17	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67[B]:ARG:HG3	1:A:67[B]:ARG:NH1	2.04	0.73
1:B:9:CSX:OD	1:B:15:THR:HB	1.93	0.69
2:C:6[A]:THR:HG21	8:C:353:HOH:O	1.95	0.66
2:C:67[B]:ARG:NH2	1:D:70:ASP:OD1	2.30	0.64
1:A:67[B]:ARG:HH11	1:A:67[B]:ARG:HG3	1.64	0.60
1:A:9:CSX:OD	1:A:15[B]:THR:OG1	2.22	0.57
1:D:67[B]:ARG:NH2	1:D:67[B]:ARG:CG	2.62	0.56
1:D:92[B]:THR:HA	1:D:93:PRO:C	2.26	0.55
1:A:67[B]:ARG:CG	1:A:67[B]:ARG:NH1	2.65	0.54
2:C:6[A]:THR:CG2	8:C:353:HOH:O	2.54	0.52
1:D:37[B]:MET:HE3	1:D:39:ILE:HD11	1.94	0.49
1:A:9:CSX:OD	1:A:15[A]:THR:CB	2.48	0.47
1:D:67[A]:ARG:NH1	2:E:70:ASP:OD1	2.42	0.46
1:B:12:TYR:CZ	2:C:32:ALA:HB1	2.52	0.44
1:D:9:CSX:HA	1:D:86:CSX:OD	2.18	0.44
1:B:86:CSX:OD	1:B:97:ALA:HB3	2.17	0.44
1:B:9:CSX:OD	1:B:15:THR:CB	2.62	0.44
1:B:79[B]:GLU:OE1	8:B:303:HOH:O	2.20	0.43
1:A:3:GLN:NE2	1:B:92:THR:HG22	2.33	0.43
1:A:67[B]:ARG:HD2	8:B:365:HOH:O	2.19	0.43
1:B:35:ARG:HA	1:B:37[B]:MET:HE3	2.00	0.43
1:A:35:ARG:HD2	1:A:37[B]:MET:HE1	2.01	0.43
1:A:35:ARG:HD2	1:A:37[B]:MET:CE	2.50	0.42
2:E:92[A]:THR:HG23	8:E:350:HOH:O	2.20	0.42
2:E:92[B]:THR:HA	2:E:93:PRO:C	2.41	0.42
1:D:68[B]:MET:HE1	1:D:99:ILE:HB	2.01	0.41
3:D:201:7BN:CAB	3:D:201:7BN:CAH	2.98	0.41
1:A:68[B]:MET:HB2	1:B:31:LEU:HD23	2.03	0.41
1:D:37[B]:MET:HE2	1:D:39:ILE:HD11	1.99	0.41
1:A:32:ALA:HB1	2:E:12:TYR:CZ	2.56	0.41
1:B:37[B]:MET:HE2	1:B:37[B]:MET:HB3	1.92	0.41
1:D:9:CSX:OD	1:D:9:CSX:C	2.59	0.41
1:D:14:ASN:O	1:D:88:TRP:HA	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:D:384:HOH:O	8:D:384:HOH:O[2_556]	1.05	1.15
5:E:203:CA:CA	5:E:203:CA:CA[2_556]	1.63	0.57

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	106/103 (103%)	105 (99%)	1 (1%)	0	100	100
1	B	103/103 (100%)	102 (99%)	1 (1%)	0	100	100
1	D	105/103 (102%)	104 (99%)	1 (1%)	0	100	100
2	C	104/103 (101%)	101 (97%)	3 (3%)	0	100	100
2	E	106/103 (103%)	105 (99%)	1 (1%)	0	100	100
All	All	524/515 (102%)	517 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	94/87 (108%)	94 (100%)	0	100	100
1	B	91/87 (105%)	91 (100%)	0	100	100
1	D	93/87 (107%)	93 (100%)	0	100	100
2	C	92/89 (103%)	92 (100%)	0	100	100
2	E	94/89 (106%)	94 (100%)	0	100	100
All	All	464/439 (106%)	464 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
2	C	3	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSX	A	86	1	4,6,7	1.99	1 (25%)	2,6,8	1.03	0
1	CSX	A	9	1	4,6,7	1.58	1 (25%)	2,6,8	0.83	0
1	CSX	B	86	1	4,6,7	1.30	1 (25%)	2,6,8	1.07	0
1	CSX	B	9	1	4,6,7	0.91	0	2,6,8	1.02	0
1	CSX	D	86	1	4,6,7	2.57	1 (25%)	2,6,8	2.78	2 (100%)
1	CSX	D	9	1	4,6,7	1.49	1 (25%)	2,6,8	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSX	A	86	1	-	0/1/5/7	0/0/0/0
1	CSX	A	9	1	-	0/1/5/7	0/0/0/0
1	CSX	B	86	1	-	0/1/5/7	0/0/0/0
1	CSX	B	9	1	-	0/1/5/7	0/0/0/0
1	CSX	D	86	1	-	0/1/5/7	0/0/0/0
1	CSX	D	9	1	-	0/1/5/7	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	86	CSX	CA-C	-3.34	1.45	1.50
1	B	86	CSX	O-C	2.32	1.29	1.19
1	D	9	CSX	CB-CA	2.44	1.59	1.53
1	A	9	CSX	CA-C	2.94	1.54	1.50
1	D	86	CSX	CA-C	4.53	1.56	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	86	CSX	CA-CB-SG	-3.18	106.33	113.47
1	D	86	CSX	O-C-CA	-2.30	118.67	125.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	9	CSX	5	0
1	B	86	CSX	1	0
1	B	9	CSX	2	0
1	D	86	CSX	1	0
1	D	9	CSX	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 3 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	7BN	A	201	-	39,45,45	1.36	6 (15%)	46,65,65	1.37	6 (13%)
3	7BN	B	201	-	39,45,45	1.59	9 (23%)	46,65,65	1.53	10 (21%)
4	MRD	B	202	-	7,7,7	0.44	0	9,10,10	0.65	0
3	7BN	C	201	-	39,45,45	1.45	6 (15%)	46,65,65	1.61	11 (23%)
3	7BN	D	201	-	39,45,45	1.80	10 (25%)	46,65,65	1.84	12 (26%)
6	TRS	D	202	-	7,7,7	0.46	0	9,9,9	0.86	0
3	7BN	E	201	-	39,45,45	1.24	4 (10%)	46,65,65	1.18	4 (8%)
7	PEG	E	202	-	6,6,6	0.42	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	7BN	A	201	-	-	0/27/77/77	0/3/3/3
3	7BN	B	201	-	-	0/27/77/77	0/3/3/3
4	MRD	B	202	-	-	0/5/5/5	0/0/0/0
3	7BN	C	201	-	-	1/27/77/77	0/3/3/3
3	7BN	D	201	-	-	1/27/77/77	0/3/3/3
6	TRS	D	202	-	-	0/9/9/9	0/0/0/0
3	7BN	E	201	-	-	1/27/77/77	0/3/3/3
7	PEG	E	202	-	-	0/4/4/4	0/0/0/0

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	201	7BN	CAH-NAI	-3.97	1.31	1.35
3	B	201	7BN	NAG-NAJ	-3.11	1.29	1.34
3	D	201	7BN	C4-C5	-2.74	1.47	1.53
3	E	201	7BN	C4-C5	-2.66	1.47	1.53
3	A	201	7BN	CAH-NAI	-2.56	1.33	1.35
3	B	201	7BN	OAS-CAT	-2.51	1.39	1.44
3	B	201	7BN	CAH-CAF	-2.32	1.32	1.36
3	A	201	7BN	CAT-CAO	-2.26	1.49	1.53
3	D	201	7BN	CAC-CAB	-2.10	1.47	1.51
3	B	201	7BN	CAH-NAI	-2.06	1.33	1.35
3	A	201	7BN	OBD-CAP	2.04	1.47	1.43
3	B	201	7BN	O2-C2	2.06	1.47	1.43
3	C	201	7BN	C3-C4	2.07	1.57	1.52
3	D	201	7BN	CAE-CAF	2.10	1.55	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	201	7BN	CAV-CAU	2.12	1.57	1.53
3	A	201	7BN	O3-C3	2.14	1.47	1.43
3	D	201	7BN	CAK-CAL	2.15	1.55	1.50
3	B	201	7BN	CAU-CAT	2.16	1.55	1.52
3	A	201	7BN	O2-C2	2.40	1.48	1.43
3	E	201	7BN	O4-C4	2.42	1.48	1.43
3	B	201	7BN	C3-C4	2.45	1.58	1.52
3	D	201	7BN	OBD-CAP	2.49	1.48	1.43
3	D	201	7BN	CAQ-CAP	2.51	1.57	1.53
3	C	201	7BN	CAQ-CAP	2.64	1.57	1.53
3	C	201	7BN	CAC-CAB	2.78	1.56	1.51
3	C	201	7BN	CAV-CAU	2.80	1.59	1.53
3	E	201	7BN	O5-C1	2.81	1.51	1.44
3	D	201	7BN	C3-C4	2.87	1.59	1.52
3	C	201	7BN	O2-C2	2.90	1.49	1.43
3	B	201	7BN	CAC-CAB	3.29	1.57	1.51
3	D	201	7BN	O2-C2	3.57	1.51	1.43
3	A	201	7BN	CAP-CAO	3.77	1.56	1.53
3	B	201	7BN	CAP-CAO	4.18	1.57	1.53
3	D	201	7BN	CAV-CAU	4.40	1.62	1.53
3	D	201	7BN	O5-C1	4.49	1.55	1.44

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	201	7BN	O5-C1-CBQ	-4.90	95.33	108.31
3	B	201	7BN	C5-O5-C1	-4.47	105.62	113.12
3	D	201	7BN	C5-O5-C1	-4.08	106.26	113.12
3	C	201	7BN	C5-O5-C1	-4.05	106.32	113.12
3	D	201	7BN	CBE-NAA-CAB	-3.75	115.64	122.84
3	C	201	7BN	C4-C3-C2	-3.25	105.11	110.84
3	A	201	7BN	CAH-CAF-NAG	-3.20	106.61	111.41
3	B	201	7BN	O3-C3-C4	-3.11	103.60	110.36
3	E	201	7BN	CAV-CAU-CAT	-2.97	107.27	113.04
3	D	201	7BN	O2-C2-C3	-2.91	104.03	110.36
3	A	201	7BN	CAP-CAO-NAN	-2.83	104.56	110.40
3	B	201	7BN	O5-C1-C2	-2.81	104.48	109.66
3	B	201	7BN	O5-C1-CBQ	-2.80	100.89	108.31
3	A	201	7BN	CAH-NAI-NAJ	-2.73	106.42	112.06
3	A	201	7BN	OAS-CAT-CAU	-2.71	103.46	107.41
3	E	201	7BN	CAH-NAI-NAJ	-2.68	106.53	112.06
3	C	201	7BN	O2-C2-C3	-2.64	104.61	110.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	201	7BN	C3-C4-C5	-2.64	105.56	110.22
3	C	201	7BN	OAS-CAT-CAU	-2.52	103.75	107.41
3	D	201	7BN	OAY-CAV-CAU	-2.46	102.98	109.09
3	D	201	7BN	CAH-CAF-NAG	-2.42	107.78	111.41
3	C	201	7BN	CAK-CAL-NAN	-2.42	111.74	116.11
3	C	201	7BN	CAH-CAF-NAG	-2.39	107.84	111.41
3	B	201	7BN	CAH-CAF-NAG	-2.34	107.90	111.41
3	B	201	7BN	OAS-CAT-CAU	-2.27	104.11	107.41
3	D	201	7BN	C4-C3-C2	-2.26	106.86	110.84
3	B	201	7BN	C6-C5-C4	-2.19	107.87	113.00
3	A	201	7BN	O5-C1-CBQ	-2.12	102.69	108.31
3	B	201	7BN	C4-C3-C2	-2.10	107.14	110.84
3	C	201	7BN	CAC-CAB-NAA	-2.08	112.90	116.49
3	D	201	7BN	CAW-CAV-CAU	-2.08	107.77	112.41
3	C	201	7BN	O5-C1-C2	-2.07	105.85	109.66
3	C	201	7BN	O3-C3-C4	-2.05	105.89	110.36
3	C	201	7BN	CBE-NAA-CAB	-2.03	118.95	122.84
3	E	201	7BN	O5-C5-C4	2.39	114.06	109.66
3	D	201	7BN	O3-C3-C2	2.48	115.75	110.36
3	C	201	7BN	OAZ-CAU-CAT	2.68	115.44	109.46
3	D	201	7BN	OAZ-CAU-CAT	2.69	115.47	109.46
3	B	201	7BN	CAQ-CAR-NAI	2.85	112.49	109.35
3	D	201	7BN	O5-C5-C4	3.02	115.22	109.66
3	E	201	7BN	CAQ-CAR-NAI	3.06	112.72	109.35
3	A	201	7BN	CAQ-CAR-NAI	3.16	112.82	109.35
3	D	201	7BN	CBQ-C1-C2	4.08	120.70	114.04

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	201	7BN	CAC-CAE-CAF-NAG
3	C	201	7BN	CAC-CAE-CAF-NAG
3	D	201	7BN	CAC-CAE-CAF-NAG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	201	7BN	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	101/103 (98%)	-0.09	0 100 100	12, 16, 23, 27	0
1	B	101/103 (98%)	-0.14	0 100 100	12, 17, 25, 30	0
1	D	101/103 (98%)	-0.04	1 (0%) 82 79	10, 17, 27, 37	0
2	C	103/103 (100%)	-0.22	1 (0%) 82 79	10, 14, 24, 38	0
2	E	103/103 (100%)	-0.19	2 (1%) 67 64	11, 15, 23, 37	0
All	All	509/515 (98%)	-0.14	4 (0%) 86 83	10, 16, 25, 38	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	103	ASN	2.4
2	E	103	ASN	2.3
2	E	13[A]	HIS	2.3
2	C	103	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CSX	A	9	7/8	0.97	0.07	-	15,17,18,19	1
1	CSX	B	9	7/8	0.98	0.09	-	14,15,16,17	1
1	CSX	D	9	7/8	0.98	0.09	-	16,17,19,19	1
1	CSX	D	86	7/8	0.98	0.07	-	14,15,16,17	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	CSX	B	86	7/8	0.97	0.07	-	14,14,15,18	1
1	CSX	A	86	7/8	0.99	0.06	-	14,15,15,16	1

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	MRD	B	202	8/8	0.75	0.16	6.45	29,31,39,40	8
7	PEG	E	202	7/7	0.74	0.13	5.56	35,36,38,39	7
6	TRS	D	202	8/8	0.61	0.23	3.66	41,46,51,52	0
3	7BN	D	201	43/43	0.89	0.11	2.53	14,19,26,29	0
5	CA	C	203	1/1	0.68	0.09	1.43	60,60,60,60	0
3	7BN	C	201	43/43	0.96	0.07	1.00	10,14,16,18	0
3	7BN	B	201	43/43	0.94	0.08	0.99	14,17,21,22	0
3	7BN	A	201	43/43	0.90	0.09	0.37	16,21,23,25	0
3	7BN	E	201	43/43	0.91	0.09	-0.03	15,23,29,35	0
5	CA	C	202	1/1	0.93	0.08	-0.59	49,49,49,49	0
5	CA	E	203	1/1	0.87	0.25	-	98,98,98,98	0

6.5 Other polymers [i](#)

There are no such residues in this entry.