



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 12:07 am GMT

PDB ID : 4M0X
Title : Crystal structure of 2-chloromuconate cycloisomerase from *Rhodococcus opacus* 1CP
Authors : Ferraroni, M.; Kolomytseva, M.
Deposited on : 2013-08-02
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2259 (2.70-2.70)
Clashscore	112137	2590 (2.70-2.70)
Ramachandran outliers	110173	2550 (2.70-2.70)
Sidechain outliers	110143	2550 (2.70-2.70)
RSRZ outliers	101464	2275 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	371	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 73%, green 24%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 73% 24% .. </div> </div>
1	B	371	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 71%, green 24%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 71% 24% .. </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	A	402	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5541 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chloromuconate cycloisomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	369	Total	C	N	O	S	0	0	0
			2727	1712	475	530	10			
1	B	361	Total	C	N	O	S	0	0	0
			2619	1651	445	513	10			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	113	ASP	THR	CONFLICT	UNP Q8G9L1
A	114	ASP	SER	CONFLICT	UNP Q8G9L1
A	115	ARG	ALA	CONFLICT	UNP Q8G9L1
B	113	ASP	THR	CONFLICT	UNP Q8G9L1
B	114	ASP	SER	CONFLICT	UNP Q8G9L1
B	115	ARG	ALA	CONFLICT	UNP Q8G9L1

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mn	0	0
			1	1		
2	A	1	Total	Mn	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	106	Total 106	O 106	0	0
4	B	86	Total 86	O 86	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chloromuconate cycloisomerase



- Molecule 1: Chloromuconate cycloisomerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	121.11Å 121.11Å 110.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.70 28.71 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.7 (20.00-2.70) 98.9 (28.71-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.32 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.177 , 0.265 0.180 , 0.264	Depositor DCC
R_{free} test set	1152 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.637	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 36.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5541	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/2778	0.85	5/3781 (0.1%)
1	B	0.56	0/2664	0.82	2/3630 (0.1%)
All	All	0.58	0/5442	0.83	7/7411 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	132	ARG	NE-CZ-NH1	6.96	123.78	120.30
1	A	320	LEU	CB-CG-CD1	-6.18	100.49	111.00
1	A	250	VAL	CB-CA-C	-5.38	101.19	111.40
1	A	349	LEU	CA-CB-CG	5.33	127.56	115.30
1	B	115	ARG	NE-CZ-NH2	-5.10	117.75	120.30
1	B	18	ARG	NE-CZ-NH1	5.09	122.84	120.30
1	A	320	LEU	CA-CB-CG	5.01	126.81	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2727	0	2701	65	0
1	B	2619	0	2571	66	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
4	A	106	0	0	3	0
4	B	86	0	0	9	0
All	All	5541	0	5272	128	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (128) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:115:ARG:HG2	1:B:115:ARG:HH11	1.12	1.08
1:A:117:ARG:NH2	1:A:358:VAL:O	1.99	0.93
1:B:40:ASP:HB3	1:B:42:ASP:H	1.36	0.91
1:A:52:THR:CG2	1:A:55:GLY:H	1.86	0.88
1:B:127:LEU:HD22	4:B:503:HOH:O	1.75	0.86
1:B:115:ARG:HG2	1:B:115:ARG:NH1	1.87	0.85
1:B:39:ARG:HD3	4:B:572:HOH:O	1.77	0.84
1:B:329:PRO:HG2	1:B:341:LEU:HD22	1.60	0.83
1:A:105:ALA:O	1:A:109:THR:HB	1.81	0.81
1:B:335:GLU:HG3	4:B:511:HOH:O	1.84	0.77
1:A:250:VAL:O	1:A:274:LYS:HE2	1.89	0.73
1:B:182:ASP:O	1:B:185:GLU:HG2	1.91	0.70
1:B:141:PHE:HD2	1:B:166:PHE:CD1	2.10	0.69
1:B:117:ARG:NH2	1:B:358:VAL:O	2.26	0.67
1:B:124:SER:H	1:B:312:GLN:NE2	1.92	0.67
1:A:52:THR:HG23	1:A:55:GLY:H	1.61	0.66
1:A:165:LYS:HE3	1:A:222:GLU:OE1	1.96	0.66
1:B:184:LEU:HD23	1:B:216:ALA:O	1.96	0.65
1:A:47:ILE:HG21	1:A:360:LEU:HG	1.78	0.65
1:A:75:ALA:N	1:A:76:PRO:HD2	2.09	0.65
1:A:22:LYS:HA	1:A:27:THR:HA	1.79	0.65
1:B:335:GLU:HG2	1:B:337:VAL:HG22	1.80	0.64
1:A:67:LYS:HE3	4:A:565:HOH:O	1.98	0.64
1:B:144:GLY:O	1:B:179:ARG:HD2	1.97	0.63
1:B:153:ALA:O	1:B:157:LEU:HB2	1.99	0.63
1:A:8:MET:HA	1:A:37:ARG:O	2.00	0.62
1:B:85:THR:HG22	1:B:88:VAL:HB	1.81	0.62
1:B:141:PHE:CD2	1:B:166:PHE:CD1	2.88	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:237:ARG:HA	1:B:244:ILE:HG12	1.82	0.61
1:A:178:ARG:NH1	1:A:182:ASP:OD1	2.33	0.61
1:B:193:VAL:H	1:B:219:THR:HB	1.66	0.61
1:B:75:ALA:N	1:B:76:PRO:HD2	2.16	0.60
1:B:206:ALA:O	1:B:210:LEU:HB2	2.01	0.60
1:B:271:LYS:HG3	1:B:298:ALA:HB3	1.83	0.60
1:A:240:LEU:HD22	1:A:244:ILE:HD11	1.84	0.59
1:A:51:VAL:HG11	1:A:302:GLU:HB3	1.83	0.59
1:B:127:LEU:HD21	1:B:278:LEU:HD21	1.84	0.59
1:A:228:TRP:CD2	1:B:254:GLN:HG2	2.38	0.57
1:A:271:LYS:HG3	1:A:298:ALA:HB3	1.87	0.57
1:A:203:GLU:OE2	1:A:235:ARG:NH1	2.36	0.57
1:A:254:GLN:HG2	1:B:228:TRP:CD2	2.40	0.57
1:A:240:LEU:CD2	1:A:244:ILE:HD11	2.35	0.56
1:A:251:THR:H	1:A:255:GLN:NE2	2.04	0.55
1:A:252:THR:OG1	1:A:255:GLN:HG3	2.06	0.55
1:A:114:ASP:HB3	1:A:118:ARG:HH21	1.71	0.55
1:A:300:THR:N	1:A:301:PRO:HA	2.21	0.55
1:A:155:GLN:O	1:A:158:ASP:HB3	2.07	0.55
1:A:16:PRO:HB2	1:A:333:ILE:HD11	1.89	0.54
1:B:96:LEU:HD13	4:B:578:HOH:O	2.08	0.54
1:A:74:LEU:HD21	1:A:97:VAL:HG13	1.89	0.54
1:B:200:ARG:CD	4:B:566:HOH:O	2.55	0.54
1:B:115:ARG:HD2	4:B:539:HOH:O	2.07	0.53
1:B:219:THR:HG22	1:B:220:VAL:HG23	1.91	0.53
1:B:335:GLU:O	1:B:364:ARG:HD2	2.09	0.53
1:B:219:THR:CG2	1:B:220:VAL:HG23	2.39	0.53
1:B:232:GLY:HA2	1:B:235:ARG:NH1	2.24	0.53
1:B:271:LYS:HE2	1:B:298:ALA:HB1	1.90	0.53
1:B:56:PRO:HD3	1:B:62:SER:HB3	1.90	0.53
1:A:74:LEU:HD21	1:A:97:VAL:CG1	2.40	0.52
1:B:85:THR:CG2	1:B:88:VAL:HB	2.39	0.52
1:A:6:THR:HG22	1:A:7:GLU:HG3	1.91	0.52
1:A:40:ASP:OD1	1:A:44:GLY:N	2.43	0.51
1:B:150:ILE:HG23	1:B:186:CYS:SG	2.50	0.51
1:A:251:THR:H	1:A:255:GLN:HE21	1.58	0.50
1:A:115:ARG:HD2	1:A:118:ARG:CZ	2.42	0.50
1:A:81:ARG:HG3	1:A:82:ASP:N	2.26	0.50
1:A:31:GLN:HE21	1:A:32:SER:H	1.59	0.50
1:A:271:LYS:HD3	1:A:299:ALA:O	2.12	0.49
1:A:155:GLN:O	1:A:158:ASP:CB	2.60	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:ARG:HD2	1:A:118:ARG:NH1	2.28	0.49
1:A:253:VAL:HG22	1:A:287:ILE:CD1	2.43	0.49
1:A:299:ALA:HB1	4:A:553:HOH:O	2.13	0.49
1:B:253:VAL:HG22	1:B:287:ILE:CD1	2.43	0.49
1:A:329:PRO:HG2	1:A:341:LEU:HD12	1.96	0.48
1:B:155:GLN:O	1:B:159:VAL:HG13	2.14	0.48
1:A:49:GLU:OE1	1:A:303:SER:OG	2.31	0.47
1:A:253:VAL:HG22	1:A:287:ILE:HD11	1.96	0.47
1:B:156:LYS:HA	1:B:159:VAL:HG22	1.97	0.47
1:B:49:GLU:HB3	1:B:306:MET:HB2	1.96	0.47
1:A:94:ASP:OD1	1:A:104:LYS:NZ	2.40	0.47
1:B:302:GLU:HB2	1:B:306:MET:HB3	1.97	0.47
1:B:298:ALA:HA	1:B:325:GLU:HG2	1.97	0.46
1:A:193:VAL:H	1:A:219:THR:HB	1.78	0.46
1:B:121:ILE:HD11	1:B:126:LEU:HD21	1.98	0.46
1:A:302:GLU:CD	1:A:302:GLU:N	2.69	0.46
1:A:269:ALA:O	1:A:274:LYS:NZ	2.49	0.46
1:A:300:THR:H	1:A:301:PRO:HA	1.80	0.46
1:A:222:GLU:O	1:A:223:GLN:C	2.54	0.45
1:B:213:LEU:HB3	1:B:218:VAL:HG21	1.97	0.45
1:A:75:ALA:HB3	1:A:76:PRO:CD	2.46	0.45
1:A:8:MET:CE	1:A:75:ALA:HA	2.47	0.45
1:B:156:LYS:HB3	1:B:161:HIS:HD2	1.80	0.45
1:A:51:VAL:HG11	1:A:302:GLU:CB	2.47	0.45
1:B:52:THR:CG2	1:B:55:GLY:H	2.30	0.44
1:B:52:THR:HG21	1:B:55:GLY:CA	2.48	0.44
1:A:97:VAL:HB	1:B:56:PRO:HB2	2.00	0.44
1:A:298:ALA:HA	1:A:325:GLU:HG2	1.99	0.43
1:B:167:LYS:HE3	1:B:198:ASN:HD21	1.83	0.43
1:A:296:PHE:CD1	1:A:322:VAL:HG12	2.53	0.43
1:B:38:VAL:O	1:B:45:GLU:HA	2.19	0.43
1:B:124:SER:HB3	1:B:312:GLN:HE21	1.84	0.42
1:A:271:LYS:CG	1:A:298:ALA:HB3	2.49	0.42
1:A:193:VAL:HB	1:A:218:VAL:HA	2.02	0.42
1:B:272:ILE:N	1:B:273:PRO:CD	2.82	0.42
1:B:142:SER:C	1:B:144:GLY:H	2.24	0.41
1:A:145:SER:O	1:A:149:LEU:HG	2.21	0.41
1:B:124:SER:CB	1:B:312:GLN:HE21	2.33	0.41
1:B:15:LEU:HD21	1:B:336:VAL:HG13	2.03	0.41
1:B:85:THR:HG22	1:B:85:THR:O	2.20	0.41
1:A:259:LEU:O	1:A:260:ALA:C	2.59	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:132:ARG:HD3	4:B:557:HOH:O	2.20	0.41
1:A:179:ARG:O	1:A:182:ASP:HB2	2.20	0.41
1:B:167:LYS:NZ	4:B:564:HOH:O	2.48	0.41
1:B:282:ARG:O	1:B:282:ARG:HD2	2.20	0.41
1:B:11:THR:O	1:B:13:VAL:HG23	2.21	0.41
1:A:50:ALA:HB1	1:A:102:VAL:HG12	2.03	0.41
1:B:150:ILE:CG2	1:B:186:CYS:SG	3.09	0.41
1:A:115:ARG:HD2	1:A:118:ARG:NH2	2.36	0.41
1:A:141:PHE:HB3	1:A:149:LEU:HD22	2.02	0.41
1:A:101:SER:O	1:A:276:GLY:HA2	2.22	0.40
1:A:50:ALA:HB1	1:A:66:ILE:HD13	2.03	0.40
1:B:319:ASP:OD1	1:B:320:LEU:N	2.55	0.40
1:B:339:GLU:HA	1:B:340:PRO:HD2	1.82	0.40
1:A:99:GLY:O	1:A:104:LYS:HE3	2.21	0.40
1:A:50:ALA:HA	4:A:518:HOH:O	2.21	0.40
1:B:329:PRO:HG2	1:B:341:LEU:CD2	2.40	0.40
1:B:352:THR:HG22	4:B:585:HOH:O	2.22	0.40
1:B:75:ALA:N	1:B:76:PRO:CD	2.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	367/371 (99%)	349 (95%)	17 (5%)	1 (0%)	44	73
1	B	357/371 (96%)	322 (90%)	34 (10%)	1 (0%)	44	73
All	All	724/742 (98%)	671 (93%)	51 (7%)	2 (0%)	44	73

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	144	GLY
1	B	146	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/295 (96%)	260 (92%)	23 (8%)	14	31
1	B	266/295 (90%)	243 (91%)	23 (9%)	12	28
All	All	549/590 (93%)	503 (92%)	46 (8%)	13	29

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	ILE
1	A	45	GLU
1	A	52	THR
1	A	81	ARG
1	A	97	VAL
1	A	108	GLU
1	A	109	THR
1	A	111	LEU
1	A	159	VAL
1	A	167	LYS
1	A	178	ARG
1	A	200	ARG
1	A	204	LEU
1	A	210	LEU
1	A	219	THR
1	A	233	LEU
1	A	250	VAL
1	A	261	ASP
1	A	274	LYS
1	A	282	ARG
1	A	302	GLU
1	A	321	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	333	ILE
1	B	30	HIS
1	B	35	LEU
1	B	37	ARG
1	B	51	VAL
1	B	81	ARG
1	B	111	LEU
1	B	115	ARG
1	B	145	SER
1	B	157	LEU
1	B	184	LEU
1	B	193	VAL
1	B	204	LEU
1	B	219	THR
1	B	233	LEU
1	B	240	LEU
1	B	250	VAL
1	B	253	VAL
1	B	257	ILE
1	B	261	ASP
1	B	282	ARG
1	B	293	LEU
1	B	302	GLU
1	B	352	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	HIS
1	A	31	GLN
1	A	255	GLN
1	B	30	HIS
1	B	161	HIS
1	B	198	ASN
1	B	312	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	369/371 (99%)	-0.50	2 (0%) 90 92	18, 30, 55, 92	0
1	B	361/371 (97%)	-0.41	2 (0%) 89 90	23, 38, 65, 83	0
All	All	730/742 (98%)	-0.46	4 (0%) 90 92	18, 34, 64, 92	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	4.3
1	B	2	THR	3.0
1	A	26	THR	2.3
1	A	20	PRO	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CL	A	402	1/1	0.93	0.27	2.44	46,46,46,46	0
2	MN	B	401	1/1	0.96	0.15	-0.09	39,39,39,39	0
2	MN	A	401	1/1	0.99	0.11	-1.16	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.