



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 03:57 am GMT

PDB ID : 4M2S  
Title : Corrected Structure of Mouse P-glycoprotein bound to QZ59-RRR  
Authors : Li, J.; Jaimes, K.F.; Aller, S.G.  
Deposited on : 2013-08-05  
Resolution : 4.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 4.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1024 (5.08-3.62)
Clashscore	112137	1021 (5.08-3.70)
Ramachandran outliers	110173	1018 (5.08-3.66)
Sidechain outliers	110143	1000 (5.08-3.66)
RSRZ outliers	101464	1007 (5.08-3.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1282	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: center;">%</div> <div style="position: absolute; top: 10px; left: 0; width: 100%; text-align: center;"> <span style="display: inline-block; width: 65%; height: 10px; background-color: green;"></span> <span style="display: inline-block; width: 27%; height: 10px; background-color: yellow;"></span> <span style="display: inline-block; width: 7%; height: 10px; background-color: grey;"></span> </div> </div> </div>
1	B	1282	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: center;">%</div> <div style="position: absolute; top: 10px; left: 0; width: 100%; text-align: center;"> <span style="display: inline-block; width: 66%; height: 10px; background-color: green;"></span> <span style="display: inline-block; width: 24%; height: 10px; background-color: yellow;"></span> <span style="display: inline-block; width: 8%; height: 10px; background-color: grey;"></span> </div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	OJZ	A	6001	-	-	-	X
2	OJZ	B	6002	-	-	-	X

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 18439 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Multidrug resistance protein 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1188	Total	C	N	O	S	0	0	0
			9214	5923	1559	1694	38			
1	B	1180	Total	C	N	O	S	0	0	0
			9153	5887	1550	1678	38			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	83	GLN	ASN	CONFLICT	UNP P21447
A	87	GLN	ASN	CONFLICT	UNP P21447
A	90	GLN	ASN	CONFLICT	UNP P21447
A	1277	HIS	-	EXPRESSION TAG	UNP P21447
A	1278	HIS	-	EXPRESSION TAG	UNP P21447
A	1279	HIS	-	EXPRESSION TAG	UNP P21447
A	1280	HIS	-	EXPRESSION TAG	UNP P21447
A	1281	HIS	-	EXPRESSION TAG	UNP P21447
A	1282	HIS	-	EXPRESSION TAG	UNP P21447
B	83	GLN	ASN	CONFLICT	UNP P21447
B	87	GLN	ASN	CONFLICT	UNP P21447
B	90	GLN	ASN	CONFLICT	UNP P21447
B	1277	HIS	-	EXPRESSION TAG	UNP P21447
B	1278	HIS	-	EXPRESSION TAG	UNP P21447
B	1279	HIS	-	EXPRESSION TAG	UNP P21447
B	1280	HIS	-	EXPRESSION TAG	UNP P21447
B	1281	HIS	-	EXPRESSION TAG	UNP P21447
B	1282	HIS	-	EXPRESSION TAG	UNP P21447

- Molecule 2 is (4R,11R,18R)-4,11,18-TRI(PROPAN-2-YL)-6,13,20-TRISELENA-3,10,17,22,23,24-HEXAAZATETRACYCLO[17.2.1.1 5,8 .1 12,15 ]TETRACOSA-1(21),5(24),7,12(23),14,19(22)-HEXAENE-2,9,16-TRIONE (three-letter code: 0JZ) (formula: C<sub>24</sub>H<sub>30</sub>N<sub>6</sub>O<sub>3</sub>Se<sub>3</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	Se	0	0
			36	24	6	3	3		
2	B	1	Total	C	N	O	Se	0	0
			36	24	6	3	3		





• Molecule 1: Multidrug resistance protein 1A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.63Å 115.09Å 374.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.51 – 4.40 49.02 – 4.31	Depositor EDS
% Data completeness (in resolution range)	94.7 (24.51-4.40) 87.3 (49.02-4.31)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.43 (at 4.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.227 , 0.294 0.228 , 0.294	Depositor DCC
$R_{free}$ test set	4603 reflections (9.56%)	DCC
Wilson B-factor (Å <sup>2</sup> )	205.2	Xtriage
Anisotropy	0.356	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.23 , 112.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	18439	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	209.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 0JZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.26	0/9383	0.43	0/12685
1	B	0.26	0/9322	0.44	0/12602
All	All	0.26	0/18705	0.43	0/25287

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9214	0	9388	235	0
1	B	9153	0	9338	212	0
2	A	36	0	27	4	0
2	B	36	0	27	6	0
All	All	18439	0	18780	439	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 439 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:794:ARG:NH1	1:A:1015:ASP:OD2	1.91	1.02
1:A:46:ASP:OD1	1:A:138:ARG:NH1	2.07	0.88
1:A:433:VAL:HG13	1:A:549:LEU:HD23	1.62	0.81
1:A:1106:ARG:O	1:A:1188:ARG:NH1	2.14	0.80
1:A:801:ASP:OD2	1:A:1083:TYR:OH	2.01	0.78

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1184/1282 (92%)	1104 (93%)	72 (6%)	8 (1%)	25	68
1	B	1176/1282 (92%)	1083 (92%)	83 (7%)	10 (1%)	20	63
All	All	2360/2564 (92%)	2187 (93%)	155 (7%)	18 (1%)	22	67

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	33	VAL
1	B	1136	VAL
1	A	1036	VAL
1	A	1204	THR
1	B	32	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	979/1063 (92%)	953 (97%)	26 (3%)	50	75
1	B	973/1063 (92%)	944 (97%)	29 (3%)	46	73
All	All	1952/2126 (92%)	1897 (97%)	55 (3%)	49	74

5 of 55 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1167	ASP
1	B	277	LEU
1	B	967	PHE
1	A	1168	LYS
1	B	90	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1114	GLN
1	B	60	HIS
1	B	292	ASN
1	A	1103	GLN
1	B	139	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	OJZ	A	6001	-	24,39,39	2.08	3 (12%)	27,57,57	2.38	11 (40%)
2	OJZ	B	6002	-	24,39,39	2.05	3 (12%)	27,57,57	2.13	10 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OJZ	A	6001	-	-	0/24/48/48	0/0/4/4
2	OJZ	B	6002	-	-	0/24/48/48	0/0/4/4

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	6001	OJZ	C16-N17	4.40	1.44	1.34
2	B	6002	OJZ	C16-N17	4.68	1.44	1.34
2	B	6002	OJZ	C02-N03	5.30	1.46	1.34
2	A	6001	OJZ	C09-N10	5.60	1.46	1.34
2	B	6002	OJZ	C09-N10	5.98	1.47	1.34

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	6002	OJZ	C12-C11-N10	-4.84	105.40	112.37
2	A	6001	OJZ	C19-C18-N17	-3.60	107.18	112.37
2	B	6002	OJZ	O27-C16-N17	-2.99	117.00	122.46
2	B	6002	OJZ	C05-C04-N03	-2.76	108.39	112.37
2	A	6001	OJZ	C32-C31-C11	-2.48	109.03	111.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	6001	OJZ	4	0
2	B	6002	OJZ	6	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1188/1282 (92%)	-0.05	10 (0%) 86 80	117, 200, 297, 422	0
1	B	1180/1282 (92%)	-0.08	18 (1%) 74 66	123, 198, 304, 458	0
All	All	2368/2564 (92%)	-0.06	28 (1%) 79 71	117, 198, 303, 458	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1199	THR	5.8
1	A	1084	ASP	4.2
1	B	1162	ASN	3.5
1	B	825	THR	3.2
1	B	1198	ALA	3.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	OJZ	B	6002	36/36	0.75	0.44	0.67	159,210,345,550	0
2	OJZ	A	6001	36/36	0.62	0.42	0.54	180,230,476,517	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.