



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:04 am GMT

PDB ID : 1M4L  
Title : STRUCTURE OF NATIVE CARBOXYPEPTIDASE A AT 1.25 RESOLUTION  
Authors : Kilshtain-Vardi, A.; Glick, M.; Greenblatt, H.M.; Goldblum, A.; Shoham, G.  
Deposited on : 2002-07-03  
Resolution : 1.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1781 (1.30-1.22)
Ramachandran outliers	110173	1716 (1.30-1.22)
Sidechain outliers	110143	1714 (1.30-1.22)
RSRZ outliers	101464	1693 (1.30-1.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2838 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CARBOXYPEPTIDASE A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	307	Total	C	N	O	S	0	9	0
			2462	1574	410	473	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	228	ALA	GLU	SEE REMARK 999	UNP P00730
A	305	VAL	LEU	SEE REMARK 999	UNP P00730

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	375	Total	O	0	0
			375	375		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: CARBOXYPEPTIDASE A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.69Å 60.27Å 47.45Å 90.00° 97.47° 90.00°	Depositor
Resolution (Å)	30.00 – 1.25 32.61 – 1.25	Depositor EDS
% Data completeness (in resolution range)	86.3 (30.00-1.25) 84.8 (32.61-1.25)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.50 (at 1.25Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.104 , 0.145 0.105 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	11.9	Xtriage
Anisotropy	0.371	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 65.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	2838	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.76	2/2576 (0.1%)	1.37	33/3500 (0.9%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	194	SER	CB-OG	-5.44	1.35	1.42
1	A	34	SER	CB-OG	-5.37	1.35	1.42

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	276[A]	ARG	CD-NE-CZ	16.94	147.32	123.60
1	A	276[B]	ARG	CD-NE-CZ	16.94	147.32	123.60
1	A	276[A]	ARG	NE-CZ-NH1	12.19	126.39	120.30
1	A	276[B]	ARG	NE-CZ-NH1	12.19	126.39	120.30
1	A	2[A]	ARG	NE-CZ-NH2	11.61	126.11	120.30
1	A	2[B]	ARG	NE-CZ-NH2	11.61	126.11	120.30
1	A	40	ARG	NE-CZ-NH1	-10.85	114.87	120.30
1	A	20	ASP	CB-CG-OD1	8.95	126.35	118.30
1	A	15	LEU	CB-CG-CD2	8.86	126.07	111.00
1	A	16	ASP	CB-CG-OD1	8.16	125.64	118.30
1	A	82	PHE	CB-CG-CD1	7.95	126.37	120.80
1	A	238	TYR	CB-CG-CD2	-7.68	116.39	121.00
1	A	82	PHE	CB-CG-CD2	-7.37	115.64	120.80
1	A	127	ARG	NE-CZ-NH1	-7.28	116.66	120.30
1	A	127	ARG	CD-NE-CZ	7.23	133.72	123.60
1	A	19	TYR	CB-CG-CD1	7.00	125.20	121.00
1	A	59	ARG	NE-CZ-NH2	-6.80	116.90	120.30
1	A	89	ASP	CB-CG-OD1	6.54	124.19	118.30
1	A	86	PHE	CB-CG-CD2	-6.33	116.37	120.80

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	133	THR	N-CA-CB	6.14	121.96	110.30
1	A	40	ARG	NH1-CZ-NH2	6.08	126.09	119.40
1	A	145	ARG	NE-CZ-NH1	-5.68	117.46	120.30
1	A	19	TYR	CB-CG-CD2	-5.63	117.62	121.00
1	A	238	TYR	CB-CG-CD1	5.58	124.34	121.00
1	A	185	ASP	CB-CG-OD1	5.53	123.27	118.30
1	A	86	PHE	CB-CG-CD1	5.41	124.59	120.80
1	A	240	TYR	CB-CG-CD1	-5.28	117.83	121.00
1	A	248	TYR	CB-CG-CD2	5.19	124.11	121.00
1	A	127	ARG	NH1-CZ-NH2	5.12	125.03	119.40
1	A	280	LEU	CB-CG-CD1	5.12	119.70	111.00
1	A	2[A]	ARG	NE-CZ-NH1	-5.10	117.75	120.30
1	A	2[B]	ARG	NE-CZ-NH1	-5.10	117.75	120.30
1	A	16	ASP	CB-CG-OD2	-5.05	113.75	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2462	0	2367	33	0
2	A	1	0	0	0	0
3	A	375	0	0	14	0
All	All	2838	0	2367	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:SER:HB3	3:A:1682:HOH:O	1.58	1.04
1:A:15:LEU:HG	3:A:1683:HOH:O	1.63	0.98
1:A:276[B]:ARG:NE	3:A:1625:HOH:O	2.00	0.94

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:HIS:HD2	1:A:188:ASN:H	1.27	0.79
1:A:298:LEU:HG	1:A:302[B]:GLU:OE1	1.86	0.76
1:A:239:LYS:HD3	3:A:1407:HOH:O	1.88	0.73
1:A:168:LYS:HG2	3:A:1499:HOH:O	1.88	0.73
1:A:276[B]:ARG:CZ	3:A:1625:HOH:O	2.35	0.71
1:A:62:ILE:HG13	3:A:1678:HOH:O	1.93	0.67
1:A:137:LEU:HD22	1:A:137:LEU:H	1.61	0.65
1:A:62:ILE:HD12	1:A:191:ALA:HB3	1.79	0.64
1:A:303:HIS:HB2	3:A:1511:HOH:O	1.97	0.64
1:A:186:HIS:CD2	1:A:188:ASN:H	2.17	0.54
1:A:272:ARG:HH11	1:A:285:GLN:HE21	1.56	0.54
1:A:137:LEU:HD22	3:A:1645:HOH:O	2.08	0.53
1:A:133:THR:HG23	3:A:1512:HOH:O	2.11	0.51
1:A:186:HIS:HE1	3:A:1357:HOH:O	1.96	0.48
1:A:137:LEU:N	1:A:137:LEU:HD22	2.28	0.48
1:A:15:LEU:HD21	1:A:19:TYR:CE2	2.49	0.46
1:A:286:ILE:HG22	1:A:287:ILE:HD13	1.97	0.46
1:A:127:ARG:NH2	3:A:1497:HOH:O	2.49	0.45
1:A:71:ARG:HD3	1:A:125:LEU:O	2.17	0.44
1:A:272:ARG:HH11	1:A:285:GLN:NE2	2.15	0.44
1:A:42:TYR:OH	1:A:132:VAL:HG22	2.19	0.42
1:A:171:ASN:HB3	1:A:177:LYS:HD3	2.01	0.42
1:A:86:PHE:HE1	1:A:294:TRP:CZ3	2.38	0.42
1:A:276[B]:ARG:NH2	3:A:1625:HOH:O	2.51	0.42
1:A:40:ARG:HD3	1:A:40:ARG:HH11	1.58	0.41
1:A:62:ILE:CG1	3:A:1678:HOH:O	2.63	0.41
1:A:190:LYS:HA	1:A:190:LYS:HD3	1.94	0.40
1:A:218:GLU:OE1	1:A:264:LYS:HE2	2.20	0.40
1:A:33:VAL:HA	1:A:51:LYS:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	314/307 (102%)	306 (98%)	7 (2%)	1 (0%)	44 16

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	199	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	272/263 (103%)	263 (97%)	9 (3%)	43 6

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3[A]	SER
1	A	3[B]	SER
1	A	33	VAL
1	A	135	SER
1	A	137	LEU
1	A	239	LYS
1	A	280	LEU
1	A	287	ILE
1	A	295	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	31	GLN
1	A	37	GLN
1	A	171	ASN
1	A	186	HIS
1	A	220	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	249	GLN
1	A	285	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	307/307 (100%)	-0.42	10 (3%) 47 40	9, 15, 35, 69	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	135	SER	4.8
1	A	137	LEU	4.4
1	A	133	THR	3.6
1	A	307	ASN	3.5
1	A	248	TYR	3.2
1	A	306	ASN	2.5
1	A	57[A]	SER	2.3
1	A	1	ALA	2.2
1	A	134[A]	SER	2.2
1	A	31	GLN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	A	1308	1/1	1.00	0.01	-6.77	10,10,10,10	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.