



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 01:00 am GMT

PDB ID : 1M5R
Title : Ternary complex of T4 phage BGT with UDP and a 13 mer DNA duplex
Authors : Lariviere, L.; Morera, S.
Deposited on : 2002-07-10
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	13	<div> <div>8%</div> <div>92%</div> <div>8%</div> </div>
1	D	13	<div> <div>15%</div> <div>77%</div> <div>23%</div> </div>
2	E	13	<div> <div>38%</div> <div>69%</div> <div>31%</div> </div>
2	F	13	<div> <div>23%</div> <div>54%</div> <div>46%</div> </div>
3	A	351	<div> <div>2%</div> <div>83%</div> <div>16%</div> <div>.</div> </div>
3	B	351	<div> <div>%</div> <div>80%</div> <div>19%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	TRS	A	501	-	-	-	X
5	TRS	B	501	-	-	-	X
5	TRS	B	502	-	-	-	X
6	MPD	A	502	-	-	-	X
6	MPD	B	504	-	-	-	X
6	MPD	B	505	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7662 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*GP*AP*TP*AP*CP*TP*3DRP*AP*GP*AP*TP*AP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	13	Total	C	N	O	P	0	0	0
			258	124	49	73	12			
1	D	13	Total	C	N	O	P	0	0	0
			258	124	49	73	12			


- Molecule 2 is a DNA chain called 5'-D(*CP*TP*AP*TP*CP*TP*GP*AP*GP*TP*AP*TP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	13	Total	C	N	O	P	15	0	0
			261	127	44	78	12			
2	F	13	Total	C	N	O	P	0	0	0
			261	127	44	78	12			

- Molecule 3 is a protein called DNA beta-glucosyltransferase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	351	Total	C	N	O	S	0	0	0
			2869	1854	479	525	11			
3	B	351	Total	C	N	O	S	0	0	0
			2869	1854	479	525	11			

- Molecule 4 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C₉H₁₄N₂O₁₂P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
4	B	1	Total	C	N	O	P	0	0
			25	9	2	12	2		

- Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			8	4	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			8	4	1	3		
5	A	1	Total	C	N	O	0	0
			8	4	1	3		
5	B	1	Total	C	N	O	0	0
			8	4	1	3		
5	B	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 6 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	344	Total	O	0	0
			344	344		
7	B	352	Total	O	0	0
			352	352		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	29	Total 29	O 29	0	0
7	D	24	Total 24	O 24	0	0
7	E	18	Total 18	O 18	0	0
7	F	5	Total 5	O 5	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: 5'-D(*GP*AP*TP*AP*CP*TP*3DRP*AP*GP*AP*TP*AP*G)-3'




- Molecule 1: 5'-D(*GP*AP*TP*AP*CP*TP*3DRP*AP*GP*AP*TP*AP*G)-3'





- Molecule 2: 5'-D(*CP*TP*AP*TP*CP*TP*GP*AP*GP*TP*AP*TP*C)-3'




- Molecule 2: 5'-D(*CP*TP*AP*TP*CP*TP*GP*AP*GP*TP*AP*TP*C)-3'



- Molecule 3: DNA beta-glucosyltransferase



● Molecule 3: DNA beta-glucosyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	83.00Å 55.90Å 99.50Å 90.00° 91.10° 90.00°	Depositor
Resolution (Å)	20.00 – 1.80 19.90 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.1 (20.00-1.80) 98.0 (19.90-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 1.80Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.191 , 0.224 0.192 , 0.226	Depositor DCC
R_{free} test set	4209 reflections (5.06%)	DCC
Wilson B-factor (Å ²)	23.6	Xtriage
Anisotropy	0.177	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 49.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7662	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, UDP, TRS, 3DR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.24	0/277	0.62	0/424
1	D	0.21	0/277	0.60	0/424
2	E	0.25	0/291	0.65	0/447
2	F	0.24	0/291	0.65	0/447
3	A	0.33	0/2931	0.59	1/3953 (0.0%)
3	B	0.33	0/2931	0.60	1/3953 (0.0%)
All	All	0.32	0/6998	0.60	2/9648 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	266	ILE	N-CA-C	-6.05	94.67	111.00
3	B	266	ILE	N-CA-C	-5.37	96.50	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	258	0	145	1	0
1	D	258	0	145	3	0
2	E	261	0	150	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	261	0	150	6	0
3	A	2869	0	2903	42	0
3	B	2869	0	2903	60	0
4	A	25	0	11	0	0
4	B	25	0	11	1	0
5	A	16	0	24	0	0
5	B	24	0	36	0	0
6	A	8	0	13	0	0
6	B	16	0	26	3	0
7	A	344	0	0	3	0
7	B	352	0	0	8	0
7	C	29	0	0	0	0
7	D	24	0	0	1	0
7	E	18	0	0	0	0
7	F	5	0	0	0	0
All	All	7662	0	6517	120	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (120) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:217:ARG:H	3:A:220:GLN:HE21	1.22	0.85
3:A:217:ARG:H	3:A:220:GLN:NE2	1.73	0.85
2:E:24:DA:H2''	2:E:25:DT:H5''	1.59	0.85
6:B:505:MPD:HM2	6:B:505:MPD:H52	1.58	0.83
3:B:217:ARG:H	3:B:220:GLN:HE21	1.26	0.81
3:A:343:ASP:OD1	3:A:347:LYS:HE2	1.82	0.79
3:B:84:GLN:HE22	3:B:108:SER:H	1.37	0.72
3:A:155:ILE:HG13	3:A:156:GLU:HG3	1.71	0.72
3:B:127:LEU:HD23	3:B:129:LYS:NZ	2.06	0.70
3:A:55:ASP:HB3	3:A:58:ASP:OD2	1.91	0.69
3:A:84:GLN:HE22	3:A:108:SER:H	1.42	0.68
3:B:111:ASN:HD21	3:B:115:ARG:HH21	1.41	0.68
2:E:24:DA:C2'	2:E:25:DT:H5''	2.25	0.66
3:B:253:ALA:HB3	3:B:273:THR:HG23	1.80	0.63
3:B:14:ASN:OD1	3:B:16:LYS:HD3	1.99	0.62
3:A:215:ASN:ND2	3:A:237:LYS:HD3	2.17	0.60
2:E:14:DC:C2'	2:E:15:DT:H71	2.31	0.60
3:A:2:LYS:NZ	3:A:36:ASN:HD21	2.00	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:215:ASN:HD21	3:A:237:LYS:HD3	1.69	0.58
3:A:184:VAL:HB	3:A:210:ILE:HG12	1.86	0.57
3:B:111:ASN:C	3:B:111:ASN:HD22	2.07	0.57
3:A:75:LYS:HB2	3:A:75:LYS:NZ	2.20	0.57
3:B:111:ASN:ND2	3:B:115:ARG:HE	2.02	0.57
2:E:24:DA:H2''	2:E:25:DT:C5'	2.34	0.56
3:A:253:ALA:HB3	3:A:273:THR:HG23	1.88	0.56
3:B:150:LYS:HB2	3:B:150:LYS:HZ2	1.71	0.56
2:E:14:DC:H2''	2:E:15:DT:H71	1.87	0.56
3:B:111:ASN:HD21	3:B:115:ARG:HE	1.54	0.55
3:B:14:ASN:CG	3:B:16:LYS:HD3	2.27	0.55
3:B:225:LYS:NZ	3:B:225:LYS:HB2	2.21	0.55
3:A:312:GLU:HG3	7:A:1119:HOH:O	2.07	0.55
3:B:111:ASN:HD21	3:B:115:ARG:NH2	2.04	0.55
2:F:23:DT:H2''	2:F:24:DA:H5'	1.89	0.54
6:B:505:MPD:HM2	6:B:505:MPD:C5	2.35	0.52
3:A:109:TRP:HB3	3:A:110:PRO:HD3	1.89	0.52
3:B:109:TRP:CZ2	3:B:123:GLU:HB2	2.44	0.52
3:B:18:VAL:HB	3:B:19:PRO:HD3	1.92	0.52
3:A:2:LYS:NZ	3:A:36:ASN:ND2	2.58	0.52
3:B:305:GLU:HG3	7:B:2300:HOH:O	2.10	0.52
2:F:24:DA:H2''	2:F:25:DT:O5'	2.09	0.52
3:B:217:ARG:H	3:B:220:GLN:NE2	2.01	0.51
3:B:225:LYS:HZ2	3:B:225:LYS:HB2	1.74	0.51
3:B:289:LYS:NZ	7:B:2249:HOH:O	2.33	0.51
3:A:287:ASP:OD1	3:A:291:ARG:HG2	2.11	0.51
3:B:109:TRP:HB3	3:B:110:PRO:HD3	1.93	0.51
3:B:76:PRO:HG3	3:B:117:TRP:CD2	2.46	0.51
1:C:1:DG:O6	3:B:198:LYS:HG2	2.11	0.51
3:B:127:LEU:HD23	3:B:129:LYS:HZ2	1.76	0.50
1:D:7:3DR:H2''	1:D:8:DA:OP1	2.11	0.50
3:B:308:ASP:O	3:B:312:GLU:HG3	2.12	0.50
3:A:77:ASN:HB3	3:A:80:ILE:HG22	1.94	0.50
2:E:14:DC:H2'	2:E:15:DT:H71	1.93	0.49
1:D:2:DA:H2'	7:D:2296:HOH:O	2.12	0.49
3:B:219:LYS:HE3	7:B:2163:HOH:O	2.12	0.49
1:D:7:3DR:H1'2	7:B:2115:HOH:O	2.12	0.49
3:B:69:ILE:HD12	3:B:106:SER:OG	2.13	0.49
3:B:117:TRP:HA	3:B:119:TYR:CE1	2.48	0.48
3:A:141:LEU:O	3:A:145:LYS:HG3	2.12	0.48
2:F:25:DT:H2'	2:F:26:DC:C6	2.48	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:200:VAL:HA	3:B:204:PHE:CD1	2.49	0.47
2:F:14:DC:H2'	2:F:15:DT:H72	1.94	0.47
3:A:2:LYS:HZ2	3:A:36:ASN:ND2	2.13	0.47
3:B:308:ASP:HB2	7:B:2260:HOH:O	2.14	0.46
3:A:76:PRO:HG3	3:A:117:TRP:CD2	2.49	0.46
3:B:209:ASN:HB3	7:B:2377:HOH:O	2.14	0.46
3:A:53:GLU:HG2	7:A:1336:HOH:O	2.16	0.46
3:B:150:LYS:HB2	3:B:150:LYS:NZ	2.31	0.45
3:A:216:ALA:HA	3:A:220:GLN:HE22	1.82	0.45
3:A:187:GLY:HA2	3:A:213:PHE:O	2.17	0.45
3:B:4:ALA:HB2	3:B:59:TYR:CD1	2.52	0.45
3:A:90:TYR:CE2	3:A:92:SER:HB2	2.52	0.45
3:A:84:GLN:NE2	3:A:108:SER:H	2.12	0.45
3:A:219:LYS:HB3	3:A:219:LYS:NZ	2.32	0.44
3:A:2:LYS:HG3	3:A:59:TYR:CD1	2.53	0.44
3:B:111:ASN:HD21	3:B:115:ARG:NE	2.15	0.44
3:B:127:LEU:HD23	3:B:129:LYS:HZ1	1.82	0.44
3:B:300:VAL:HG12	3:B:305:GLU:HG2	1.98	0.44
3:A:4:ALA:HB2	3:A:59:TYR:CG	2.52	0.44
3:B:187:GLY:HA2	3:B:213:PHE:O	2.18	0.44
3:B:84:GLN:NE2	3:B:108:SER:H	2.11	0.44
3:A:289:LYS:HE2	7:A:1390:HOH:O	2.18	0.44
3:A:122:THR:OG1	3:A:125:GLU:HG3	2.17	0.44
3:B:331:LEU:HD23	3:B:331:LEU:C	2.38	0.44
2:F:14:DC:C2'	2:F:15:DT:H72	2.48	0.44
3:B:163:GLU:HG3	3:B:266:ILE:O	2.17	0.43
3:B:69:ILE:HD11	3:B:106:SER:O	2.17	0.43
3:B:245:GLU:HG2	7:B:2098:HOH:O	2.19	0.43
3:A:4:ALA:HB2	3:A:59:TYR:CD1	2.52	0.43
3:B:292:ILE:HG22	3:B:293:ILE:HG13	2.00	0.43
3:B:184:VAL:HB	3:B:210:ILE:HG12	1.99	0.43
3:B:225:LYS:HD2	3:B:226:TYR:CZ	2.54	0.43
3:B:183:ASP:OD1	6:B:504:MPD:H13	2.17	0.43
3:A:329:ASP:OD1	3:A:333:LYS:HE2	2.19	0.43
3:A:8:MET:O	3:A:80:ILE:HA	2.18	0.43
3:B:347:LYS:HD3	3:B:347:LYS:C	2.39	0.43
3:B:160:PHE:CZ	3:B:345:PHE:HB2	2.53	0.43
3:B:228:TRP:CZ3	3:B:231:ALA:HB2	2.54	0.43
3:A:14:ASN:OD1	3:A:16:LYS:HB2	2.19	0.42
3:A:75:LYS:HZ3	3:A:75:LYS:HB2	1.83	0.42
3:B:4:ALA:HB2	3:B:59:TYR:CG	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:331:LEU:C	3:A:331:LEU:HD23	2.39	0.42
3:A:2:LYS:HZ3	3:A:36:ASN:HD21	1.66	0.42
3:B:27:PHE:CB	3:B:39:ILE:HD11	2.50	0.42
3:A:310:VAL:O	3:A:314:LYS:HG3	2.20	0.42
3:B:30:ILE:O	3:B:33:MET:HB2	2.20	0.42
3:B:216:ALA:HA	3:B:220:GLN:HE22	1.83	0.42
3:A:2:LYS:HZ2	3:A:36:ASN:HD21	1.65	0.42
3:B:276:SER:O	3:B:328:HIS:HE1	2.03	0.42
3:B:111:ASN:C	3:B:111:ASN:ND2	2.72	0.41
2:F:23:DT:H1'	2:F:24:DA:H5''	2.02	0.41
3:B:195:ARG:CZ	3:B:257:GLY:HA2	2.50	0.41
3:A:217:ARG:N	3:A:220:GLN:NE2	2.53	0.41
3:A:90:TYR:CZ	3:A:92:SER:HB2	2.55	0.41
3:B:225:LYS:NZ	3:B:225:LYS:CB	2.83	0.41
3:B:243:VAL:HG21	4:B:401:UDP:H2'	2.03	0.41
3:A:160:PHE:CZ	3:A:345:PHE:HB2	2.55	0.40
3:B:149:LYS:O	3:B:149:LYS:HD3	2.22	0.40
3:B:329:ASP:OD2	3:B:333:LYS:HE2	2.21	0.40
2:E:14:DC:H2''	2:E:15:DT:C6	2.56	0.40
3:B:308:ASP:HB3	7:B:2300:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	349/351 (99%)	343 (98%)	6 (2%)	0	100	100
3	B	349/351 (99%)	339 (97%)	10 (3%)	0	100	100
All	All	698/702 (99%)	682 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	317/317 (100%)	310 (98%)	7 (2%)	57	44
3	B	317/317 (100%)	312 (98%)	5 (2%)	68	58
All	All	634/634 (100%)	622 (98%)	12 (2%)	62	50

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	36	ASN
3	A	66	ASN
3	A	78	LEU
3	A	166	LYS
3	A	212	PHE
3	A	219	LYS
3	A	317	ASP
3	B	36	ASN
3	B	66	ASN
3	B	111	ASN
3	B	212	PHE
3	B	268	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	10	ASN
3	A	36	ASN
3	A	84	GLN
3	A	170	ASN
3	A	194	GLN
3	A	215	ASN
3	A	220	GLN
3	A	260	ASN
3	A	328	HIS
3	A	332	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	B	10	ASN
3	B	36	ASN
3	B	84	GLN
3	B	111	ASN
3	B	170	ASN
3	B	194	GLN
3	B	220	GLN
3	B	260	ASN
3	B	328	HIS
3	B	332	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	3DR	C	7	1	8,11,12	0.60	0	8,14,17	0.71	0
1	3DR	D	7	1	8,11,12	0.68	0	8,14,17	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	3DR	C	7	1	-	0/3/15/16	0/1/1/1
1	3DR	D	7	1	-	0/3/15/16	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	7	3DR	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	UDP	A	400	-	21,26,26	1.10	2 (9%)	22,40,40	3.35	2 (9%)
5	TRS	A	500	-	7,7,7	1.68	2 (28%)	9,9,9	1.98	4 (44%)
5	TRS	A	501	-	7,7,7	1.58	2 (28%)	9,9,9	2.10	4 (44%)
6	MPD	A	502	-	7,7,7	1.21	1 (14%)	9,10,10	0.32	0
4	UDP	B	401	-	21,26,26	1.04	2 (9%)	22,40,40	3.35	2 (9%)
5	TRS	B	501	-	7,7,7	1.76	3 (42%)	9,9,9	1.97	3 (33%)
5	TRS	B	502	-	7,7,7	1.75	2 (28%)	9,9,9	1.88	4 (44%)
5	TRS	B	503	-	7,7,7	1.88	3 (42%)	9,9,9	1.87	3 (33%)
6	MPD	B	504	-	7,7,7	1.20	1 (14%)	9,10,10	0.30	0
6	MPD	B	505	-	7,7,7	1.29	1 (14%)	9,10,10	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	UDP	A	400	-	-	0/12/32/32	0/2/2/2
5	TRS	A	500	-	-	0/9/9/9	0/0/0/0
5	TRS	A	501	-	-	0/9/9/9	0/0/0/0
6	MPD	A	502	-	-	0/5/5/5	0/0/0/0
4	UDP	B	401	-	-	0/12/32/32	0/2/2/2
5	TRS	B	501	-	-	0/9/9/9	0/0/0/0
5	TRS	B	502	-	-	0/9/9/9	0/0/0/0
5	TRS	B	503	-	-	0/9/9/9	0/0/0/0
6	MPD	B	504	-	-	0/5/5/5	0/0/0/0
6	MPD	B	505	-	-	0/5/5/5	0/0/0/0

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	505	MPD	O2-C2	-3.16	1.36	1.44
6	A	502	MPD	O2-C2	-3.08	1.36	1.44
6	B	504	MPD	O2-C2	-3.06	1.36	1.44
5	B	503	TRS	C-N	-2.14	1.42	1.49
5	A	500	TRS	C-N	-2.13	1.42	1.49
5	B	501	TRS	C-N	-2.12	1.42	1.49
5	A	501	TRS	C-N	-2.05	1.42	1.49
4	B	401	UDP	PB-O3A	2.06	1.63	1.60
5	B	501	TRS	C2-C	2.22	1.58	1.52
5	B	502	TRS	C2-C	2.22	1.58	1.52
5	B	503	TRS	C2-C	2.27	1.58	1.52
4	A	400	UDP	PB-O3A	2.33	1.63	1.60
5	A	501	TRS	C1-C	2.75	1.59	1.52
4	B	401	UDP	C4-N3	2.90	1.38	1.33
5	A	500	TRS	C1-C	3.01	1.59	1.52
5	B	501	TRS	C1-C	3.05	1.60	1.52
5	B	502	TRS	C1-C	3.10	1.60	1.52
4	A	400	UDP	C4-N3	3.23	1.38	1.33
5	B	503	TRS	C1-C	3.29	1.60	1.52

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	400	UDP	C5-C4-N3	-3.55	114.65	123.12
4	B	401	UDP	C5-C4-N3	-3.53	114.69	123.12
5	B	503	TRS	C1-C-N	-2.58	102.23	107.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	501	TRS	C2-C-C1	-2.43	104.17	111.06
5	A	500	TRS	O1-C1-C	-2.42	103.47	110.47
5	B	501	TRS	O1-C1-C	-2.41	103.48	110.47
5	A	501	TRS	O1-C1-C	-2.39	103.55	110.47
5	A	500	TRS	C2-C-C1	-2.26	104.66	111.06
5	B	502	TRS	O1-C1-C	-2.21	104.08	110.47
5	B	502	TRS	C2-C-C1	-2.07	105.19	111.06
5	B	503	TRS	C2-C-N	2.27	112.56	107.73
5	B	503	TRS	C3-C-N	2.50	113.04	107.73
5	B	502	TRS	C2-C-N	2.70	113.47	107.73
5	A	500	TRS	C3-C-N	2.81	113.70	107.73
5	B	501	TRS	C2-C-N	2.95	114.00	107.73
5	A	500	TRS	C2-C-N	3.13	114.37	107.73
5	B	502	TRS	C3-C-N	3.17	114.47	107.73
5	B	501	TRS	C3-C-N	3.19	114.51	107.73
5	A	501	TRS	C3-C-N	3.27	114.68	107.73
5	A	501	TRS	C2-C-N	3.33	114.80	107.73
4	A	400	UDP	C4-N3-C2	14.97	126.99	114.13
4	B	401	UDP	C4-N3-C2	15.08	127.08	114.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	401	UDP	1	0
6	B	504	MPD	1	0
6	B	505	MPD	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	12/13 (92%)	0.37	1 (8%) 12 10	26, 36, 42, 48	0
1	D	12/13 (92%)	1.01	2 (16%) 2 1	29, 43, 64, 66	0
2	E	13/13 (100%)	1.66	5 (38%) 0 0	31, 43, 71, 75	1 (7%)
2	F	13/13 (100%)	1.45	3 (23%) 1 0	34, 47, 58, 61	0
3	A	351/351 (100%)	0.11	8 (2%) 61 57	14, 21, 35, 47	0
3	B	351/351 (100%)	0.05	5 (1%) 75 72	13, 21, 31, 41	0
All	All	752/754 (99%)	0.15	24 (3%) 48 43	13, 22, 41, 75	1 (0%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	26	DC	5.8
3	A	149	LYS	4.5
3	A	119	TYR	3.4
3	A	152	ASP	3.4
3	A	315	HIS	3.1
2	E	25	DT	3.0
3	B	315	HIS	2.8
3	B	46	VAL	2.8
2	F	26	DC	2.7
2	E	23	DT	2.7
1	D	13	DG	2.7
2	F	25	DT	2.6
3	A	116	PRO	2.5
1	D	12	DA	2.5
3	A	120	LEU	2.5
2	F	22	DG	2.4
3	B	203	LEU	2.4
3	A	155	ILE	2.3
3	B	149	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	E	22	DG	2.1
3	A	151	VAL	2.1
2	E	24	DA	2.1
3	B	16	LYS	2.1
1	C	13	DG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	3DR	C	7	11/12	0.92	0.18	-	29,33,34,35	0
1	3DR	D	7	11/12	0.91	0.16	-	26,34,36,37	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	MPD	A	502	8/8	0.71	0.29	10.56	46,47,49,49	0
6	MPD	B	504	8/8	0.49	0.34	9.63	45,47,50,51	0
6	MPD	B	505	8/8	0.74	0.29	9.03	48,49,49,49	0
5	TRS	A	501	8/8	0.84	0.31	6.52	69,69,70,70	0
5	TRS	B	502	8/8	0.81	0.18	3.86	25,27,29,31	0
5	TRS	B	501	8/8	0.87	0.16	2.55	26,28,30,31	0
4	UDP	B	401	25/25	0.98	0.09	-0.76	15,17,19,19	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	TRS	A	500	8/8	0.97	0.08	-1.36	16,19,20,21	0
4	UDP	A	400	25/25	0.98	0.07	-1.68	12,13,15,16	0
5	TRS	B	503	8/8	0.65	0.35	-	62,63,63,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.