



Full wwPDB NMR Structure Validation Report ⓘ

Feb 13, 2017 – 12:52 am GMT

PDB ID : 2MAF
Title : Solution structure of opa60 from n. gonorrhoeae
Authors : Columbus, L.; Fox, D.A.; Larsson, P.; Lo, R.H.; Kroncke, B.M.; Kasson, P.M.
Deposited on : 2013-07-08

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk28760
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 33%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	238	

2 Ensemble composition and analysis

This entry contains 20 models. The atoms present in the NMR models are not consistent. Some calculations may have failed as a result. All residues are included in the validation scores. Model 20 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:7-A:15, A:55-A:72, A:117-A:142, A:190-A:217, A:230-A:238 (90)	0.94	20

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 4 single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 19, 20
2	16, 17
Single-model clusters	3; 11; 15; 18

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 3718 atoms, of which 1830 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Opacity protein opA60.

Mol	Chain	Residues	Atoms					Trace
1	A	233	Total	C	H	N	O	0
			3718	1185	1830	352	351	

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Opacity protein opA60



4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Opacity protein opA60




4.2.2 Score per residue for model 2

- Molecule 1: Opacity protein opA60



4.2.3 Score per residue for model 3

- Molecule 1: Opacity protein opA60



4.2.4 Score per residue for model 4

- Molecule 1: Opacity protein opA60




4.2.5 Score per residue for model 5

- Molecule 1: Opacity protein opA60




4.2.6 Score per residue for model 6

- Molecule 1: Opacity protein opA60




4.2.7 Score per residue for model 7

- Molecule 1: Opacity protein opA60



4.2.8 Score per residue for model 8

- Molecule 1: Opacity protein opA60



4.2.9 Score per residue for model 9

- Molecule 1: Opacity protein opA60




4.2.10 Score per residue for model 10

- Molecule 1: Opacity protein opA60



4.2.11 Score per residue for model 11

- Molecule 1: Opacity protein opA60




4.2.12 Score per residue for model 12

- Molecule 1: Opacity protein opA60




4.2.13 Score per residue for model 13

- Molecule 1: Opacity protein opA60




4.2.14 Score per residue for model 14

- Molecule 1: Opacity protein opA60




4.2.15 Score per residue for model 15

- Molecule 1: Opacity protein opA60




4.2.17 Score per residue for model 17

- Molecule 1: Opacity protein opA60



4.2.18 Score per residue for model 18

- Molecule 1: Opacity protein opA60



4.2.19 Score per residue for model 19

- Molecule 1: Opacity protein opA60



4.2.20 Score per residue for model 20 (medoid)

- Molecule 1: Opacity protein opA60

Chain A:

35%

60%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 400 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	2.31
GROMACS	refinement	4.5

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2maf_cs.str
Number of chemical shift lists	1
Total number of shifts	530
Number of shifts mapped to atoms	530
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	33%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.83±0.01	0±0/732 (0.0±0.0%)	1.27±0.09	8±2/992 (0.8±0.2%)
All	All	0.83	0/14640 (0.0%)	1.28	166/19840 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	2.4±1.8
All	All	0	48

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	237	ARG	NE-CZ-NH2	-15.32	112.64	120.30	18	11
1	A	237	ARG	NE-CZ-NH1	14.65	127.63	120.30	16	14
1	A	235	ARG	NE-CZ-NH1	14.54	127.57	120.30	7	15
1	A	213	ARG	NE-CZ-NH1	13.87	127.24	120.30	4	15
1	A	66	ARG	NE-CZ-NH1	12.87	126.74	120.30	11	17
1	A	235	ARG	NE-CZ-NH2	-12.83	113.89	120.30	8	11
1	A	55	ARG	NE-CZ-NH2	-12.77	113.91	120.30	19	8
1	A	66	ARG	NE-CZ-NH2	-11.16	114.72	120.30	20	8
1	A	213	ARG	NE-CZ-NH2	-11.12	114.74	120.30	15	9
1	A	55	ARG	NE-CZ-NH1	10.63	125.62	120.30	13	15
1	A	138	ARG	NE-CZ-NH1	10.47	125.53	120.30	17	11
1	A	134	TYR	CB-CG-CD2	-10.11	114.94	121.00	16	1
1	A	7	ARG	NE-CZ-NH2	-9.94	115.33	120.30	8	1
1	A	7	ARG	NE-CZ-NH1	9.56	125.08	120.30	13	6
1	A	209	ASP	CB-CG-OD1	7.56	125.11	118.30	3	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	134	TYR	CB-CG-CD1	7.40	125.44	121.00	16	2
1	A	212	TYR	CB-CG-CD2	-6.83	116.90	121.00	17	2
1	A	138	ARG	NE-CZ-NH2	-6.64	116.98	120.30	17	4
1	A	214	TYR	CB-CG-CD2	-6.29	117.22	121.00	19	1
1	A	212	TYR	CB-CG-CD1	6.11	124.67	121.00	17	2
1	A	214	TYR	CB-CG-CD1	5.95	124.57	121.00	19	2
1	A	236	TYR	CB-CG-CD1	-5.45	117.73	121.00	9	1
1	A	70	ASP	CB-CG-OD1	5.37	123.13	118.30	2	2
1	A	129	ASP	CB-CA-C	5.32	121.04	110.40	2	1
1	A	237	ARG	NH1-CZ-NH2	-5.28	113.59	119.40	12	1
1	A	61	ASP	CB-CG-OD1	5.19	122.97	118.30	12	2
1	A	201	ASP	CB-CG-OD1	5.16	122.94	118.30	2	1
1	A	10	TYR	CB-CG-CD2	-5.03	117.98	121.00	18	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	235	ARG	Sidechain	7
1	A	237	ARG	Sidechain	6
1	A	132	LYS	Peptide	5
1	A	66	ARG	Sidechain	5
1	A	10	TYR	Sidechain	4
1	A	60	TYR	Sidechain	3
1	A	134	TYR	Sidechain	3
1	A	213	ARG	Sidechain	3
1	A	55	ARG	Sidechain	3
1	A	236	TYR	Sidechain	2
1	A	214	TYR	Sidechain	2
1	A	138	ARG	Sidechain	2
1	A	7	ARG	Sidechain	1
1	A	209	ASP	Peptide	1
1	A	71	TYR	Sidechain	1

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	711	693	693	0±1
All	All	14220	13860	13860	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:213:ARG:HH21	1:A:215:HIS:CE1	0.55	2.20	19	1
1:A:132:LYS:HZ3	1:A:201:ASP:CG	0.45	2.14	8	1
1:A:132:LYS:HZ1	1:A:201:ASP:CG	0.42	2.18	19	1
1:A:124:ASP:HB3	1:A:134:TYR:CE2	0.41	2.51	8	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	89/238 (37%)	84±2 (94±3%)	4±2 (5±2%)	1±1 (1±1%)	23	70
All	All	1780/4760 (37%)	1676 (94%)	86 (5%)	18 (1%)	23	70

All 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	217	TRP	3
1	A	231	SER	3
1	A	55	ARG	3
1	A	214	TYR	2
1	A	230	ALA	2
1	A	129	ASP	1
1	A	130	LYS	1
1	A	126	GLN	1
1	A	190	VAL	1
1	A	203	THR	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	67/202 (33%)	64±1 (95±2%)	3±1 (5±2%)	36	81
All	All	1340/4040 (33%)	1278 (95%)	62 (5%)	36	81

All 31 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	132	LYS	6
1	A	61	ASP	4
1	A	237	ARG	4
1	A	129	ASP	4
1	A	15	LEU	3
1	A	201	ASP	3
1	A	66	ARG	3
1	A	213	ARG	3
1	A	60	TYR	2
1	A	217	TRP	2
1	A	10	TYR	2
1	A	70	ASP	2
1	A	215	HIS	2
1	A	216	ASN	2
1	A	14	ASP	2
1	A	194	VAL	2
1	A	214	TYR	2
1	A	134	TYR	1
1	A	212	TYR	1
1	A	141	TYR	1
1	A	207	THR	1
1	A	235	ARG	1
1	A	55	ARG	1
1	A	206	LEU	1
1	A	127	ILE	1
1	A	124	ASP	1
1	A	128	ASN	1
1	A	130	LYS	1
1	A	209	ASP	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	190	VAL	1
1	A	203	THR	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 33% for the well-defined parts and 17% for the entire structure.

7.1 Chemical shift list 1

File name: 2maf_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	530
Number of shifts mapped to atoms	530
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	113	0.51 ± 0.12	Should be applied
$^{13}\text{C}_\beta$	93	0.42 ± 0.19	None needed (< 0.5 ppm)
$^{13}\text{C}'$	112	2.13 ± 0.12	Should be applied
^{15}N	106	0.84 ± 0.26	Should be applied

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 33%, i.e. 361 atoms were assigned a chemical shift out of a possible 1099. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	298/444 (67%)	73/177 (41%)	152/180 (84%)	73/87 (84%)
Sidechain	63/507 (12%)	0/296 (0%)	63/183 (34%)	0/28 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	0/148 (0%)	0/77 (0%)	0/68 (0%)	0/3 (0%)
Overall	361/1099 (33%)	73/550 (13%)	215/431 (50%)	73/118 (62%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 17%, i.e. 506 atoms were assigned a chemical shift out of a possible 2953. 0 out of 25 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	417/1145 (36%)	101/456 (22%)	215/466 (46%)	101/223 (45%)
Sidechain	89/1507 (6%)	0/886 (0%)	89/521 (17%)	0/100 (0%)
Aromatic	0/301 (0%)	0/158 (0%)	0/131 (0%)	0/12 (0%)
Overall	506/2953 (17%)	101/1500 (7%)	304/1118 (27%)	101/335 (30%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

