



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 01:02 pm GMT

PDB ID : 3MAD
Title : Crystal structure of StSPL (symmetric form)
Authors : Bourquin, F.; Grutter, M.G.; Capitani, G.
Deposited on : 2010-03-23
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	514	<div> <div>2%</div> <div> <div></div> <div>75%</div> <div>12%</div> <div>12%</div> </div> </div>
1	B	514	<div> <div>2%</div> <div> <div></div> <div>72%</div> <div>15%</div> <div>12%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	B	514	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7266 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sphingosine-1-phosphate lyase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	451	Total	C	N	O	P	S	0	2	0
			3450	2205	604	627	1	13			
1	B	451	Total	C	N	O	P	S	0	3	0
			3456	2208	602	632	1	13			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	EXPRESSION TAG	UNP Q67PY4
A	1	PRO	-	EXPRESSION TAG	UNP Q67PY4
A	508	HIS	-	EXPRESSION TAG	UNP Q67PY4
A	509	HIS	-	EXPRESSION TAG	UNP Q67PY4
A	510	HIS	-	EXPRESSION TAG	UNP Q67PY4
A	511	HIS	-	EXPRESSION TAG	UNP Q67PY4
A	512	HIS	-	EXPRESSION TAG	UNP Q67PY4
A	513	HIS	-	EXPRESSION TAG	UNP Q67PY4
B	0	MET	-	EXPRESSION TAG	UNP Q67PY4
B	1	PRO	-	EXPRESSION TAG	UNP Q67PY4
B	508	HIS	-	EXPRESSION TAG	UNP Q67PY4
B	509	HIS	-	EXPRESSION TAG	UNP Q67PY4
B	510	HIS	-	EXPRESSION TAG	UNP Q67PY4
B	511	HIS	-	EXPRESSION TAG	UNP Q67PY4
B	512	HIS	-	EXPRESSION TAG	UNP Q67PY4
B	513	HIS	-	EXPRESSION TAG	UNP Q67PY4

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	173	Total	O	0	0
			173	173		
3	B	177	Total	O	0	0
			177	177		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Sphingosine-1-phosphate lyase



• Molecule 1: Sphingosine-1-phosphate lyase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.63Å 126.76Å 137.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.45 – 2.00 29.45 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.7 (29.45-2.00) 98.8 (29.45-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.60 (at 2.00Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.4_29)	Depositor
R, R_{free}	0.210 , 0.247 0.202 , 0.241	Depositor DCC
R_{free} test set	949 reflections (1.40%)	DCC
Wilson B-factor (Å ²)	25.3	Xtrriage
Anisotropy	0.307	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7266	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 42.24 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.1006e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/3526	0.48	0/4816
1	B	0.28	0/3528	0.47	0/4821
All	All	0.29	0/7054	0.48	0/9637

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3450	0	3378	41	0
1	B	3456	0	3378	51	0
2	A	5	0	0	1	0
2	B	5	0	0	0	0
3	A	173	0	0	2	0
3	B	177	0	0	1	0
All	All	7266	0	6756	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:168:GLY:HA3	1:B:311:LLP:H5'2	1.59	0.84
1:A:372:GLU:HG2	1:B:73:ALA:HA	1.70	0.74
1:A:311:LLP:H4'1	3:A:580:HOH:O	1.88	0.72
1:B:494:ARG:HH11	1:B:495:GLN:HG3	1.55	0.71
1:A:499:GLY:O	1:A:503:LEU:HD22	1.90	0.71
1:B:494:ARG:NH1	1:B:495:GLN:HG3	2.07	0.69
1:A:76:PRO:HB2	1:A:79:GLU:OE2	1.94	0.68
1:B:170:GLU:HG3	1:B:350:PHE:HD2	1.67	0.59
1:B:441:HIS:C	1:B:441:HIS:CD2	2.76	0.59
1:B:60:TYR:HB3	1:B:64:PHE:HD2	1.68	0.57
1:B:311:LLP:OP4	1:B:311:LLP:H4'1	2.06	0.56
1:B:311:LLP:OP4	1:B:311:LLP:C4'	2.55	0.55
1:A:441:HIS:C	1:A:441:HIS:CD2	2.80	0.54
1:A:294:PRO:HB2	1:A:299:LEU:HD21	1.88	0.54
1:A:256:PRO:O	1:A:259:GLU:HG2	2.08	0.54
1:A:196:VAL:HG12	1:A:243:ALA:HB3	1.90	0.52
1:B:206:LYS:HG2	1:B:210:TYR:CZ	2.43	0.52
1:B:419:GLN:OE1	1:B:475:ALA:HA	2.09	0.52
1:A:225:ARG:HD2	1:A:254:VAL:O	2.09	0.52
1:A:311:LLP:HE3	2:A:514:PO4:O4	2.10	0.51
1:A:299:LEU:O	1:A:325:ARG:NH2	2.43	0.51
1:A:91:GLU:OE2	1:B:134:PRO:HD2	2.11	0.51
1:A:72:ARG:O	1:B:372:GLU:HG3	2.11	0.50
1:A:153:GLY:HA2	1:A:160:VAL:HG21	1.93	0.50
1:B:379:ARG:NH2	3:B:529:HOH:O	2.45	0.50
1:A:196:VAL:O	1:A:217:ARG:HA	2.11	0.50
1:A:63[A]:ARG:HG2	1:A:64:PHE:CE2	2.47	0.49
1:A:63[A]:ARG:HG2	1:A:64:PHE:CD2	2.47	0.49
1:A:210:TYR:CE2	1:B:350:PHE:CD1	3.01	0.48
1:B:503:LEU:HD22	1:B:503:LEU:O	2.13	0.48
1:B:170:GLU:OE2	1:B:349:THR:HB	2.13	0.48
1:B:180:ARG:HG3	1:B:192:PRO:HG3	1.95	0.48
1:A:192:PRO:HD2	3:A:537:HOH:O	2.13	0.48
1:A:299:LEU:HB2	1:A:302:VAL:HG23	1.95	0.48
1:A:340:TRP:CD1	1:A:341:PRO:HD2	2.49	0.47
1:B:196:VAL:O	1:B:217:ARG:HA	2.14	0.47
1:B:491:GLU:HB2	1:B:494:ARG:HH22	1.80	0.47
1:B:169:THR:O	1:B:173:LEU:HG	2.15	0.46
1:A:317:LYS:HD3	1:A:317:LYS:HA	1.64	0.46
1:A:431:LEU:HD13	1:A:440:PHE:CD1	2.51	0.46
1:A:136:THR:O	1:A:140:GLU:HG3	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:326:ARG:NH1	1:A:328:ASP:OD2	2.49	0.46
1:B:387:ARG:HD3	1:B:454:VAL:CG1	2.46	0.46
1:B:389:LYS:O	1:B:393:ARG:HG3	2.16	0.45
1:A:134:PRO:HD2	1:B:91:GLU:OE2	2.15	0.45
1:B:92:SER:N	1:B:93:PRO:CD	2.79	0.45
1:A:434:LEU:HD11	1:A:441:HIS:HB3	1.99	0.45
1:A:83:GLU:O	1:A:87:MET:HG3	2.16	0.45
1:B:170:GLU:HG3	1:B:350:PHE:CD2	2.49	0.45
1:A:493:VAL:HG12	1:A:497:LEU:HD22	1.98	0.44
1:B:387:ARG:HD3	1:B:454:VAL:HG11	1.99	0.44
1:B:276:CYS:O	1:B:308:ASP:HB2	2.17	0.44
1:A:276:CYS:O	1:A:308:ASP:HB2	2.17	0.44
1:B:396:PRO:O	1:B:397:SER:HB2	2.17	0.44
1:A:281:ILE:HD11	1:A:312:TYR:CD2	2.52	0.44
1:A:311:LLP:NZ	1:A:311:LLP:O3	2.48	0.43
1:B:153:GLY:HA2	1:B:160:VAL:HG21	2.00	0.43
1:B:292:VAL:HA	1:B:293:PRO:HD3	1.71	0.43
1:A:341:PRO:HA	1:B:435:HIS:O	2.19	0.43
1:B:369:LEU:HD13	1:B:374:TYR:CE2	2.54	0.43
1:A:397:SER:CB	1:A:469:ARG:HH22	2.32	0.42
1:B:441:HIS:C	1:B:441:HIS:HD2	2.19	0.42
1:A:102:GLY:O	1:A:103:ALA:HB3	2.19	0.42
1:A:156:ALA:HB3	1:A:160:VAL:HG23	1.99	0.42
1:A:446:LEU:HA	1:A:446:LEU:HD12	1.92	0.42
1:B:173:LEU:HB2	1:B:350:PHE:HE2	1.85	0.42
1:B:493:VAL:O	1:B:497:LEU:HD23	2.19	0.42
1:B:274:ASP:OD1	1:B:311:LLP:N1	2.53	0.42
1:B:83:GLU:O	1:B:87:MET:HG3	2.20	0.42
1:B:340:TRP:CD1	1:B:341:PRO:HD2	2.55	0.41
1:B:92:SER:HA	1:B:95:TRP:NE1	2.35	0.41
1:B:388:LEU:HD13	1:B:444:LEU:HD21	2.01	0.41
1:B:492:LEU:C	1:B:492:LEU:HD13	2.40	0.41
1:B:325:ARG:HG2	1:B:326:ARG:HG3	2.01	0.41
1:B:376:ASP:OD1	1:B:379:ARG:NH1	2.52	0.41
1:B:236:THR:HB	1:B:237:PRO:CD	2.50	0.41
1:B:151:LEU:O	1:B:296:ASP:HB2	2.20	0.41
1:B:444:LEU:HA	1:B:448:HIS:ND1	2.35	0.41
1:B:195:VAL:HB	1:B:242:VAL:HG12	2.03	0.41
1:B:299:LEU:O	1:B:325:ARG:NH2	2.53	0.41
1:A:441:HIS:C	1:A:441:HIS:HD2	2.24	0.41
1:B:395:ILE:HA	1:B:396:PRO:HD3	1.96	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:LEU:HD23	1:A:207:ALA:HB2	2.03	0.41
1:A:169:THR:O	1:A:173:LEU:HG	2.21	0.41
1:A:131:ASP:HB2	1:B:430:ARG:HH21	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	450/514 (88%)	441 (98%)	9 (2%)	0	100	100
1	B	451/514 (88%)	436 (97%)	15 (3%)	0	100	100
All	All	901/1028 (88%)	877 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	339/395 (86%)	325 (96%)	14 (4%)	35	31
1	B	340/395 (86%)	325 (96%)	15 (4%)	33	28
All	All	679/790 (86%)	650 (96%)	29 (4%)	33	29

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	LYS
1	A	61	ARG
1	A	128	LEU
1	A	263	LEU
1	A	282	LEU
1	A	310	HIS
1	A	330	LEU
1	A	410	VAL
1	A	441	HIS
1	A	446	LEU
1	A	461	LEU
1	A	491	GLU
1	A	497	LEU
1	A	503	LEU
1	B	70	LEU
1	B	128	LEU
1	B	233	GLU
1	B	253	VAL
1	B	263	LEU
1	B	282	LEU
1	B	292	VAL
1	B	310	HIS
1	B	317	LYS
1	B	328	ASP
1	B	330	LEU
1	B	410	VAL
1	B	441	HIS
1	B	446	LEU
1	B	461	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	441	HIS
1	B	441	HIS
1	B	495	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	A	311	1	24,24,25	1.80	6 (25%)	28,32,34	1.51	5 (17%)
1	LLP	B	311	1	24,24,25	1.75	4 (16%)	28,32,34	1.47	5 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	311	1	-	0/15/17/19	0/1/1/1
1	LLP	B	311	1	-	0/15/17/19	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	311	LLP	O3-C3	-5.82	1.23	1.37
1	A	311	LLP	O3-C3	-5.79	1.23	1.37
1	A	311	LLP	C6-N1	2.00	1.38	1.34
1	A	311	LLP	C2-N1	2.12	1.38	1.33
1	B	311	LLP	C4'-NZ	2.25	1.34	1.27
1	A	311	LLP	CA-C	2.29	1.53	1.50
1	B	311	LLP	C4-C4'	2.37	1.50	1.46
1	B	311	LLP	C2-N1	2.42	1.38	1.33
1	A	311	LLP	C4'-NZ	2.48	1.34	1.27
1	A	311	LLP	C4-C4'	2.82	1.51	1.46

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	311	LLP	C4-C4'-NZ	-3.27	108.77	124.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	311	LLP	C4-C4'-NZ	-3.10	109.59	124.66
1	B	311	LLP	C5-C6-N1	-2.77	119.19	123.87
1	B	311	LLP	OP4-P-OP1	-2.71	98.87	106.47
1	A	311	LLP	OP4-P-OP1	-2.60	99.18	106.47
1	A	311	LLP	C5-C6-N1	-2.40	119.80	123.87
1	B	311	LLP	CE-NZ-C4'	-2.15	112.79	119.03
1	A	311	LLP	O-C-CA	-2.04	119.40	125.02
1	B	311	LLP	OP4-C5'-C5	2.96	115.27	109.32
1	A	311	LLP	OP4-C5'-C5	4.77	118.92	109.32

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	311	LLP	3	0
1	B	311	LLP	4	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	514	-	4,4,4	0.79	0	6,6,6	0.50	0
2	PO4	B	514	-	4,4,4	0.75	0	6,6,6	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	514	-	-	0/0/0/0	0/0/0/0
2	PO4	B	514	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	514	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	450/514 (87%)	-0.01	11 (2%) 59 59	17, 26, 43, 76	0
1	B	450/514 (87%)	0.01	8 (1%) 69 68	17, 27, 43, 78	0
All	All	900/1028 (87%)	-0.00	19 (2%) 64 63	17, 26, 43, 78	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	508	HIS	3.9
1	A	61	ARG	3.5
1	B	60	TYR	3.4
1	B	61	ARG	3.2
1	A	494	ARG	3.0
1	B	503	LEU	3.0
1	A	62	ASP	3.0
1	A	249	TYR	2.7
1	B	72	ARG	2.6
1	B	62	ASP	2.5
1	A	60	TYR	2.4
1	B	275	ALA	2.4
1	A	63[A]	ARG	2.4
1	A	504	LEU	2.3
1	B	350	PHE	2.3
1	A	72	ARG	2.2
1	A	350	PHE	2.2
1	A	276	CYS	2.1
1	B	494	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	LLP	A	311	24/25	0.92	0.15	-	19,28,36,38	0
1	LLP	B	311	24/25	0.94	0.16	-	21,28,39,41	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	PO4	B	514	5/5	0.92	0.21	3.23	26,30,36,41	5
2	PO4	A	514	5/5	0.94	0.14	0.31	26,29,43,45	5

6.5 Other polymers [i](#)

There are no such residues in this entry.