



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 30, 2017 – 06:20 PM EDT

PDB ID : 3MDL
Title : X-ray crystal structure of 1-arachidonoyl glycerol bound to the cyclooxygenase channel of cyclooxygenase-2
Authors : Vecchio, A.J.; Malkowski, M.G.
Deposited on : unknown
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	587	<div> <div>3%</div> <div>86%</div> <div>8%</div> <div>6%</div> </div>
1	B	587	<div> <div>4%</div> <div>87%</div> <div>7%</div> <div>6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	1AG	A	1	-	-	-	X
2	1AG	B	1	-	-	-	X
3	AKR	A	2	-	-	X	-
3	AKR	B	3	-	-	X	-
4	GOL	B	6	-	-	-	X
4	GOL	B	7	-	-	-	X
5	COH	A	614	-	-	-	X
6	NAG	A	661	X	-	-	X
6	NAG	A	681	-	-	-	X
6	NAG	B	661	-	-	-	X
6	NAG	B	681	-	-	-	X
7	MAN	A	673	X	-	-	-
8	BOG	A	703	-	-	-	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 10254 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Prostaglandin G/H synthase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	552	Total	C	N	O	S	0	4	0
			4485	2897	753	810	25			
1	B	551	Total	C	N	O	S	0	2	0
			4455	2877	747	806	25			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	28	ASN	-	EXPRESSION TAG	UNP Q05769
A	29	HIS	-	EXPRESSION TAG	UNP Q05769
A	30	HIS	-	EXPRESSION TAG	UNP Q05769
A	31	HIS	-	EXPRESSION TAG	UNP Q05769
A	32	HIS	-	EXPRESSION TAG	UNP Q05769
A	33	HIS	-	EXPRESSION TAG	UNP Q05769
A	34	HIS	-	EXPRESSION TAG	UNP Q05769
A	594	ALA	ASN	ENGINEERED	UNP Q05769
B	28	ASN	-	EXPRESSION TAG	UNP Q05769
B	29	HIS	-	EXPRESSION TAG	UNP Q05769
B	30	HIS	-	EXPRESSION TAG	UNP Q05769
B	31	HIS	-	EXPRESSION TAG	UNP Q05769
B	32	HIS	-	EXPRESSION TAG	UNP Q05769
B	33	HIS	-	EXPRESSION TAG	UNP Q05769
B	34	HIS	-	EXPRESSION TAG	UNP Q05769
B	594	ALA	ASN	ENGINEERED	UNP Q05769

- Molecule 2 is (2S)-2,3-dihydroxypropyl (5Z,8Z,11Z,14Z)-icosa-5,8,11,14-tetraenoate (three-letter code: 1AG) (formula: C₂₃H₃₈O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			27	23	4		
2	B	1	Total	C	O	0	0
			27	23	4		

- Molecule 3 is ACRYLIC ACID (three-letter code: AKR) (formula: C₃H₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			5	3	2		
3	A	1	Total	C	O	0	0
			5	3	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			5	3	2		
3	B	1	Total	C	O	0	0
			5	3	2		
3	B	1	Total	C	O	0	0
			5	3	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING CO (three-letter code: COH) (formula: $C_{34}H_{32}CoN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Co	N	O	0	0
			43	34	1	4	4		
5	B	1	Total	C	Co	N	O	0	0
			43	34	1	4	4		

- Molecule 6 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 8 is B-OCTYLGLUCOSIDE (three-letter code: BOG) (formula: C₁₄H₂₈O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			20	14	6		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	476	Total	O	0	0
			476	476		
9	B	484	Total	O	0	0
			484	484		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Prostaglandin G/H synthase 2



• Molecule 1: Prostaglandin G/H synthase 2



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	118.96Å 131.77Å 179.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.97 – 2.20 19.97 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.2 (19.97-2.20) 99.2 (19.97-2.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.19Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, R_{free}	0.165 , 0.215 0.193 , 0.238	Depositor DCC
R_{free} test set	3597 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	28.7	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 37.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10254	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAG, AKR, 1AG, COH, BOG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/4624	0.53	0/6272
1	B	0.42	0/4588	0.53	1/6227 (0.0%)
All	All	0.42	0/9212	0.53	1/12499 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	531	LEU	CA-CB-CG	6.35	129.90	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4485	0	4363	40	0
1	B	4455	0	4306	40	0
2	A	27	0	36	12	0
2	B	27	0	36	18	0
3	A	10	0	6	2	0
3	B	15	0	9	2	0
4	A	6	0	8	0	0
4	B	12	0	16	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	43	0	30	2	0
5	B	43	0	30	1	0
6	A	70	0	62	5	0
6	B	70	0	63	2	0
7	A	11	0	10	1	0
8	A	20	0	28	0	0
9	A	476	0	0	0	0
9	B	484	0	0	3	0
All	All	10254	0	9003	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 86 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:205:PHE:HE2	2:B:1:1AG:H15	1.02	1.10
1:B:205:PHE:CE2	2:B:1:1AG:H15	1.93	1.02
1:B:477:SER:HB2	3:B:3:AKR:HA1	1.50	0.94
1:A:216:ARG:HH11	6:A:672:NAG:H83	1.38	0.88
1:A:216:ARG:NH1	6:A:672:NAG:H83	1.90	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	554/587 (94%)	543 (98%)	11 (2%)	0	100	100
1	B	551/587 (94%)	541 (98%)	10 (2%)	0	100	100
All	All	1105/1174 (94%)	1084 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	489/525 (93%)	484 (99%)	5 (1%)	80	89
1	B	482/525 (92%)	476 (99%)	6 (1%)	75	86
All	All	971/1050 (92%)	960 (99%)	11 (1%)	78	88

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	556	PHE
1	B	108	LEU
1	B	385	TYR
1	A	484	GLU
1	B	289	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	386	HIS
1	B	374	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	1AG	A	1	-	26,26,26	2.25	4 (15%)	27,27,27	1.00	0
3	AKR	A	2	-	1,4,4	1.59	0	0,4,4	0.00	-
3	AKR	A	3	-	1,4,4	1.57	0	0,4,4	0.00	-
4	GOL	A	4	-	5,5,5	0.30	0	5,5,5	0.36	0
5	COH	A	614	1	28,50,50	4.47	19 (67%)	21,82,82	4.69	8 (38%)
6	NAG	A	661	1,6	14,14,15	0.54	0	15,19,21	1.50	1 (6%)
6	NAG	A	662	6	14,14,15	0.50	0	15,19,21	0.59	0
6	NAG	A	671	1,6	14,14,15	0.50	0	15,19,21	1.25	1 (6%)
6	NAG	A	672	7,6	14,14,15	0.51	0	15,19,21	1.20	1 (6%)
7	MAN	A	673	6	11,11,12	0.56	0	13,15,17	2.15	3 (23%)
6	NAG	A	681	1	14,14,15	0.53	0	15,19,21	0.73	0
8	BOG	A	703	-	20,20,20	0.45	0	25,25,25	0.54	0
2	1AG	B	1	-	26,26,26	2.30	4 (15%)	27,27,27	0.85	1 (3%)
3	AKR	B	3	-	1,4,4	1.56	0	0,4,4	0.00	-
3	AKR	B	4	-	1,4,4	1.58	0	0,4,4	0.00	-
3	AKR	B	5	-	1,4,4	1.55	0	0,4,4	0.00	-
4	GOL	B	6	-	5,5,5	0.34	0	5,5,5	0.17	0
5	COH	B	615	1	28,50,50	4.46	20 (71%)	21,82,82	4.84	9 (42%)
6	NAG	B	661	1,6	14,14,15	0.62	0	15,19,21	0.79	0
6	NAG	B	662	6	14,14,15	0.55	0	15,19,21	2.28	4 (26%)
6	NAG	B	671	1,6	14,14,15	0.45	0	15,19,21	1.13	1 (6%)
6	NAG	B	672	6	14,14,15	0.54	0	15,19,21	0.98	1 (6%)
6	NAG	B	681	1	14,14,15	0.56	0	15,19,21	0.86	1 (6%)
4	GOL	B	7	-	5,5,5	0.36	0	5,5,5	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1AG	A	1	-	-	0/26/26/26	0/0/0/0
3	AKR	A	2	-	-	0/0/2/2	0/0/0/0
3	AKR	A	3	-	-	0/0/2/2	0/0/0/0
4	GOL	A	4	-	-	0/4/4/4	0/0/0/0
5	COH	A	614	1	-	0/8/94/94	0/0/8/8
6	NAG	A	661	1,6	1/1/5/7	0/6/23/26	0/1/1/1
6	NAG	A	662	6	-	0/6/23/26	0/1/1/1
6	NAG	A	671	1,6	-	0/6/23/26	0/1/1/1
6	NAG	A	672	7,6	-	0/6/23/26	0/1/1/1
7	MAN	A	673	6	1/1/4/5	0/2/19/22	0/1/1/1
6	NAG	A	681	1	-	0/6/23/26	0/1/1/1
8	BOG	A	703	-	-	0/11/31/31	0/1/1/1
2	1AG	B	1	-	-	0/26/26/26	0/0/0/0
3	AKR	B	3	-	-	0/0/2/2	0/0/0/0
3	AKR	B	4	-	-	0/0/2/2	0/0/0/0
3	AKR	B	5	-	-	0/0/2/2	0/0/0/0
4	GOL	B	6	-	-	0/4/4/4	0/0/0/0
5	COH	B	615	1	-	0/8/94/94	0/0/8/8
6	NAG	B	661	1,6	-	0/6/23/26	0/1/1/1
6	NAG	B	662	6	-	0/6/23/26	0/1/1/1
6	NAG	B	671	1,6	-	0/6/23/26	0/1/1/1
6	NAG	B	672	6	-	0/6/23/26	0/1/1/1
6	NAG	B	681	1	-	0/6/23/26	0/1/1/1
4	GOL	B	7	-	-	0/4/4/4	0/0/0/0

The worst 5 of 47 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	1AG	O01-C21	-8.69	1.25	1.45
2	A	1	1AG	O01-C21	-8.41	1.26	1.45
5	B	615	COH	CHA-C1A	-7.90	1.38	1.53
5	A	614	COH	CHA-C1A	-7.77	1.38	1.53
5	B	615	COH	CHB-C4A	-7.76	1.38	1.53

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	673	MAN	C1-C2-C3	-3.96	104.63	109.65
5	B	615	COH	CBA-CAA-C2A	-3.92	107.13	114.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	614	COH	CBD-CAD-C3D	-2.98	106.78	112.48
5	B	615	COH	CBD-CAD-C3D	-2.96	106.82	112.48
5	A	614	COH	C3C-C4C-NC	-2.42	107.17	109.03

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
7	A	673	MAN	C1
6	A	661	NAG	C5

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 44 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	1AG	12	0
3	A	2	AKR	2	0
5	A	614	COH	2	0
6	A	672	NAG	5	0
7	A	673	MAN	1	0
2	B	1	1AG	18	0
3	B	3	AKR	2	0
5	B	615	COH	1	0
6	B	672	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	552/587 (94%)	-0.13	18 (3%) 47 44	8, 23, 41, 48	1 (0%)
1	B	551/587 (93%)	0.06	25 (4%) 34 32	6, 27, 42, 51	1 (0%)
All	All	1103/1174 (93%)	-0.04	43 (3%) 40 38	6, 25, 42, 51	2 (0%)

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	74	PHE	4.6
1	B	82	LEU	4.6
1	B	399	ASP	4.0
1	B	78	ILE	3.9
1	A	115	TYR	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	NAG	A	681	14/15	0.90	0.34	3.78	39,43,45,46	0
6	NAG	B	681	14/15	0.85	0.29	3.65	41,45,47,48	0
6	NAG	A	661	14/15	0.87	0.22	3.44	40,46,49,53	0
6	NAG	B	661	14/15	0.84	0.31	3.40	43,51,53,58	0
4	GOL	B	6	6/6	0.96	0.24	3.34	32,33,34,34	0
8	BOG	A	703	20/20	0.83	0.21	2.99	45,46,47,47	0
2	1AG	A	1	27/27	0.72	0.28	2.65	36,49,58,58	0
5	COH	A	614	43/43	0.86	0.20	2.19	43,48,57,60	0
2	1AG	B	1	27/27	0.79	0.25	2.08	26,46,56,57	0
4	GOL	B	7	6/6	0.76	0.28	2.06	77,78,78,78	0
4	GOL	A	4	6/6	0.94	0.16	1.89	27,29,31,31	0
3	AKR	A	3	5/5	0.86	0.24	1.77	39,40,40,40	0
3	AKR	A	2	5/5	0.74	0.22	1.15	51,51,51,52	0
5	COH	B	615	43/43	0.86	0.19	1.14	40,44,54,58	0
3	AKR	B	4	5/5	0.83	0.27	0.98	53,54,54,54	0
6	NAG	B	671	14/15	0.97	0.09	0.04	18,24,27,33	0
6	NAG	A	671	14/15	0.95	0.11	0.03	21,26,30,33	0
6	NAG	B	672	14/15	0.78	0.29	-	39,43,44,45	0
3	AKR	B	5	5/5	0.54	0.33	-	102,102,102,102	0
7	MAN	A	673	11/12	0.81	0.41	-	53,55,57,57	0
3	AKR	B	3	5/5	0.80	0.21	-	53,54,54,54	0
6	NAG	A	672	14/15	0.81	0.17	-	38,40,45,49	0
6	NAG	A	662	14/15	0.82	0.40	-	57,60,61,61	0
6	NAG	B	662	14/15	0.75	0.45	-	62,65,67,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.