



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 30, 2017 – 09:05 PM EDT

PDB ID : 3ME8  
Title : Crystal structure of putative electron transfer protein aq\_2194 from Aquifex  
aeolicus VF5  
Authors : Malashkevich, V.N.; Toro, R.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New  
York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : unknown  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2279 (1.50-1.50)
Clashscore	112137	2503 (1.50-1.50)
Ramachandran outliers	110173	2445 (1.50-1.50)
Sidechain outliers	110143	2443 (1.50-1.50)
RSRZ outliers	101464	2305 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	<div> <div>8%</div> <div> <div></div> <div>82%</div> <div>16%</div> <div>..</div> </div> </div>
1	B	170	<div> <div>10%</div> <div> <div></div> <div>79%</div> <div>11%</div> <div>..</div> <div>8%</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2969 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	169	Total	C	N	O	S	0	4	0
			1391	917	229	242	3			
1	B	156	Total	C	N	O	S	0	3	0
			1273	838	203	229	3			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP O67938
A	2	SER	-	EXPRESSION TAG	UNP O67938
A	164	GLY	-	EXPRESSION TAG	UNP O67938
A	165	HIS	-	EXPRESSION TAG	UNP O67938
A	166	HIS	-	EXPRESSION TAG	UNP O67938
A	167	HIS	-	EXPRESSION TAG	UNP O67938
A	168	HIS	-	EXPRESSION TAG	UNP O67938
A	169	HIS	-	EXPRESSION TAG	UNP O67938
A	170	HIS	-	EXPRESSION TAG	UNP O67938
B	1	MET	-	EXPRESSION TAG	UNP O67938
B	2	SER	-	EXPRESSION TAG	UNP O67938
B	164	GLY	-	EXPRESSION TAG	UNP O67938
B	165	HIS	-	EXPRESSION TAG	UNP O67938
B	166	HIS	-	EXPRESSION TAG	UNP O67938
B	167	HIS	-	EXPRESSION TAG	UNP O67938
B	168	HIS	-	EXPRESSION TAG	UNP O67938
B	169	HIS	-	EXPRESSION TAG	UNP O67938
B	170	HIS	-	EXPRESSION TAG	UNP O67938

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	164	Total	O	0	0
			164	164		

*Continued on next page...*


*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	141	Total	O	0	0
			141	141		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative uncharacterized protein

Chain A: 



- Molecule 1: Putative uncharacterized protein

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.57Å 134.77Å 37.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.50 19.49 – 1.50	Depositor EDS
% Data completeness (in resolution range)	95.2 (20.00-1.50) 95.2 (19.49-1.50)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.10 (at 1.50Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.188 , 0.220 0.193 , 0.230	Depositor DCC
$R_{free}$ test set	2718 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.9	Xtriage
Anisotropy	0.213	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 39.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2969	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.62	0/1443	0.69	0/1953
1	B	0.65	0/1312	0.68	0/1776
All	All	0.64	0/2755	0.68	0/3729

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1391	0	1413	29	0
1	B	1273	0	1307	27	0
2	A	164	0	0	4	0
2	B	141	0	0	2	0
All	All	2969	0	2720	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:ASP:OD1	1:A:93:LYS:HG2	1.54	1.08

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:118:MET:HB2	1:B:93:LYS:HZ1	1.33	0.94
1:B:118:MET:SD	1:B:126:HIS:HE1	2.01	0.82
1:B:37:TYR:H	1:B:125:ILE:HD12	1.43	0.81
1:B:40:CYS:SG	1:B:125:ILE:HG22	2.21	0.80
1:A:118:MET:HB2	1:B:93:LYS:NZ	2.05	0.71
1:A:70[B]:PHE:HZ	1:A:106:LEU:CD1	2.09	0.66
1:A:118:MET:CB	1:B:93:LYS:HZ1	2.09	0.62
1:A:91:ASP:OD1	1:A:93:LYS:CG	2.42	0.60
1:A:2:SER:N	1:A:142:TYR:HH	2.00	0.59
1:B:118:MET:SD	1:B:126:HIS:CE1	2.92	0.58
1:B:40:CYS:HB2	1:B:125:ILE:HG22	1.85	0.57
1:B:21:PHE:HB2	1:B:25[B]:ASN:OD1	2.04	0.56
1:B:125:ILE:HG13	2:B:257:HOH:O	2.05	0.56
1:A:118:MET:CB	1:B:93:LYS:NZ	2.67	0.56
1:B:40:CYS:CB	1:B:125:ILE:HG22	2.35	0.55
1:B:126:HIS:ND1	1:B:127:PRO:HD3	2.21	0.54
1:A:78:LEU:HD21	2:A:262:HOH:O	2.07	0.54
1:A:14:VAL:HG22	1:A:20:GLU:HG2	1.88	0.54
1:A:82:LYS:HE3	1:A:86:LYS:HE3	1.90	0.54
1:A:75[B]:LYS:HD3	2:A:242:HOH:O	2.08	0.54
1:B:40:CYS:SG	1:B:125:ILE:CG2	2.94	0.53
1:A:118:MET:CG	1:B:93:LYS:HZ2	2.22	0.53
1:A:70[B]:PHE:HZ	1:A:106:LEU:HD12	1.74	0.53
1:A:70[B]:PHE:CZ	1:A:106:LEU:CD1	2.93	0.51
1:A:70[B]:PHE:HZ	1:A:106:LEU:HD13	1.78	0.48
1:A:149:ASN:O	1:A:152:GLU:HG2	2.15	0.47
1:B:45:PRO:HD2	1:B:125:ILE:HG23	1.96	0.47
1:A:168:HIS:CD2	1:A:170:HIS:H	2.33	0.47
1:B:37:TYR:HB2	1:B:125:ILE:HB	1.98	0.46
1:A:148:TYR:HB3	1:A:153:PHE:HE1	1.80	0.46
1:A:75[A]:LYS:HG2	2:A:339:HOH:O	2.15	0.45
1:A:119:THR:HG23	2:A:202:HOH:O	2.16	0.44
1:B:40:CYS:SG	1:B:44:CYS:CB	3.05	0.44
1:A:70[B]:PHE:CZ	1:A:106:LEU:HD13	2.52	0.44
1:B:25[B]:ASN:ND2	2:B:366:HOH:O	2.49	0.43
1:A:165:HIS:CE1	1:A:166:HIS:CE1	3.06	0.43
1:A:148:TYR:HB3	1:A:153:PHE:CE1	2.53	0.42
1:B:87:GLU:HG2	1:B:88:TYR:CD1	2.54	0.42
1:A:36:ILE:O	1:A:71:THR:HA	2.19	0.42
1:A:70[B]:PHE:CZ	1:A:109:LEU:HD23	2.55	0.42
1:A:33[A]:LEU:HD11	1:A:133:LEU:HD11	2.02	0.42

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:118:MET:CG	1:B:93:LYS:NZ	2.82	0.42
1:B:115:PHE:HZ	1:B:129:VAL:CG1	2.33	0.41
1:B:40:CYS:CB	1:B:44:CYS:SG	3.07	0.41
1:B:70:PHE:CZ	1:B:109:LEU:HD23	2.55	0.41
1:B:14:VAL:HG22	1:B:20[B]:GLU:HG2	2.02	0.41
1:B:115:PHE:CZ	1:B:129:VAL:CG1	3.04	0.41
1:B:40:CYS:HB2	1:B:125:ILE:CG2	2.51	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	171/170 (101%)	165 (96%)	6 (4%)	0	100	100
1	B	155/170 (91%)	150 (97%)	4 (3%)	1 (1%)	28	7
All	All	326/340 (96%)	315 (97%)	10 (3%)	1 (0%)	44	19

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	126	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	153/150 (102%)	148 (97%)	5 (3%)	43	11
1	B	142/150 (95%)	136 (96%)	6 (4%)	34	6
All	All	295/300 (98%)	284 (96%)	11 (4%)	45	9

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	75[A]	LYS
1	A	75[B]	LYS
1	A	91	ASP
1	A	122	ASN
1	A	147	ASN
1	B	25[A]	ASN
1	B	25[B]	ASN
1	B	87	GLU
1	B	126	HIS
1	B	128	ASN
1	B	151	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	147	ASN
1	A	165	HIS
1	A	168	HIS
1	B	126	HIS
1	B	128	ASN
1	B	147	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	169/170 (99%)	0.39	14 (8%) <span>12</span> <span>14</span>	10, 15, 25, 32	0
1	B	156/170 (91%)	0.78	17 (10%) <span>6</span> <span>7</span>	12, 17, 37, 42	0
All	All	325/340 (95%)	0.58	31 (9%) <span>9</span> <span>10</span>	10, 16, 28, 42	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	42	ALA	11.2
1	B	41	ARG	10.3
1	B	40	CYS	10.1
1	B	43	ALA	7.9
1	B	125	ILE	5.9
1	B	44	CYS	5.1
1	A	121	GLY	4.8
1	B	39	HIS	4.6
1	B	126	HIS	4.3
1	A	122	ASN	4.2
1	A	168	HIS	3.9
1	A	170	HIS	3.6
1	A	169	HIS	3.5
1	A	70[A]	PHE	3.2
1	B	45	PRO	3.2
1	A	120	ALA	3.0
1	A	33[A]	LEU	2.8
1	B	46	LEU	2.7
1	B	163	GLU	2.7
1	B	118	MET	2.7
1	B	83	ARG	2.5
1	B	37	TYR	2.3
1	B	116	ARG	2.2
1	A	78	LEU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	165	HIS	2.2
1	B	79	GLU	2.1
1	A	167	HIS	2.1
1	A	147	ASN	2.1
1	A	32	ILE	2.1
1	A	93	LYS	2.1
1	B	6	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.