



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 15, 2017 – 12:55 am GMT

PDB ID : 1MG0  
Title : Horse Liver Alcohol Dehydrogenase Complexed With NAD<sup>+</sup> and 2,3-Difluorobenzyl Alcohol  
Authors : Rubach, J.K.; Plapp, B.V.  
Deposited on : 2002-08-14  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*





The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	 97%
1	B	374	 95% 5%
1	C	374	 96%
1	D	374	 95% 5%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12301 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alcohol Dehydrogenase E chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	374	Total	C	N	O	S	0	0	0
			2784	1769	472	520	23			
1	B	374	Total	C	N	O	S	0	0	0
			2784	1769	472	520	23			
1	C	374	Total	C	N	O	S	0	0	0
			2784	1769	472	520	23			
1	D	374	Total	C	N	O	S	0	0	0
			2784	1769	472	520	23			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Zn	0	0
			2	2		
2	A	2	Total	Zn	0	0
			2	2		
2	D	2	Total	Zn	0	0
			2	2		
2	C	2	Total	Zn	0	0
			2	2		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is 2,3-DIFLUOROBENZYL ALCOHOL (three-letter code: DFB) (formula: C<sub>7</sub>H<sub>6</sub>F<sub>2</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	F	O	0	1
			19	14	4	1		
4	B	1	Total	C	F	O	0	1
			19	14	4	1		
4	C	1	Total	C	F	O	0	1
			19	14	4	1		
4	D	1	Total	C	F	O	0	1
			19	14	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	203	Total	O	0	0
			203	203		
5	B	216	Total	O	0	0
			216	216		
5	C	223	Total	O	0	0
			223	223		
5	D	263	Total	O	0	0
			263	263		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alcohol Dehydrogenase E chain

Chain A:  97% .



- Molecule 1: Alcohol Dehydrogenase E chain

Chain B:  95% 5% .



- Molecule 1: Alcohol Dehydrogenase E chain

Chain C:  96% .



- Molecule 1: Alcohol Dehydrogenase E chain

Chain D:  95% 5% .



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.11Å 180.31Å 86.91Å 90.00° 105.99° 90.00°	Depositor
Resolution (Å)	20.00 – 1.80 19.84 – 1.80	Depositor EDS
% Data completeness (in resolution range)	88.0 (20.00-1.80) 88.0 (19.84-1.80)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.16 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.1.27	Depositor
R, $R_{free}$	0.179 , 0.214 0.184 , 0.216	Depositor DCC
$R_{free}$ test set	1822 reflections (1.54%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.7	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 48.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	12301	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 63.03 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.0293e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NAD, DFB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.56	0/2836	0.70	0/3834
1	B	0.57	0/2836	0.71	3/3834 (0.1%)
1	C	0.59	0/2836	0.72	2/3834 (0.1%)
1	D	0.59	0/2836	0.71	2/3834 (0.1%)
All	All	0.58	0/11344	0.71	7/15336 (0.0%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	263	ASP	CB-CG-OD2	6.17	123.85	118.30
1	B	223	ASP	CB-CG-OD2	5.62	123.36	118.30
1	B	115	ASP	CB-CG-OD2	5.47	123.22	118.30
1	C	245	ASP	CB-CG-OD2	5.24	123.01	118.30
1	D	50	ASP	CB-CG-OD2	5.22	123.00	118.30
1	D	343	ASP	CB-CG-OD2	5.06	122.85	118.30
1	C	125	ASP	CB-CG-OD2	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2784	0	2848	7	0
1	B	2784	0	2848	9	0
1	C	2784	0	2848	6	0
1	D	2784	0	2848	11	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	44	0	26	2	0
3	B	44	0	26	2	0
3	C	44	0	26	2	0
3	D	44	0	26	2	0
4	A	19	0	10	1	0
4	B	19	0	10	1	0
4	C	19	0	10	0	0
4	D	19	0	10	0	0
5	A	203	0	0	0	0
5	B	216	0	0	0	0
5	C	223	0	0	0	0
5	D	263	0	0	0	0
All	All	12301	0	11536	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:100:CYS:HB2	1:D:112:LEU:HD12	1.86	0.57
1:D:40:MET:CA	1:D:40:MET:HE2	2.35	0.56
1:D:40:MET:HA	1:D:40:MET:HE2	1.88	0.55
1:D:70:ALA:HB1	1:D:166:LEU:HD22	1.89	0.54
1:A:279:LEU:HD22	1:A:312:ARG:HD3	1.89	0.52
1:C:295:PRO:HG3	1:D:305:PRO:HG2	1.93	0.51
1:A:178:THR:HG21	3:A:377:NAD:C4N	2.43	0.49
1:B:178:THR:HG21	3:B:377:NAD:C4N	2.44	0.48
1:B:174:CYS:SG	3:B:377:NAD:H5N	2.55	0.47
1:D:40:MET:CE	1:D:40:MET:CA	2.92	0.46
1:A:57:LEU:HD21	4:A:378[A]:DFB:HC5	1.96	0.46
1:C:90:ILE:HD12	1:C:160:ILE:HG21	1.97	0.46
1:D:61:LEU:HB3	1:D:62:PRO:HA	1.98	0.46
1:C:305:PRO:HG2	1:D:295:PRO:HG3	1.99	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:295:PRO:HG3	1:B:305:PRO:HG2	1.99	0.45
1:D:271:ARG:HB2	1:D:274:THR:OG1	2.17	0.44
1:B:118:MET:HE3	1:B:118:MET:HB3	1.85	0.44
1:B:318:ILE:HD12	4:B:378[B]:DFB:HC5	2.00	0.44
1:D:178:THR:HG21	3:D:377:NAD:C4N	2.47	0.43
1:C:242:ASN:OD1	1:C:244:GLN:HB2	2.18	0.43
1:A:292:VAL:O	3:A:377:NAD:H2N	2.20	0.42
1:A:32:LYS:O	1:A:77:GLY:HA3	2.20	0.42
1:C:292:VAL:O	3:C:377:NAD:H2N	2.20	0.42
1:A:305:PRO:HG2	1:B:295:PRO:HG3	2.03	0.41
1:B:70:ALA:HB1	1:B:166:LEU:HD22	2.00	0.41
1:B:350:LEU:O	1:B:372:LEU:HA	2.20	0.41
1:B:47:ARG:HA	1:B:47:ARG:HD3	1.92	0.41
1:D:174:CYS:SG	3:D:377:NAD:H5N	2.61	0.41
1:C:178:THR:HG21	3:C:377:NAD:C4N	2.51	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	372/374 (100%)	360 (97%)	12 (3%)	0	100	100
1	B	372/374 (100%)	361 (97%)	11 (3%)	0	100	100
1	C	372/374 (100%)	360 (97%)	12 (3%)	0	100	100
1	D	372/374 (100%)	359 (96%)	13 (4%)	0	100	100
All	All	1488/1496 (100%)	1440 (97%)	48 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/308 (100%)	303 (98%)	5 (2%)	68	58
1	B	308/308 (100%)	302 (98%)	6 (2%)	62	50
1	C	308/308 (100%)	304 (99%)	4 (1%)	73	66
1	D	308/308 (100%)	303 (98%)	5 (2%)	68	58
All	All	1232/1232 (100%)	1212 (98%)	20 (2%)	68	58

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	SER
1	A	78	GLU
1	A	251	GLN
1	A	279	LEU
1	A	300	ASN
1	B	1	SER
1	B	116	LEU
1	B	118	MET
1	B	239	GLU
1	B	256	GLU
1	B	300	ASN
1	C	1	SER
1	C	16	GLU
1	C	116	LEU
1	C	185	LYS
1	D	1	SER
1	D	164	SER
1	D	239	GLU
1	D	251	GLN
1	D	300	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	300	ASN
1	B	300	ASN
1	C	124	GLN
1	C	300	ASN
1	D	300	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 8 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	NAD	A	377	-	41,48,48	0.56	0	43,73,73	1.16	3 (6%)
4	DFB	A	378[A]	-	10,10,10	3.48	2 (20%)	12,13,13	2.18	3 (25%)
4	DFB	A	378[B]	-	10,10,10	3.28	2 (20%)	12,13,13	1.45	2 (16%)
3	NAD	B	377	-	41,48,48	0.64	0	43,73,73	1.42	4 (9%)
4	DFB	B	378[A]	-	10,10,10	3.38	2 (20%)	12,13,13	2.10	3 (25%)
4	DFB	B	378[B]	-	10,10,10	3.52	2 (20%)	12,13,13	1.67	3 (25%)
3	NAD	C	377	-	41,48,48	0.68	1 (2%)	43,73,73	1.25	4 (9%)
4	DFB	C	378[A]	-	10,10,10	3.47	2 (20%)	12,13,13	2.05	3 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	DFB	C	378[B]	-	10,10,10	3.52	2 (20%)	12,13,13	1.79	2 (16%)
3	NAD	D	377	-	41,48,48	0.60	0	43,73,73	1.34	3 (6%)
4	DFB	D	378[A]	-	10,10,10	3.38	2 (20%)	12,13,13	1.64	2 (16%)
4	DFB	D	378[B]	-	10,10,10	3.40	2 (20%)	12,13,13	2.28	3 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	377	-	-	0/22/62/62	0/5/5/5
4	DFB	A	378[A]	-	-	0/2/2/2	0/1/1/1
4	DFB	A	378[B]	-	-	0/2/2/2	0/1/1/1
3	NAD	B	377	-	-	0/22/62/62	0/5/5/5
4	DFB	B	378[A]	-	-	0/2/2/2	0/1/1/1
4	DFB	B	378[B]	-	-	0/2/2/2	0/1/1/1
3	NAD	C	377	-	-	0/22/62/62	0/5/5/5
4	DFB	C	378[A]	-	-	0/2/2/2	0/1/1/1
4	DFB	C	378[B]	-	-	0/2/2/2	0/1/1/1
3	NAD	D	377	-	-	0/22/62/62	0/5/5/5
4	DFB	D	378[A]	-	-	0/2/2/2	0/1/1/1
4	DFB	D	378[B]	-	-	0/2/2/2	0/1/1/1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	377	NAD	C4N-C3N	2.01	1.42	1.39
4	A	378[A]	DFB	C3-C2	4.86	1.47	1.37
4	D	378[B]	DFB	C3-C2	4.96	1.47	1.37
4	B	378[A]	DFB	C3-C2	5.19	1.48	1.37
4	C	378[A]	DFB	C3-C2	5.27	1.48	1.37
4	A	378[B]	DFB	C3-C2	5.28	1.48	1.37
4	C	378[B]	DFB	C3-C2	5.35	1.48	1.37
4	D	378[A]	DFB	C3-C2	5.36	1.48	1.37
4	B	378[B]	DFB	C3-C2	5.44	1.48	1.37
4	A	378[B]	DFB	C1-C2	8.82	1.48	1.38
4	D	378[A]	DFB	C1-C2	9.13	1.48	1.38
4	B	378[A]	DFB	C1-C2	9.26	1.48	1.38
4	D	378[B]	DFB	C1-C2	9.46	1.48	1.38
4	C	378[A]	DFB	C1-C2	9.51	1.48	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	378[B]	DFB	C1-C2	9.63	1.48	1.38
4	C	378[B]	DFB	C1-C2	9.65	1.48	1.38
4	A	378[A]	DFB	C1-C2	9.78	1.49	1.38

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	377	NAD	N3A-C2A-N1A	-6.04	123.60	128.86
3	D	377	NAD	N3A-C2A-N1A	-4.91	124.58	128.86
3	C	377	NAD	N3A-C2A-N1A	-4.87	124.61	128.86
3	A	377	NAD	N3A-C2A-N1A	-4.72	124.75	128.86
4	B	378[A]	DFB	C4-C3-C2	-3.10	118.41	121.27
4	C	378[A]	DFB	C4-C3-C2	-2.66	118.81	121.27
4	A	378[A]	DFB	C4-C3-C2	-2.66	118.82	121.27
4	D	378[B]	DFB	C4-C3-C2	-2.49	118.97	121.27
4	B	378[B]	DFB	C4-C3-C2	-2.11	119.32	121.27
3	B	377	NAD	C5A-C6A-N1A	-2.10	113.35	119.70
3	C	377	NAD	C4B-O4B-C1B	-2.07	107.56	109.77
3	D	377	NAD	O2N-PN-O1N	2.20	123.68	112.28
3	A	377	NAD	C2A-N1A-C6A	2.39	122.95	118.77
4	A	378[B]	DFB	F2-C2-C1	2.45	120.44	117.85
3	C	377	NAD	C2A-N1A-C6A	2.48	123.11	118.77
4	A	378[B]	DFB	C6-C1-C2	2.51	118.26	116.43
3	A	377	NAD	C5A-C6A-N6A	2.70	125.98	120.47
4	C	378[B]	DFB	C6-C1-C2	2.72	118.41	116.43
3	C	377	NAD	C5A-C6A-N6A	2.77	126.12	120.47
3	B	377	NAD	C5A-C6A-N6A	2.82	126.21	120.47
4	A	378[A]	DFB	C6-C1-C2	2.82	118.48	116.43
3	D	377	NAD	C2A-N1A-C6A	2.89	123.83	118.77
4	D	378[A]	DFB	F2-C2-C1	2.91	120.93	117.85
4	B	378[B]	DFB	C6-C1-C2	3.03	118.64	116.43
4	B	378[B]	DFB	F2-C2-C1	3.35	121.40	117.85
4	B	378[A]	DFB	F2-C2-C1	3.42	121.47	117.85
4	D	378[A]	DFB	C6-C1-C2	3.45	118.94	116.43
3	B	377	NAD	C2A-N1A-C6A	3.50	124.89	118.77
4	C	378[A]	DFB	C6-C1-C2	3.90	119.27	116.43
4	C	378[A]	DFB	F2-C2-C1	4.17	122.27	117.85
4	B	378[A]	DFB	C6-C1-C2	4.27	119.54	116.43
4	D	378[B]	DFB	C6-C1-C2	4.53	119.73	116.43
4	C	378[B]	DFB	F2-C2-C1	4.58	122.70	117.85
4	D	378[B]	DFB	F2-C2-C1	5.02	123.16	117.85
4	A	378[A]	DFB	F2-C2-C1	5.70	123.89	117.85

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	377	NAD	2	0
4	A	378[A]	DFB	1	0
3	B	377	NAD	2	0
4	B	378[B]	DFB	1	0
3	C	377	NAD	2	0
3	D	377	NAD	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	374/374 (100%)	-0.35	0 100 100	10, 18, 27, 34	0
1	B	374/374 (100%)	-0.35	1 (0%) 93 92	12, 19, 28, 37	0
1	C	374/374 (100%)	-0.35	0 100 100	12, 18, 28, 34	0
1	D	374/374 (100%)	-0.39	1 (0%) 93 92	11, 17, 26, 34	0
All	All	1496/1496 (100%)	-0.36	2 (0%) 95 93	10, 18, 27, 37	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	SER	2.6
1	D	84	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	DFB	A	378[B]	10/10	0.96	0.09	1.04	14,15,17,18	9
4	DFB	B	378[A]	10/10	0.97	0.08	0.75	18,19,20,21	9
4	DFB	B	378[B]	10/10	0.97	0.08	0.75	18,19,21,21	9
4	DFB	A	378[A]	10/10	0.96	0.09	0.62	14,18,19,19	9
4	DFB	D	378[A]	10/10	0.96	0.09	0.42	17,18,19,20	9
4	DFB	C	378[B]	10/10	0.96	0.08	0.35	17,19,19,21	9
4	DFB	C	378[A]	10/10	0.96	0.08	0.16	17,18,19,19	9
4	DFB	D	378[B]	10/10	0.96	0.09	0.14	18,20,22,23	9
3	NAD	A	377	44/44	0.97	0.07	-0.78	10,15,21,22	0
2	ZN	A	375	1/1	1.00	0.04	-1.04	23,23,23,23	0
3	NAD	D	377	44/44	0.98	0.06	-1.05	10,14,16,20	0
3	NAD	C	377	44/44	0.97	0.06	-1.10	11,16,22,23	0
3	NAD	B	377	44/44	0.98	0.05	-1.33	10,16,19,19	0
2	ZN	D	375	1/1	1.00	0.04	-1.40	22,22,22,22	0
2	ZN	A	376	1/1	1.00	0.02	-1.47	17,17,17,17	0
2	ZN	C	376	1/1	1.00	0.03	-1.58	18,18,18,18	0
2	ZN	B	376	1/1	1.00	0.04	-1.59	21,21,21,21	0
2	ZN	D	376	1/1	0.99	0.03	-1.68	18,18,18,18	0
2	ZN	B	375	1/1	1.00	0.04	-2.31	23,23,23,23	0
2	ZN	C	375	1/1	1.00	0.02	-3.41	24,24,24,24	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.