



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 09:10 pm GMT

PDB ID : 4MKH
Title : Crystal structure of a stable adenylate kinase variant AKv18
Authors : Moon, S.; Bae, E.
Deposited on : 2013-09-05
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2279 (1.50-1.50)
Clashscore	112137	2503 (1.50-1.50)
Ramachandran outliers	110173	2445 (1.50-1.50)
Sidechain outliers	110143	2443 (1.50-1.50)
RSRZ outliers	101464	2305 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	222	<div> <div></div> <div>86%</div> <div>12%</div> <div></div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 1914 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenylate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	218	Total	C	N	O	S	0	0	0
			1706	1071	291	332	12			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP P16304
A	0	GLY	-	EXPRESSION TAG	UNP P16304
A	3	ILE	LEU	ENGINEERED MUTATION	UNP P16304
A	16	LEU	GLN	ENGINEERED MUTATION	UNP P16304
A	17	ALA	GLY	ENGINEERED MUTATION	UNP P16304
A	19	LYS	ARG	ENGINEERED MUTATION	UNP P16304
A	23	LYS	ASP	ENGINEERED MUTATION	UNP P16304
A	69	ARG	LYS	ENGINEERED MUTATION	UNP P16304
A	73	SER	GLY	ENGINEERED MUTATION	UNP P16304
A	75	SER	ASP	ENGINEERED MUTATION	UNP P16304
A	103	MET	TYR	ENGINEERED MUTATION	UNP P16304
A	105	ARG	LYS	ENGINEERED MUTATION	UNP P16304
A	114	GLN	GLU	ENGINEERED MUTATION	UNP P16304
A	116	ARG	ASP	ENGINEERED MUTATION	UNP P16304
A	118	GLU	ASP	ENGINEERED MUTATION	UNP P16304
A	119	GLU	VAL	ENGINEERED MUTATION	UNP P16304
A	169	THR	SER	ENGINEERED MUTATION	UNP P16304
A	179	MET	THR	ENGINEERED MUTATION	UNP P16304
A	180	ALA	GLN	ENGINEERED MUTATION	UNP P16304
A	184	ALA	ASP	ENGINEERED MUTATION	UNP P16304
A	187	ASP	SER	ENGINEERED MUTATION	UNP P16304
A	188	SER	GLU	ENGINEERED MUTATION	UNP P16304
A	190	GLU	GLY	ENGINEERED MUTATION	UNP P16304
A	191	VAL	TYR	ENGINEERED MUTATION	UNP P16304
A	193	VAL	ALA	ENGINEERED MUTATION	UNP P16304
A	198	GLU	GLN	ENGINEERED MUTATION	UNP P16304
A	199	ARG	GLN	ENGINEERED MUTATION	UNP P16304

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	202	GLU	GLN	ENGINEERED MUTATION	UNP P16304
A	205	PHE	TYR	ENGINEERED MUTATION	UNP P16304
A	209	ASP	LYS	ENGINEERED MUTATION	UNP P16304
A	210	VAL	ASP	ENGINEERED MUTATION	UNP P16304
A	211	ILE	LEU	ENGINEERED MUTATION	UNP P16304
A	213	LEU	-	EXPRESSION TAG	UNP P16304
A	214	GLU	-	EXPRESSION TAG	UNP P16304
A	215	HIS	-	EXPRESSION TAG	UNP P16304
A	216	HIS	-	EXPRESSION TAG	UNP P16304
A	217	HIS	-	EXPRESSION TAG	UNP P16304
A	218	HIS	-	EXPRESSION TAG	UNP P16304
A	219	HIS	-	EXPRESSION TAG	UNP P16304
A	220	HIS	-	EXPRESSION TAG	UNP P16304

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0

- Molecule 3 is BIS(ADENOSINE)-5'-PENTAPHOSPHATE (three-letter code: AP5) (formula: C₂₀H₂₉N₁₀O₂₂P₅).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O P 60 20 10 24 6	0	1

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	135	Total	O	0	0
			135	135		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adenylate kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.19Å 55.19Å 46.38Å 90.00° 110.61° 90.00°	Depositor
Resolution (Å)	27.60 – 1.50 27.60 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.6 (27.60-1.50) 99.7 (27.60-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.171 , 0.218 0.174 , 0.222	Depositor DCC
R_{free} test set	1687 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	18.4	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.026 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1914	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, EDO, AP5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.30	2/1732 (0.1%)	1.49	23/2336 (1.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	86	PHE	CG-CD1	5.66	1.47	1.38
1	A	105	ARG	CZ-NH2	5.46	1.40	1.33

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	88	ARG	NE-CZ-NH2	14.48	127.54	120.30
1	A	88	ARG	NE-CZ-NH1	-12.90	113.85	120.30
1	A	160	ARG	NE-CZ-NH1	11.19	125.89	120.30
1	A	1	MET	CG-SD-CE	-11.04	82.53	100.20
1	A	123	ARG	NE-CZ-NH2	-10.03	115.28	120.30
1	A	69	ARG	NE-CZ-NH1	9.35	124.97	120.30
1	A	54	ASP	CB-CG-OD1	-9.13	110.08	118.30
1	A	16	LEU	CA-CB-CG	9.11	136.25	115.30
1	A	160	ARG	NE-CZ-NH2	-8.70	115.95	120.30
1	A	171	ARG	NE-CZ-NH1	-7.96	116.32	120.30
1	A	105	ARG	NE-CZ-NH1	7.84	124.22	120.30
1	A	151	ASP	CB-CG-OD1	6.25	123.93	118.30
1	A	200	ASP	CB-CG-OD1	6.18	123.86	118.30
1	A	108	ASP	CB-CG-OD2	6.15	123.83	118.30
1	A	103	MET	CG-SD-CE	5.92	109.68	100.20
1	A	6	MET	CG-SD-CE	5.71	109.34	100.20
1	A	139	LEU	CB-CG-CD1	-5.61	101.47	111.00
1	A	88	ARG	CD-NE-CZ	5.48	131.27	123.60
1	A	79	ARG	NE-CZ-NH1	-5.42	117.59	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	76	ASP	CB-CG-OD1	5.17	122.96	118.30
1	A	105	ARG	CB-CG-CD	5.14	124.95	111.60
1	A	109	TYR	CB-CG-CD2	-5.02	117.99	121.00
1	A	186	TYR	CD1-CE1-CZ	-5.01	115.29	119.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1706	0	1708	4	0
2	A	1	0	0	0	0
3	A	60	0	0	0	0
4	A	12	0	18	0	0
5	A	135	0	0	0	0
All	All	1914	0	1726	4	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:16:LEU:HD13	1:A:113:ILE:HD13	1.86	0.55
1:A:111:ILE:HD13	1:A:208:VAL:HG13	2.00	0.43
1:A:17:ALA:HA	1:A:82:LEU:HD21	2.02	0.41
1:A:21:VAL:HG21	1:A:28:HIS:HB2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	216/222 (97%)	215 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/191 (98%)	187 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	AP5	A	302[A]	-	48,62,62	1.36	4 (8%)	43,98,98	1.99	12 (27%)
3	AP5	A	302[B]	-	42,58,62	1.44	4 (9%)	38,85,98	2.01	11 (28%)
4	EDO	A	303	-	3,3,3	0.55	0	2,2,2	0.83	0
4	EDO	A	304	-	3,3,3	0.85	0	2,2,2	0.74	0
4	EDO	A	305	-	3,3,3	0.87	0	2,2,2	1.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AP5	A	302[A]	-	-	0/36/76/76	0/6/6/6
3	AP5	A	302[B]	-	-	0/18/62/76	0/6/6/6
4	EDO	A	303	-	-	0/1/1/1	0/0/0/0
4	EDO	A	304	-	-	0/1/1/1	0/0/0/0
4	EDO	A	305	-	-	0/1/1/1	0/0/0/0

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	302[B]	AP5	PA-O2A	-2.24	1.43	1.55
3	A	302[A]	AP5	PA-O2A	-2.24	1.43	1.55
3	A	302[B]	AP5	C2A-N1A	2.14	1.37	1.33
3	A	302[A]	AP5	C2A-N1A	2.14	1.37	1.33
3	A	302[B]	AP5	O4J-C1J	2.98	1.45	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	302[A]	AP5	O4J-C1J	2.98	1.45	1.41
3	A	302[B]	AP5	O4F-C1F	5.93	1.49	1.41
3	A	302[A]	AP5	O4F-C1F	5.93	1.49	1.41

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	302[B]	AP5	N3A-C2A-N1A	-6.94	122.82	128.86
3	A	302[A]	AP5	N3A-C2A-N1A	-6.94	122.82	128.86
3	A	302[B]	AP5	N3B-C2B-N1B	-3.48	125.83	128.86
3	A	302[A]	AP5	N3B-C2B-N1B	-3.48	125.83	128.86
3	A	302[B]	AP5	C5B-C6B-N1B	-2.69	111.56	119.70
3	A	302[A]	AP5	C5B-C6B-N1B	-2.69	111.56	119.70
3	A	302[B]	AP5	C5A-C6A-N1A	-2.16	113.17	119.70
3	A	302[A]	AP5	C5A-C6A-N1A	-2.16	113.17	119.70
3	A	302[B]	AP5	C1F-N9A-C4A	-2.10	123.01	126.64
3	A	302[A]	AP5	C1F-N9A-C4A	-2.10	123.01	126.64
3	A	302[B]	AP5	C4F-O4F-C1F	-2.10	107.54	109.77
3	A	302[A]	AP5	C4F-O4F-C1F	-2.10	107.54	109.77
3	A	302[B]	AP5	N6B-C6B-N1B	2.11	122.95	118.77
3	A	302[A]	AP5	N6B-C6B-N1B	2.11	122.95	118.77
3	A	302[B]	AP5	C5B-C6B-N6B	2.34	125.24	120.47
3	A	302[A]	AP5	C5B-C6B-N6B	2.34	125.24	120.47
3	A	302[B]	AP5	C2B-N1B-C6B	2.70	123.50	118.77
3	A	302[A]	AP5	C2B-N1B-C6B	2.70	123.50	118.77
3	A	302[A]	AP5	O2D-PD-O1D	2.99	127.75	112.28
3	A	302[B]	AP5	N6A-C6A-N1A	3.33	125.38	118.77
3	A	302[A]	AP5	N6A-C6A-N1A	3.33	125.38	118.77
3	A	302[B]	AP5	C2A-N1A-C6A	4.36	126.40	118.77
3	A	302[A]	AP5	C2A-N1A-C6A	4.36	126.40	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	218/222 (98%)	0.01	2 (0%) 84 87	14, 20, 30, 40	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	ILE	2.6
1	A	152	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	EDO	A	304	4/4	0.98	0.15	1.53	20,22,22,24	0
4	EDO	A	305	4/4	0.95	0.09	0.68	27,27,28,29	0
4	EDO	A	303	4/4	0.95	0.10	0.14	25,25,28,30	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	AP5	A	302[B]	55/57	0.96	0.09	0.03	13,19,25,27	3
3	AP5	A	302[A]	57/57	0.96	0.09	0.03	13,19,25,27	5
2	ZN	A	301	1/1	0.99	0.04	-1.46	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.