



wwPDB NMR Structure Validation Summary Report ⓘ

Feb 13, 2017 – 01:07 am GMT

PDB ID : 2MLW
Title : New Cyt-like delta-endotoxins from *Dickeya dadantii* - CytC protein
Authors : Loth, K.; Costechareyre, D.; Effantin, G.; Rahbe, Y.; Condemine, G.; Landon, C.; Da Silva, P.
Deposited on : 2014-03-05

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk28760
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 79%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	203	<div> <div></div> <div>29%</div> <div>40%</div> <div>14%</div> <div>16%</div> </div>

2 Ensemble composition and analysis

This entry contains 20 models. Model 11 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:97, A:125-A:171, A:178-A:203 (170)	0.75	11

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 6 single-model clusters were found.

Cluster number	Models
1	1, 5, 11, 12, 13, 15, 16, 19
2	3, 4, 6, 7, 9, 10
Single-model clusters	2; 8; 14; 17; 18; 20

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 3146 atoms, of which 1568 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Type-1Ba cytolytic delta-endotoxin.


Mol	Chain	Residues	Atoms						Trace
1	A	203	Total	C	H	N	O	S	0
			3146	1006	1568	254	313	5	

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Type-1Ba cytolytic delta-endotoxin



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 11. Colouring as in section 4.1 above.

- Molecule 1: Type-1Ba cytolytic delta-endotoxin



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 500 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA	structure solution	2.3
ARIA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2mlw_cs.str
Number of chemical shift lists	1
Total number of shifts	2103
Number of shifts mapped to atoms	2103
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	79%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.42±0.01	0±0/1325 (0.0±0.0%)	0.59±0.02	0±0/1802 (0.0±0.0%)
All	All	0.42	0/26500 (0.0%)	0.59	1/36040 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.5±0.8
All	All	0	10

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	145	ILE	N-CA-CB	5.39	123.20	110.80	19	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	170	ALA	Peptide	4
1	A	169	THR	Peptide	3
1	A	141	TYR	Peptide	2
1	A	142	SER	Peptide	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1302	1311	1307	100±9
All	All	26040	26220	26140	2006

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

5 of 519 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:127:ASN:O	1:A:137:THR:HA	0.91	1.65	4	18
1:A:165:VAL:HB	1:A:183:THR:O	0.85	1.70	17	19
1:A:70:MET:HB3	1:A:92:ILE:HG21	0.84	1.46	4	2
1:A:97:GLY:O	1:A:199:VAL:HA	0.83	1.73	20	10
1:A:161:LEU:O	1:A:186:LYS:HB2	0.83	1.74	11	20

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	168/203 (83%)	115±1 (68±1%)	39±2 (23±1%)	14±2 (8±1%)	2	14
All	All	3360/4060 (83%)	2299 (68%)	785 (23%)	276 (8%)	2	14

5 of 31 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	201	PRO	20
1	A	4	GLU	20
1	A	18	PHE	20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	12	THR	20
1	A	6	TYR	20

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/174 (82%)	109±3 (77±2%)	33±3 (23±2%)	3	29
All	All	2840/3480 (82%)	2178 (77%)	662 (23%)	3	29

5 of 79 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	33	LEU	20
1	A	68	ASN	20
1	A	19	LEU	20
1	A	168	ASN	20
1	A	46	GLN	20

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 79% for the well-defined parts and 76% for the entire structure.

7.1 Chemical shift list 1

File name: 2mlw_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	2103
Number of shifts mapped to atoms	2103
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	27

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	200	-0.24 ± 0.17	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	176	-0.07 ± 0.26	None needed (< 0.5 ppm)
$^{13}\text{C}'$	182	-0.08 ± 0.09	None needed (< 0.5 ppm)
^{15}N	189	0.76 ± 0.29	Should be applied

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 79%, i.e. 1589 atoms were assigned a chemical shift out of a possible 2004. 16 out of 29 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	814/836 (97%)	329/333 (99%)	324/340 (95%)	161/163 (99%)
Sidechain	739/1031 (72%)	430/596 (72%)	288/403 (71%)	21/32 (66%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	36/137 (26%)	35/71 (49%)	0/57 (0%)	1/9 (11%)
Overall	1589/2004 (79%)	794/1000 (79%)	612/800 (76%)	183/204 (90%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	181	ARG	NE	111.90	92.63 – 76.73	17.1
1	A	51	LYS	CE	52.53	46.00 – 37.80	13.0
1	A	51	LYS	NZ	71.88	49.86 – 18.16	11.9
1	A	178	LYS	NZ	71.56	49.86 – 18.16	11.8
1	A	107	ILE	CG2	29.33	24.63 – 10.43	8.3
1	A	100	GLN	CG	24.60	39.38 – 28.18	-8.2
1	A	7	PRO	CG	36.02	32.66 – 21.76	8.1
1	A	6	TYR	HD2	9.15	8.44 – 5.44	7.4
1	A	6	TYR	HD1	9.15	8.44 – 5.44	7.4
1	A	174	ILE	CG2	27.08	24.63 – 10.43	6.7
1	A	91	LYS	CE	47.32	46.00 – 37.80	6.6
1	A	92	ILE	CG2	26.77	24.63 – 10.43	6.5
1	A	180	TRP	CB	42.17	40.02 – 19.92	6.1
1	A	189	ILE	CG2	26.06	24.63 – 10.43	6.0
1	A	201	PRO	CD	56.26	55.31 – 45.41	6.0
1	A	62	LEU	CG	33.51	32.55 – 21.05	5.8
1	A	107	ILE	CD1	23.20	21.91 – 5.01	5.8
1	A	49	LEU	CG	33.32	32.55 – 21.05	5.7
1	A	15	TYR	HD1	8.61	8.44 – 5.44	5.6
1	A	15	TYR	HD2	8.61	8.44 – 5.44	5.6
1	A	2	SER	HB2	5.32	5.18 – 2.58	5.5
1	A	186	LYS	HE2	1.88	3.87 – 1.97	-5.5
1	A	14	PRO	CD	55.79	55.31 – 45.41	5.5
1	A	112	PHE	HZ	9.29	9.11 – 4.91	5.4
1	A	115	LEU	CD1	16.03	32.77 – 16.57	-5.3
1	A	73	LYS	CB	41.94	41.68 – 23.88	5.1
1	A	2	SER	HB3	5.29	5.25 – 2.45	5.1

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from

the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

