



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 13, 2017 – 05:27 PM EST

PDB ID : 5MLN
Title : The crystal structure of alcohol dehydrogenase 10 from *Candida magnoliae*
Authors : Castellanos, J.R.G.; Mattevi, A.
Deposited on : unknown
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2696 (1.60-1.60)
Clashscore	112137	2967 (1.60-1.60)
Ramachandran outliers	110173	2887 (1.60-1.60)
Sidechain outliers	110143	2886 (1.60-1.60)
RSRZ outliers	101464	2714 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	241	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>11%</div> <div>••</div> </div> </div>
1	B	241	<div> <div>88%</div> <div>10%</div> <div>•</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PEG	A	302	-	-	-	X
3	PEG	A	303	-	-	X	X
3	PEG	A	304	-	-	-	X
3	PEG	B	302	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4033 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alcohol dehydrogenase 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	237	Total	C	N	O	S	0	2	0
			1766	1122	300	342	2			
1	B	238	Total	C	N	O	S	0	0	0
			1762	1119	300	341	2			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	30	ASN	SER	conflict	UNP B8K244
A	81	ARG	SER	conflict	UNP B8K244
A	133	VAL	ALA	conflict	UNP B8K244
A	135	LYS	ASN	conflict	UNP B8K244
B	30	ASN	SER	conflict	UNP B8K244
B	81	ARG	SER	conflict	UNP B8K244
B	133	VAL	ALA	conflict	UNP B8K244
B	135	LYS	ASN	conflict	UNP B8K244

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



- PEG
-
- The diagram illustrates the chemical structure of Polyethylene Glycol (PEG). It features a zigzag chain of carbon (C) and oxygen (O) atoms. The chain starts with a terminal hydroxyl group (HO) labeled O1, followed by a carbon atom labeled C1, then another carbon atom labeled C2, an oxygen atom labeled O2, a carbon atom labeled C3, another carbon atom labeled C4, and finally a terminal hydroxyl group (OH) labeled O4. The atoms are color-coded: carbon atoms (C1, C2, C3, C4) are green, and oxygen atoms (O1, O2, O4) are red. The hydroxyl groups are shown as HO and OH.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 7 4 3	0	0
3	A	1	Total C O 7 4 3	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	178	Total	O	0	0
			178	178		
4	B	203	Total	O	0	0
			203	203		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alcohol dehydrogenase 3



• Molecule 1: Alcohol dehydrogenase 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α , β , γ	74.18Å 80.42Å 80.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.30 – 1.60 40.26 – 1.59	Depositor EDS
% Data completeness (in resolution range)	99.9 (40.30-1.60) 99.9 (40.26-1.59)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.13 (at 1.59Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.146 , 0.180 0.160 , 0.188	Depositor DCC
R_{free} test set	3197 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	18.7	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 61.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.018 for -h,l,k	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4033	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.21	5/1804 (0.3%)	1.11	5/2455 (0.2%)
1	B	1.25	2/1794 (0.1%)	1.11	7/2442 (0.3%)
All	All	1.23	7/3598 (0.2%)	1.11	12/4897 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	241	ALA	C-O	8.34	1.39	1.23
1	B	209	TRP	CG-CD1	6.22	1.45	1.36
1	A	37	SER	CB-OG	5.32	1.49	1.42
1	A	237	ASN	CG-OD1	5.15	1.35	1.24
1	B	89	ASN	N-CA	-5.09	1.36	1.46

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	63	ASP	CB-CG-OD1	6.31	123.98	118.30
1	B	174	GLU	OE1-CD-OE2	-6.27	115.77	123.30
1	A	241	ALA	CA-C-O	-5.93	107.64	120.10
1	B	145	VAL	CG1-CB-CG2	5.91	120.36	110.90
1	B	125	LEU	CB-CG-CD2	5.73	120.74	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	199	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1766	0	1795	23	0
1	B	1762	0	1790	10	0
2	A	48	0	25	1	0
2	B	48	0	25	1	0
3	A	21	0	30	10	0
3	B	7	0	10	0	0
4	A	178	0	0	7	6
4	B	203	0	0	6	7
All	All	4033	0	3675	35	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 35 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:241:ALA:O	4:B:401:HOH:O	1.99	0.81
1:A:6:ASN:HD21	3:A:303:PEG:C4	1.94	0.80
1:A:129:SER:OG	4:A:402:HOH:O	2.03	0.76
1:B:189:TYR:HB2	4:B:559:HOH:O	1.87	0.75
1:A:6:ASN:HD21	3:A:303:PEG:H41	1.53	0.72

The worst 5 of 7 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:448:HOH:O	4:B:468:HOH:O[2_555]	1.26	0.94

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:417:HOH:O	4:B:550:HOH:O[2_555]	1.54	0.66
4:B:520:HOH:O	4:B:520:HOH:O[2_555]	1.70	0.50
4:A:564:HOH:O	4:B:569:HOH:O[1_556]	1.92	0.28
4:A:417:HOH:O	4:B:522:HOH:O[2_555]	1.96	0.24

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	237/241 (98%)	231 (98%)	6 (2%)	0	100	100
1	B	236/241 (98%)	230 (98%)	5 (2%)	1 (0%)	38	16
All	All	473/482 (98%)	461 (98%)	11 (2%)	1 (0%)	51	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	129	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/189 (99%)	181 (97%)	6 (3%)	44	17
1	B	186/189 (98%)	181 (97%)	5 (3%)	50	22
All	All	373/378 (99%)	362 (97%)	11 (3%)	46	20

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	198	ILE
1	A	231	THR
1	B	133	VAL
1	A	193	GLU
1	B	73	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	232	ASN
1	A	237	ASN
1	B	186	ASN
1	A	186	ASN
1	B	153	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	A	301	-	44,52,52	1.25	6 (13%)	51,80,80	1.22	6 (11%)
3	PEG	A	302	-	6,6,6	0.53	0	5,5,5	0.95	0
3	PEG	A	303	-	6,6,6	0.62	0	5,5,5	1.04	0
3	PEG	A	304	-	6,6,6	0.44	0	5,5,5	0.75	0
2	NAP	B	301	-	44,52,52	1.53	4 (9%)	51,80,80	1.87	10 (19%)
3	PEG	B	302	-	6,6,6	0.64	0	5,5,5	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	301	-	-	0/27/67/67	0/5/5/5
3	PEG	A	302	-	-	0/4/4/4	0/0/0/0
3	PEG	A	303	-	-	0/4/4/4	0/0/0/0
3	PEG	A	304	-	-	0/4/4/4	0/0/0/0
2	NAP	B	301	-	-	0/27/67/67	0/5/5/5
3	PEG	B	302	-	-	0/4/4/4	0/0/0/0

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	NAP	C2D-C1D	-3.09	1.48	1.53
2	A	301	NAP	C6N-C5N	-2.62	1.32	1.38
2	A	301	NAP	P2B-O3X	-2.00	1.46	1.54
2	A	301	NAP	C5A-C4A	2.04	1.45	1.40
2	A	301	NAP	O4D-C1D	2.06	1.44	1.41

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	NAP	N3A-C2A-N1A	-7.04	122.72	128.86
2	B	301	NAP	C5N-C4N-C3N	-5.90	113.41	120.35
2	B	301	NAP	C4D-O4D-C1D	-3.30	106.26	109.77
2	A	301	NAP	C4A-C5A-N7A	-2.66	106.84	109.41
2	A	301	NAP	C1B-N9A-C4A	-2.61	122.12	126.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	NAP	1	0
3	A	303	PEG	8	0
3	A	304	PEG	2	0
2	B	301	NAP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/241 (98%)	-0.34	5 (2%) 64 64	12, 19, 38, 73	0
1	B	238/241 (98%)	-0.38	0 100 100	12, 20, 39, 47	0
All	All	475/482 (98%)	-0.36	5 (1%) 80 82	12, 19, 39, 73	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	133	VAL	7.2
1	A	129	SER	5.5
1	A	198	ILE	4.6
1	A	134	ASP	4.1
1	A	130	GLU	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	PEG	A	304	7/7	0.84	0.28	8.13	43,45,50,51	0
3	PEG	B	302	7/7	0.84	0.23	5.34	38,44,49,55	0
3	PEG	A	302	7/7	0.84	0.13	3.10	35,38,49,61	0
3	PEG	A	303	7/7	0.80	0.29	2.98	43,52,59,62	0
2	NAP	B	301	48/48	0.97	0.09	0.73	18,21,25,27	0
2	NAP	A	301	48/48	0.98	0.05	-0.81	14,16,20,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.