



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 31, 2017 – 06:21 PM EDT

PDB ID : 3MMA
Title : Dissimilatory sulfite reductase phosphate complex
Authors : Parey, K.; Warkentin, E.; Kroneck, P.M.H.; Ermler, U.
Deposited on : unknown
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	418	<div> <div>7%</div> <div> <div></div> <div>86%</div> <div>13%</div> </div> </div>
1	D	418	<div> <div>31%</div> <div> <div></div> <div>85%</div> <div>14%</div> </div> </div>
2	B	366	<div> <div>5%</div> <div> <div></div> <div>86%</div> <div>12%</div> </div> </div>
2	E	366	<div> <div>28%</div> <div> <div></div> <div>79%</div> <div>18%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria.

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SF4	E	585	-	-	X	-
5	PO4	A	590	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 12960 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Sulfite reductase, dissimilatory-type subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	417	Total	C	N	O	S	0	0	0
			3329	2134	557	612	26			
1	D	417	Total	C	N	O	S	0	0	0
			3329	2134	557	612	26			

- Molecule 2 is a protein called Sulfite reductase, dissimilatory-type subunit beta.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	363	Total	C	N	O	S	0	0	0
			2901	1862	491	526	22			
2	E	363	Total	C	N	O	S	0	0	0
			2901	1862	491	526	22			

- Molecule 3 is SIROHEME (three-letter code: SRM) (formula: $C_{42}H_{42}FeN_4O_{16}$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	
			63	42	1	4	16	
3	B	1	Total	C	Fe	N	O	
			63	42	1	4	16	
3	D	1	Total	C	Fe	N	O	
			63	42	1	4	16	
3	E	1	Total	C	Fe	N	O	
			63	42	1	4	16	

- Molecule 4 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Fe	S		
			8	4	4	0	0
4	A	1	Total	Fe	S		
			8	4	4	0	0
4	B	1	Total	Fe	S		
			8	4	4	0	0
4	B	1	Total	Fe	S		
			8	4	4	0	0
4	D	1	Total	Fe	S		
			8	4	4	0	0
4	D	1	Total	Fe	S		
			8	4	4	0	0
4	E	1	Total	Fe	S		
			8	4	4	0	0
4	E	1	Total	Fe	S		
			8	4	4	0	0

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	78	Total	O	0	0
			78	78		
6	B	84	Total	O	0	0
			84	84		
6	D	11	Total	O	0	0
			11	11		
6	E	6	Total	O	0	0
			6	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Sulfite reductase, dissimilatory-type subunit alpha




- Molecule 1: Sulfite reductase, dissimilatory-type subunit alpha





- Molecule 2: Sulfite reductase, dissimilatory-type subunit beta





● Molecule 2: Sulfite reductase, dissimilatory-type subunit beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	94.60Å 68.90Å 145.10Å 90.00° 107.50° 90.00°	Depositor
Resolution (Å)	48.34 – 2.30 48.33 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.34-2.30) 98.8 (48.33-2.30)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.81 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.6.0046	Depositor
R, R_{free}	0.192 , 0.236 0.211 , 0.247	Depositor DCC
R_{free} test set	3928 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	34.8	Xtriage
Anisotropy	0.261	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 43.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12960	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, PO4, SRM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	1/3416 (0.0%)	0.63	0/4610
1	D	0.42	1/3416 (0.0%)	0.53	0/4610
2	B	0.54	0/2984	0.65	1/4058 (0.0%)
2	E	0.40	0/2984	0.55	0/4058
All	All	0.48	2/12800 (0.0%)	0.59	1/17336 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	223	CYS	CB-SG	8.13	1.96	1.82
1	D	23	LYS	CD-CE	5.95	1.66	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	277	ARG	NE-CZ-NH1	5.13	122.87	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	317	PHE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3329	0	3276	45	0
1	D	3329	0	3276	44	0
2	B	2901	0	2837	37	0
2	E	2901	0	2838	53	0
3	A	63	0	34	12	0
3	B	63	0	34	6	0
3	D	63	0	34	6	0
3	E	63	0	34	11	0
4	A	16	0	0	0	0
4	B	16	0	0	0	0
4	D	16	0	0	1	0
4	E	16	0	0	2	0
5	A	5	0	0	2	0
6	A	78	0	0	1	0
6	B	84	0	0	1	0
6	D	11	0	0	2	0
6	E	6	0	0	0	0
All	All	12960	0	12363	167	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 167 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:570:SRM:HBD2	3:E:570:SRM:HDD2	1.33	1.07
1:A:403:LYS:H	2:E:261:ASN:HD21	1.07	0.97
2:E:230:LYS:HB2	2:E:231:ASN:HB2	1.48	0.95
1:D:317:PHE:HD2	2:E:180:ASN:HB3	1.35	0.88
2:E:177:CYS:HB2	4:E:585:SF4:S3	2.14	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	415/418 (99%)	402 (97%)	12 (3%)	1 (0%)	51	63
1	D	415/418 (99%)	393 (95%)	21 (5%)	1 (0%)	51	63
2	B	361/366 (99%)	343 (95%)	16 (4%)	2 (1%)	28	34
2	E	361/366 (99%)	331 (92%)	25 (7%)	5 (1%)	13	13
All	All	1552/1568 (99%)	1469 (95%)	74 (5%)	9 (1%)	28	34

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	318	VAL
2	E	317	HIS
1	D	318	VAL
2	E	76	LYS
2	E	231	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	353/354 (100%)	345 (98%)	8 (2%)	56	73
1	D	353/354 (100%)	344 (98%)	9 (2%)	53	70
2	B	314/317 (99%)	304 (97%)	10 (3%)	44	60

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	E	314/317 (99%)	301 (96%)	13 (4%)	35	48
All	All	1334/1342 (99%)	1294 (97%)	40 (3%)	46	63

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	96	THR
1	D	224	VAL
2	E	310	TRP
1	D	126	LEU
1	D	317	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	93	HIS
1	D	100	ASN
2	E	231	ASN
2	B	261	ASN
2	E	108	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SF4	A	575	1	0,12,12	0.00	-	0,24,24	0.00	-
4	SF4	A	576	1	0,12,12	0.00	-	0,24,24	0.00	-
3	SRM	A	580	2	34,70,70	1.93	7 (20%)	34,112,112	3.83	18 (52%)
5	PO4	A	590	-	4,4,4	0.92	0	6,6,6	0.82	0
3	SRM	B	570	1	34,70,70	2.49	10 (29%)	34,112,112	4.13	19 (55%)
4	SF4	B	585	2	0,12,12	0.00	-	0,24,24	0.00	-
4	SF4	B	586	2	0,12,12	0.00	-	0,24,24	0.00	-
4	SF4	D	575	1	0,12,12	0.00	-	0,24,24	0.00	-
4	SF4	D	576	1	0,12,12	0.00	-	0,24,24	0.00	-
3	SRM	D	580	-	34,70,70	1.95	9 (26%)	34,112,112	3.33	13 (38%)
3	SRM	E	570	-	34,70,70	2.73	10 (29%)	34,112,112	3.95	21 (61%)
4	SF4	E	585	2	0,12,12	0.00	-	0,24,24	0.00	-
4	SF4	E	586	2	0,12,12	0.00	-	0,24,24	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SF4	A	575	1	-	0/0/48/48	0/6/5/5
4	SF4	A	576	1	-	0/0/48/48	0/6/5/5
3	SRM	A	580	2	-	1/22/126/126	0/0/8/8
5	PO4	A	590	-	-	0/0/0/0	0/0/0/0
3	SRM	B	570	1	-	0/22/126/126	0/0/8/8
4	SF4	B	585	2	-	0/0/48/48	0/6/5/5
4	SF4	B	586	2	-	0/0/48/48	0/6/5/5
4	SF4	D	575	1	-	0/0/48/48	0/6/5/5
4	SF4	D	576	1	-	0/0/48/48	0/6/5/5
3	SRM	D	580	-	-	0/22/126/126	0/0/8/8
3	SRM	E	570	-	-	0/22/126/126	0/0/8/8
4	SF4	E	585	2	-	0/0/48/48	0/6/5/5
4	SF4	E	586	2	-	0/0/48/48	0/6/5/5

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	570	SRM	C4C-NC	-7.09	1.28	1.36
3	E	570	SRM	C1C-NC	-6.71	1.28	1.36
3	B	570	SRM	C4C-NC	-6.08	1.29	1.36
3	B	570	SRM	C4A-NA	-5.80	1.28	1.39
3	E	570	SRM	C4A-NA	-5.64	1.28	1.39

The worst 5 of 71 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	580	SRM	CAB-C3B-C2B	-15.26	106.27	123.52
3	B	570	SRM	CAA-C3A-C2A	-12.23	109.69	123.52
3	D	580	SRM	CAB-C3B-C2B	-11.94	110.02	123.52
3	E	570	SRM	CAA-C3A-C2A	-10.71	111.41	123.52
3	E	570	SRM	CDC-C2C-C1C	-9.58	112.92	127.39

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	580	SRM	C3D-C2D-CAD-CBD

There are no ring outliers.

7 monomers are involved in 40 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	580	SRM	12	0
5	A	590	PO4	2	0
3	B	570	SRM	6	0
4	D	575	SF4	1	0
3	D	580	SRM	6	0
3	E	570	SRM	11	0
4	E	585	SF4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	417/418 (99%)	0.39	29 (6%) 17 23	12, 19, 35, 78	0
1	D	417/418 (99%)	1.49	131 (31%) 0 0	13, 23, 36, 54	0
2	B	363/366 (99%)	0.35	17 (4%) 32 39	11, 18, 28, 63	0
2	E	363/366 (99%)	1.54	104 (28%) 1 1	6, 20, 53, 78	0
All	All	1560/1568 (99%)	0.94	281 (18%) 2 2	6, 20, 42, 78	0

The worst 5 of 281 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	67	ILE	8.8
2	E	6	VAL	8.3
2	E	4	GLU	8.1
2	E	207	ILE	7.8
2	E	229	MET	7.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	SRM	E	570	63/63	0.88	0.25	0.16	16,20,24,28	0
3	SRM	D	580	63/63	0.85	0.28	0.01	40,55,58,60	0
4	SF4	E	585	8/8	0.93	0.18	-0.67	70,74,77,78	0
3	SRM	A	580	63/63	0.95	0.15	-0.69	23,27,36,40	0
3	SRM	B	570	63/63	0.96	0.14	-0.94	13,16,20,23	0
4	SF4	D	575	8/8	0.93	0.13	-1.14	14,15,16,16	0
5	PO4	A	590	5/5	0.97	0.14	-1.25	59,59,61,63	0
4	SF4	E	586	8/8	0.96	0.12	-1.40	35,38,39,43	0
4	SF4	D	576	8/8	0.92	0.10	-1.49	17,20,21,22	0
4	SF4	A	575	8/8	0.98	0.10	-1.89	14,16,18,20	0
4	SF4	A	576	8/8	0.95	0.05	-2.64	13,14,15,16	0
4	SF4	B	586	8/8	0.98	0.06	-2.70	15,17,18,19	0
4	SF4	B	585	8/8	0.98	0.06	-2.74	21,24,28,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.