



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 19, 2017 – 06:56 PM EDT

PDB ID : 4MO1
Title : Crystal structure of antitermination protein Q from bacteriophage lambda.
Northeast Structural Genomics Consortium target OR18A.
Authors : Vorobiev, S.; Su, M.; Nickels, B.; Seetharaman, J.; Sahdev, S.; Xiao, R.;
Kogan, S.; Maglaqui, M.; Wang, D.; Everett, J.K.; Acton, T.B.; Ebright,
R.H.; Montelione, G.T.; Hunt, J.; Tong, L.; Northeast Structural Genomics
Consortium (NESG)
Deposited on : unknown
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	178	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: right;">%</div> <div style="position: absolute; top: 10px; left: 0; width: 100%; text-align: right;">75% 6% 19%</div> </div> </div>
1	B	178	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 10px; left: 0; width: 100%; text-align: right;">66% 14% 20%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ZN	B	301	-	-	-	X
3	BR	B	303	-	-	X	-
4	CL	A	303	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2372 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antitermination protein Q.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	145	Total	C	N	O	S	Se	0	0	0
			1104	693	201	201	6	3			
1	B	143	Total	C	N	O	S	Se	0	0	0
			1098	688	202	199	6	3			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	208	LEU	-	EXPRESSION TAG	UNP P03047
A	209	GLU	-	EXPRESSION TAG	UNP P03047
A	210	HIS	-	EXPRESSION TAG	UNP P03047
A	211	HIS	-	EXPRESSION TAG	UNP P03047
A	212	HIS	-	EXPRESSION TAG	UNP P03047
A	213	HIS	-	EXPRESSION TAG	UNP P03047
A	214	HIS	-	EXPRESSION TAG	UNP P03047
A	215	HIS	-	EXPRESSION TAG	UNP P03047
B	208	LEU	-	EXPRESSION TAG	UNP P03047
B	209	GLU	-	EXPRESSION TAG	UNP P03047
B	210	HIS	-	EXPRESSION TAG	UNP P03047
B	211	HIS	-	EXPRESSION TAG	UNP P03047
B	212	HIS	-	EXPRESSION TAG	UNP P03047
B	213	HIS	-	EXPRESSION TAG	UNP P03047
B	214	HIS	-	EXPRESSION TAG	UNP P03047
B	215	HIS	-	EXPRESSION TAG	UNP P03047

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total 2	Br 2	0	0
3	A	1	Total 1	Br 1	0	0


- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total 2	Cl 2	0	0
4	A	1	Total 1	Cl 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	84	Total 84	O 84	0	0
5	B	78	Total 78	O 78	0	0

- Molecule 1: Antitermination protein Q



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	54.47Å 54.47Å 109.85Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.35 – 2.10 43.35 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.3 (43.35-2.10) 99.5 (43.35-2.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.7.2_869	Depositor
R, R_{free}	0.214 , 0.249 0.194 , 0.237	Depositor DCC
R_{free} test set	1088 reflections (5.14%)	DCC
Wilson B-factor (Å ²)	32.4	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 49.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l 0.049 for h,-h-k,-l 0.029 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2372	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.80 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 6.5055e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BR, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/1121	0.60	0/1512
1	B	0.33	0/1115	0.59	0/1502
All	All	0.33	0/2236	0.59	0/3014

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1104	0	1105	4	0
1	B	1098	0	1105	13	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	2	0	0	2	0
4	A	1	0	0	0	0
4	B	2	0	0	0	0
5	A	84	0	0	2	0
5	B	78	0	0	3	0
All	All	2372	0	2210	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:ASN:O	1:A:72:GLN:HG2	1.90	0.71
1:B:68:ASN:O	1:B:72:GLN:HG2	1.99	0.62
1:B:86:LYS:HG2	5:B:408:HOH:O	2.07	0.54
1:B:62:ASP:HB3	1:B:65:LYS:HE3	1.92	0.51
1:B:71:MSE:HE2	1:B:97:GLN:HG3	1.91	0.51
1:A:173:GLN:HG2	5:A:459:HOH:O	2.10	0.51
1:B:108:CYS:SG	5:B:477:HOH:O	2.60	0.49
1:A:109:ARG:HD2	5:A:458:HOH:O	2.11	0.48
1:B:181:LYS:HB3	1:B:182:PRO:HD3	1.95	0.48
1:B:84:VAL:HA	1:B:87:LEU:HG	1.96	0.48
1:B:166:MSE:HE2	5:B:452:HOH:O	2.15	0.46
1:A:117:ARG:HD3	1:A:192:HIS:CE1	2.50	0.46
1:B:83:GLY:N	1:B:179:THR:HB	2.32	0.44
1:B:95:VAL:HG21	1:B:168:ILE:HD11	1.98	0.44
1:B:106:ASP:O	3:B:303:BR:BR	2.91	0.44
1:B:173:GLN:HB3	1:B:174:PRO:HD3	2.01	0.43
1:B:109:ARG:HB3	3:B:303:BR:BR	2.76	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	143/178 (80%)	139 (97%)	4 (3%)	0	100	100
1	B	141/178 (79%)	136 (96%)	4 (3%)	1 (1%)	25	20
All	All	284/356 (80%)	275 (97%)	8 (3%)	1 (0%)	38	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	63	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/135 (84%)	108 (96%)	5 (4%)	33	31
1	B	113/135 (84%)	108 (96%)	5 (4%)	33	31
All	All	226/270 (84%)	216 (96%)	10 (4%)	33	31

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	GLN
1	A	82	ARG
1	A	86	LYS
1	A	155	MSE
1	A	176	TRP
1	B	86	LYS
1	B	138	ARG
1	B	154	ARG
1	B	155	MSE
1	B	197	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	142/178 (79%)	-0.13	2 (1%) 75 79	21, 36, 61, 80	0
1	B	140/178 (78%)	-0.16	0 100 100	21, 35, 67, 94	0
All	All	282/356 (79%)	-0.14	2 (0%) 87 89	21, 35, 67, 94	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	206	THR	3.2
1	A	205	VAL	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	CL	A	303	1/1	0.99	0.16	7.81	29,29,29,29	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ZN	B	301	1/1	0.99	0.14	2.21	31,31,31,31	0
2	ZN	A	301	1/1	0.97	0.13	1.22	42,42,42,42	0
4	CL	B	305	1/1	0.98	0.08	-1.59	37,37,37,37	0
3	BR	B	302	1/1	0.75	0.33	-	51,51,51,51	1
4	CL	B	304	1/1	0.98	0.19	-	36,36,36,36	0
3	BR	B	303	1/1	0.99	0.17	-	35,35,35,35	1
3	BR	A	302	1/1	0.99	0.19	-	40,40,40,40	1

6.5 Other polymers [i](#)

There are no such residues in this entry.