



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 01:58 am GMT

PDB ID : 3MP3
Title : Crystal Structure of Human Lyase in complex with inhibitor HG-CoA
Authors : Fu, Z.; Runquist, J.A.; Montgomery, C.; Mizioroko, H.M.; Kim, J.-J.P.
Deposited on : 2010-04-24
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	<div> <div>63%</div> <div>32%</div> <div>• •</div> </div>
1	B	298	<div> <div>2%</div> <div>61%</div> <div>34%</div> <div>• •</div> </div>
1	C	298	<div> <div>53%</div> <div>42%</div> <div>• •</div> </div>
1	D	298	<div> <div>46%</div> <div>45%</div> <div>6%</div> <div>•</div> </div>
1	E	298	<div> <div>59%</div> <div>34%</div> <div>5%</div> <div>•</div> </div>
1	F	298	<div> <div>%</div> <div>48%</div> <div>46%</div> <div>6%</div> <div>•</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	3HG	A	400	-	-	-	X
3	MG	F	406	-	-	-	X

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 13597 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydroxymethylglutaryl-CoA lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	296	Total	C	N	O	S	0	0	0
			2189	1390	365	417	17			
1	B	296	Total	C	N	O	S	0	0	0
			2189	1390	365	417	17			
1	C	296	Total	C	N	O	S	0	0	0
			2189	1390	365	417	17			
1	D	289	Total	C	N	O	S	0	0	0
			2142	1361	357	408	16			
1	E	296	Total	C	N	O	S	0	0	0
			2189	1390	365	417	17			
1	F	296	Total	C	N	O	S	0	0	0
			2189	1390	365	417	17			

- Molecule 2 is 3-HYDROXPENTANEDIOIC ACID (three-letter code: 3HG) (formula: $C_5H_8O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	5	5		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	1	Total	Mg	0	0
			1	1		
3	E	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		
3	F	1	Total	Mg	0	0
			1	1		

- Molecule 4 is (3R,5S,9R,21S)-1-[(2R,3S,4R,5R)-5-(6-AMINO-9H-PURIN-9-YL)-4-HYDROXY-3-(PHOSPHONOOXY)TETRAHYDROFURAN-2-YL]-3,5,9,21-TETRAHYDROXY-8,8-DIMETHYL-10,14,19-TRIOXO-2,4,6-TRIOXA-18-THIA-11,15-DIAZA-3,5-DIPHOSPHATRICOSAN-23-OIC ACID 3,5-DIOXIDE (three-letter code: HGG) (formula: C₂₆H₄₂N₇O₂₀P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	S	0	0
			57	26	7	20	3	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	100	Total 100	O 100	0	0
5	B	70	Total 70	O 70	0	0
5	C	70	Total 70	O 70	0	0
5	D	65	Total 65	O 65	0	0
5	E	74	Total 74	O 74	0	0
5	F	58	Total 58	O 58	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Hydroxymethylglutaryl-CoA lyase




• Molecule 1: Hydroxymethylglutaryl-CoA lyase




• Molecule 1: Hydroxymethylglutaryl-CoA lyase






• Molecule 1: Hydroxymethylglutaryl-CoA lyase

Chain D: 46% 45% 6% •




• Molecule 1: Hydroxymethylglutaryl-CoA lyase

Chain E: 59% 34% 5% •



• Molecule 1: Hydroxymethylglutaryl-CoA lyase

Chain F: 48% 46% 6% •



G274	N275	L276	A277	T278	E279	V282	L288	G289	I290	H291	N295	L296	Q297	K298	L299	F305	I306	A309	I310	R311	R312	T314	S315	S316	A319	Q320	A321	T322	C323	LYS	LEU
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	197.37Å 117.00Å 86.87Å 90.00° 112.31° 90.00°	Depositor
Resolution (Å)	29.70 – 2.40 39.17 – 2.39	Depositor EDS
% Data completeness (in resolution range)	93.3 (29.70-2.40) 92.5 (39.17-2.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 2.39Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.212 , 0.281 0.212 , 0.280	Depositor DCC
R_{free} test set	6749 reflections (10.14%)	DCC
Wilson B-factor (Å ²)	22.8	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 22.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	13597	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HGG, MG, 3HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/2224	0.64	0/3011
1	B	0.39	0/2224	0.63	0/3011
1	C	0.39	0/2224	0.63	0/3011
1	D	0.38	0/2174	0.60	0/2941
1	E	0.38	0/2224	0.62	0/3011
1	F	0.36	0/2224	0.58	0/3011
All	All	0.38	0/13294	0.62	0/17996

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2189	0	2244	92	0
1	B	2189	0	2244	93	0
1	C	2189	0	2244	139	0
1	D	2142	0	2203	136	0
1	E	2189	0	2244	104	0
1	F	2189	0	2244	138	0
2	A	10	0	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
4	B	57	0	37	10	0
5	A	100	0	0	6	0
5	B	70	0	0	3	0
5	C	70	0	0	7	0
5	D	65	0	0	3	0
5	E	74	0	0	4	0
5	F	58	0	0	3	0
All	All	13597	0	13466	686	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 686 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:47:GLU:HG2	1:D:310:LEU:HD21	1.46	0.98
1:E:211:PRO:HG2	1:F:288:LEU:HD11	1.44	0.96
1:D:263:LEU:HD11	1:D:306:ILE:HD12	1.48	0.96
1:E:84:GLN:NE2	1:E:84:GLN:H	1.65	0.94
1:E:51:VAL:HB	1:E:55:VAL:HG21	1.48	0.93

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/298 (99%)	272 (92%)	16 (5%)	6 (2%)	9	10
1	B	294/298 (99%)	272 (92%)	16 (5%)	6 (2%)	9	10
1	C	294/298 (99%)	263 (90%)	27 (9%)	4 (1%)	13	18
1	D	285/298 (96%)	254 (89%)	25 (9%)	6 (2%)	8	9
1	E	294/298 (99%)	273 (93%)	15 (5%)	6 (2%)	9	10
1	F	294/298 (99%)	249 (85%)	35 (12%)	10 (3%)	4	3
All	All	1755/1788 (98%)	1583 (90%)	134 (8%)	38 (2%)	8	9

5 of 38 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	139	ILE
1	A	140	ASN
1	A	260	VAL
1	A	270	GLN
1	A	272	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/240 (99%)	220 (92%)	18 (8%)	15	24
1	B	238/240 (99%)	223 (94%)	15 (6%)	21	33
1	C	238/240 (99%)	222 (93%)	16 (7%)	19	30
1	D	234/240 (98%)	218 (93%)	16 (7%)	18	29
1	E	238/240 (99%)	217 (91%)	21 (9%)	12	17
1	F	238/240 (99%)	224 (94%)	14 (6%)	23	36
All	All	1424/1440 (99%)	1324 (93%)	100 (7%)	18	28

5 of 100 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	268	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	114	GLU
1	F	192	PHE
1	C	299	LEU
1	D	44	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 54 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	250	GLN
1	D	161	ASN
1	F	250	GLN
1	C	311	ASN
1	D	88	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	3HG	A	400	3	3,9,9	2.55	2 (66%)	2,11,11	1.49	0
4	HGG	B	399	3	48,59,59	3.67	23 (47%)	54,87,87	3.58	19 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3HG	A	400	3	-	0/4/8/8	0/0/0/0
4	HGG	B	399	3	-	0/52/75/75	0/3/3/3

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	399	HGG	C57-C45	-2.70	1.47	1.53
4	B	399	HGG	C62-N68	-2.17	1.41	1.46
4	B	399	HGG	C73-C74	-2.05	1.41	1.50
4	B	399	HGG	C16-C15	2.02	1.56	1.53
2	A	400	3HG	C2-C3	2.09	1.55	1.53

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	399	HGG	N3-C2-N1	-11.72	118.65	128.86
4	B	399	HGG	O1'-C1'-S81	-8.45	113.99	122.84
4	B	399	HGG	C15-N9-C4	-4.00	119.72	126.64
4	B	399	HGG	O22-C18-C19	-3.98	95.95	109.40
4	B	399	HGG	C53-C45-C46	-3.08	103.84	108.37

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	399	HGG	10	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	296/298 (99%)	-0.81	0 100 100	8, 21, 44, 60	0
1	B	296/298 (99%)	-0.62	5 (1%) 70 68	10, 27, 59, 74	0
1	C	296/298 (99%)	-0.66	0 100 100	10, 30, 52, 66	0
1	D	289/298 (96%)	-0.48	1 (0%) 93 93	17, 37, 62, 79	0
1	E	296/298 (99%)	-0.73	0 100 100	11, 26, 56, 69	0
1	F	296/298 (99%)	-0.39	2 (0%) 87 86	16, 42, 64, 73	0
All	All	1769/1788 (98%)	-0.61	8 (0%) 90 89	8, 29, 59, 79	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	268	TYR	3.1
1	B	269	ALA	2.9
1	B	272	ALA	2.8
1	F	268	TYR	2.6
1	B	270	GLN	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	3HG	A	400	10/10	0.75	0.26	3.69	61,63,66,67	0
3	MG	F	406	1/1	0.97	0.16	3.43	30,30,30,30	0
4	HGG	B	399	57/57	0.89	0.17	1.78	58,79,96,97	0
3	MG	A	403	1/1	0.97	0.10	-	16,16,16,16	0
3	MG	E	402	1/1	0.99	0.18	-	13,13,13,13	0
3	MG	C	404	1/1	0.97	0.17	-	15,15,15,15	0
3	MG	B	401	1/1	0.98	0.14	-	17,17,17,17	0
3	MG	D	405	1/1	0.98	0.09	-	28,28,28,28	0

6.5 Other polymers

There are no such residues in this entry.