



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 9, 2017 – 11:02 am GMT

PDB ID : 5MQ6
Title : Polycyclic Ketone Monooxygenase from the Thermophilic Fungus *Thermothelomyces thermophila*
Authors : Savino, S.; Furst, M.J.L.J.; Fraaije, M.W.; Mattevi, A.
Deposited on : 2016-12-20
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	recalc28906
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28906


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	655	<div> <div></div> <div>83%</div> <div>11%</div> <div>• •</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5433 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyridine nucleotide-disulfide oxidoreductase-like protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	636	4970	3155	866	934	15	0	2	0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			39	15	5	16	3		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	365	Total 365	O 365	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyridine nucleotide-disulfide oxidoreductase-like protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.20Å 91.02Å 133.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	75.21 – 2.00 75.21 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (75.21-2.00) 100.0 (75.21-2.00)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0151	Depositor
R, R_{free}	0.153 , 0.231 0.164 , 0.236	Depositor DCC
R_{free} test set	1541 reflections (3.21%)	DCC
Wilson B-factor (Å ²)	25.5	Xtriage
Anisotropy	0.025	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 55.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5433	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FAD, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.10	8/5104 (0.2%)	1.16	29/6926 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	338	SER	CB-OG	6.06	1.50	1.42
1	A	475	SER	CB-OG	-5.97	1.34	1.42
1	A	532	LEU	C-O	-5.71	1.12	1.23
1	A	402	ARG	CD-NE	-5.43	1.37	1.46
1	A	400	ARG	CD-NE	-5.42	1.37	1.46
1	A	397	GLU	CD-OE2	5.28	1.31	1.25
1	A	198	GLU	CD-OE1	5.12	1.31	1.25
1	A	135	GLU	CD-OE2	5.04	1.31	1.25

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	400	ARG	NE-CZ-NH2	-20.08	110.26	120.30
1	A	400	ARG	NE-CZ-NH1	16.91	128.75	120.30
1	A	402	ARG	NE-CZ-NH2	-16.59	112.01	120.30
1	A	402	ARG	NE-CZ-NH1	15.36	127.98	120.30
1	A	643	ASP	CB-CG-OD2	-8.33	110.80	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	332	ARG	NE-CZ-NH2	7.83	124.22	120.30
1	A	33	ARG	NE-CZ-NH1	7.82	124.21	120.30
1	A	643	ASP	CB-CG-OD1	7.70	125.23	118.30
1	A	400	ARG	CD-NE-CZ	7.62	134.26	123.60
1	A	324	ARG	NE-CZ-NH1	7.59	124.09	120.30
1	A	300	ARG	NE-CZ-NH2	-7.14	116.73	120.30
1	A	332	ARG	NE-CZ-NH1	-7.10	116.75	120.30
1	A	203	ARG	NE-CZ-NH1	6.89	123.75	120.30
1	A	402	ARG	CD-NE-CZ	6.79	133.11	123.60
1	A	96	ARG	NE-CZ-NH1	6.67	123.64	120.30
1	A	96	ARG	NE-CZ-NH2	-6.09	117.25	120.30
1	A	166	ARG	NE-CZ-NH2	-6.02	117.29	120.30
1	A	253	ASP	CB-CG-OD1	6.00	123.70	118.30
1	A	566	ASP	CB-CG-OD1	5.99	123.69	118.30
1	A	300	ARG	NE-CZ-NH1	5.95	123.28	120.30
1	A	404	ASP	CB-CG-OD2	5.78	123.50	118.30
1	A	220	LEU	CB-CG-CD1	5.68	120.66	111.00
1	A	641	ARG	NE-CZ-NH2	5.58	123.09	120.30
1	A	284	VAL	CG1-CB-CG2	-5.29	102.44	110.90
1	A	597	THR	CB-CA-C	-5.22	97.51	111.60
1	A	166	ARG	NE-CZ-NH1	5.19	122.90	120.30
1	A	217	ARG	NE-CZ-NH2	-5.15	117.73	120.30
1	A	221	LEU	CB-CG-CD2	5.13	119.73	111.00
1	A	561	ASP	CB-CG-OD1	5.05	122.85	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	20	HIS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4970	0	4875	50	1
2	A	53	0	31	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	39	0	18	2	0
4	A	6	0	8	0	0
5	A	365	0	0	9	2
All	All	5433	0	4932	50	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132[B]:CYS:SG	2:A:700:FAD:O4	1.96	1.23
1:A:193:ARG:NH1	1:A:654:ARG:O	1.93	1.01
1:A:132[B]:CYS:HG	2:A:700:FAD:C4	1.83	0.92
1:A:255:THR:HG22	1:A:256:GLY:O	1.70	0.90
1:A:300:ARG:HD3	3:A:701:NDP:O2X	1.75	0.86
1:A:132[B]:CYS:SG	2:A:700:FAD:C4	2.65	0.83
1:A:488:PRO:O	1:A:492:THR:HG23	1.79	0.82
1:A:223[A]:SER:OG	1:A:492:THR:HB	1.79	0.80
1:A:563:VAL:O	5:A:801:HOH:O	2.01	0.77
1:A:597:THR:HG21	5:A:1134:HOH:O	1.97	0.64
1:A:271:ARG:NH1	1:A:468:PRO:O	2.29	0.64
1:A:280:THR:O	1:A:284:VAL:HG23	1.97	0.63
1:A:456:ASN:HB2	1:A:457:PRO:HD2	1.80	0.62
1:A:23:SER:O	1:A:27:THR:HG23	2.01	0.61
1:A:404:ASP:OD2	1:A:417:LYS:NZ	2.28	0.60
1:A:250:TRP:HE1	1:A:255:THR:HG21	1.66	0.59
1:A:409:ASP:HB3	1:A:412:THR:HG23	1.85	0.58
1:A:271:ARG:HG3	1:A:470:ASP:OD2	2.05	0.56
1:A:223[A]:SER:CB	1:A:492:THR:HB	2.37	0.54
1:A:366:ILE:HD12	1:A:537:GLN:HG3	1.90	0.54
1:A:597:THR:CG2	5:A:1134:HOH:O	2.56	0.53
1:A:255:THR:HG23	1:A:264:LEU:HG	1.91	0.53
1:A:378:PRO:O	5:A:802:HOH:O	2.19	0.51
1:A:397:GLU:OE2	5:A:803:HOH:O	2.19	0.51
1:A:223[B]:SER:CB	1:A:492:THR:HB	2.40	0.50
1:A:253:ASP:O	1:A:266:ARG:NH1	2.41	0.50
1:A:456:ASN:HB2	1:A:457:PRO:CD	2.42	0.50
1:A:38:GLY:O	1:A:41:GLN:HG2	2.12	0.50
1:A:409:ASP:HB3	1:A:412:THR:CG2	2.43	0.48
1:A:320:THR:HG22	1:A:321:LYS:HG3	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:33:ARG:HB2	5:A:947:HOH:O	2.15	0.46
1:A:404:ASP:HB3	5:A:837:HOH:O	2.15	0.46
1:A:453:ASP:HB2	1:A:461:VAL:O	2.16	0.45
1:A:50:ASN:OD1	1:A:52:ARG:HB3	2.16	0.45
1:A:84:LEU:HD22	1:A:223[A]:SER:HB2	1.99	0.45
1:A:400:ARG:HD2	5:A:1068:HOH:O	2.17	0.44
1:A:593:VAL:HG13	1:A:602:THR:HG22	1.98	0.44
1:A:311:ARG:HD3	5:A:1114:HOH:O	2.17	0.44
1:A:223[B]:SER:HB2	1:A:492:THR:HB	2.00	0.43
1:A:223[B]:SER:OG	1:A:492:THR:HG22	2.18	0.43
1:A:454:ALA:HB3	1:A:461:VAL:HG13	2.01	0.42
1:A:453:ASP:OD2	1:A:463:ALA:N	2.52	0.41
1:A:426:ARG:HD3	3:A:701:NDP:H2D	2.01	0.41
1:A:330:ARG:HA	1:A:333:MET:HE2	2.02	0.41
1:A:43:ILE:HB	1:A:233:LYS:HE3	2.02	0.41
1:A:393:LEU:HB3	1:A:394:PRO:HD3	2.03	0.41
1:A:181:SER:O	2:A:700:FAD:H2A	2.21	0.41
1:A:33:ARG:HG3	1:A:598:PRO:HD2	2.03	0.40
1:A:221:LEU:HD13	1:A:492:THR:HG21	2.02	0.40
1:A:468:PRO:O	1:A:469:LEU:HD13	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:75:ASP:OD2	5:A:801:HOH:O[4_465]	1.96	0.24
5:A:801:HOH:O	5:A:914:HOH:O[4_565]	2.12	0.08

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	634/655 (97%)	608 (96%)	25 (4%)	1 (0%)	51	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	566	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	516/527 (98%)	489 (95%)	27 (5%)	27	22

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	GLU
1	A	18	LYS
1	A	27	THR
1	A	52	ARG
1	A	53	PHE
1	A	193	ARG
1	A	220	LEU
1	A	238	GLU
1	A	255	THR
1	A	264	LEU
1	A	316	VAL
1	A	319	LYS
1	A	325	LYS
1	A	346	GLU
1	A	402	ARG
1	A	404	ASP
1	A	412	THR
1	A	450	LYS
1	A	457	PRO
1	A	461	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	465	LYS
1	A	466	GLU
1	A	492	THR
1	A	593	VAL
1	A	597	THR
1	A	601	ILE
1	A	615	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	700	-	51,58,58	1.71	9 (17%)	54,89,89	2.37	12 (22%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NDP	A	701	-	36,42,52	1.18	3 (8%)	40,65,80	1.88	8 (20%)
4	GOL	A	702	-	5,5,5	0.59	0	5,5,5	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	700	-	-	0/28/50/50	0/6/6/6
3	NDP	A	701	-	-	0/23/56/77	0/4/4/5
4	GOL	A	702	-	-	0/4/4/4	0/0/0/0

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	701	NDP	O4D-C1D	2.12	1.48	1.43
2	A	700	FAD	C5A-C4A	2.22	1.45	1.40
3	A	701	NDP	O3D-C3D	2.29	1.48	1.43
2	A	700	FAD	O4'-C4'	3.06	1.50	1.43
3	A	701	NDP	C5A-C4A	3.12	1.47	1.40
2	A	700	FAD	C4X-N5	3.21	1.37	1.33
2	A	700	FAD	C4-C4X	3.36	1.47	1.41
2	A	700	FAD	C4X-C10	3.38	1.46	1.41
2	A	700	FAD	C8-C7	3.44	1.49	1.41
2	A	700	FAD	C6-C7	3.84	1.48	1.37
2	A	700	FAD	C9A-C5X	4.10	1.51	1.42
2	A	700	FAD	C6-C5X	4.32	1.48	1.41

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	700	FAD	C4-C4X-C10	-9.85	112.00	119.96
3	A	701	NDP	N3A-C2A-N1A	-7.06	122.71	128.86
2	A	700	FAD	C4A-C5A-N7A	-3.85	105.69	109.41
3	A	701	NDP	C1B-N9A-C4A	-3.71	120.22	126.64
2	A	700	FAD	C7-C6-C5X	-3.35	115.89	121.08
3	A	701	NDP	C4B-O4B-C1B	-2.61	106.99	109.77
2	A	700	FAD	O3'-C3'-C4'	-2.34	103.03	108.82
3	A	701	NDP	O2A-PA-O1A	2.06	122.95	112.28
3	A	701	NDP	O3X-P2B-O1X	2.09	118.68	110.50
2	A	700	FAD	O2P-P-O1P	2.18	123.55	112.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	700	FAD	O4'-C4'-C3'	2.47	115.23	109.09
2	A	700	FAD	C5A-C6A-N6A	2.52	125.61	120.47
3	A	701	NDP	N6A-C6A-N1A	2.83	124.39	118.77
2	A	700	FAD	C5X-C9A-N10	2.85	119.78	117.66
2	A	700	FAD	C2A-N1A-C6A	2.92	123.87	118.77
3	A	701	NDP	C2A-N1A-C6A	2.95	123.94	118.77
3	A	701	NDP	O3D-C3D-C2D	3.62	120.57	111.91
2	A	700	FAD	C1'-N10-C9A	4.81	122.75	118.35
2	A	700	FAD	C4-N3-C2	5.59	120.05	115.16
2	A	700	FAD	C4-C4X-N5	6.18	125.45	118.68

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	700	FAD	4	0
3	A	701	NDP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	636/655 (97%)	-0.28	9 (1%) 75 75	14, 28, 61, 79	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	599	GLY	4.0
1	A	602	THR	3.1
1	A	600	TYR	2.9
1	A	603	SER	2.4
1	A	605	GLY	2.3
1	A	601	ILE	2.3
1	A	45	LEU	2.2
1	A	38	GLY	2.2
1	A	346	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	GOL	A	702	6/6	0.98	0.11	0.49	28,29,32,32	0
3	NDP	A	701	39/48	0.97	0.14	0.46	30,42,51,58	0
2	FAD	A	700	53/53	0.98	0.10	-0.32	14,24,37,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.