



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 03:26 am GMT

PDB ID : 3MSY  
Title : Crystal Structure of Mandelate racemase/muconate lactonizing enzyme from a Marine actinobacterium  
Authors : Satyanarayana, L.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2010-04-29  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	379	<div> <div></div> <div>53% 35% 5% 8%</div> </div>
1	B	379	<div> <div>%</div> <div>53% 34% 5% 7%</div> </div>
1	C	379	<div> <div>%</div> <div>54% 35% • 6%</div> </div>
1	D	379	<div> <div>%</div> <div>55% 34% • 7%</div> </div>
1	E	379	<div> <div>%</div> <div>50% 38% 5% 8%</div> </div>
1	F	379	<div> <div>%</div> <div>53% 36% • 7%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 16869 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mandelate racemase/muconate lactonizing enzyme.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	349	Total	C	N	O	S	Se	0	0	0
			2731	1723	481	512	5	10			
1	B	351	Total	C	N	O	S	Se	0	0	0
			2747	1732	485	515	5	10			
1	C	355	Total	C	N	O	S	Se	0	0	0
			2786	1757	493	521	5	10			
1	D	351	Total	C	N	O	S	Se	0	0	0
			2747	1732	485	515	5	10			
1	E	350	Total	C	N	O	S	Se	0	0	0
			2737	1726	482	514	5	10			
1	F	351	Total	C	N	O	S	Se	0	0	0
			2747	1732	485	515	5	10			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	MSE	-	expression tag	UNP A4AFX2
A	19	SER	-	expression tag	UNP A4AFX2
A	389	GLU	-	expression tag	UNP A4AFX2
A	390	GLY	-	expression tag	UNP A4AFX2
A	391	HIS	-	expression tag	UNP A4AFX2
A	392	HIS	-	expression tag	UNP A4AFX2
A	393	HIS	-	expression tag	UNP A4AFX2
A	394	HIS	-	expression tag	UNP A4AFX2
A	395	HIS	-	expression tag	UNP A4AFX2
A	396	HIS	-	expression tag	UNP A4AFX2
B	18	MSE	-	expression tag	UNP A4AFX2
B	19	SER	-	expression tag	UNP A4AFX2
B	389	GLU	-	expression tag	UNP A4AFX2
B	390	GLY	-	expression tag	UNP A4AFX2
B	391	HIS	-	expression tag	UNP A4AFX2
B	392	HIS	-	expression tag	UNP A4AFX2
B	393	HIS	-	expression tag	UNP A4AFX2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	394	HIS	-	expression tag	UNP A4AFX2
B	395	HIS	-	expression tag	UNP A4AFX2
B	396	HIS	-	expression tag	UNP A4AFX2
C	18	MSE	-	expression tag	UNP A4AFX2
C	19	SER	-	expression tag	UNP A4AFX2
C	389	GLU	-	expression tag	UNP A4AFX2
C	390	GLY	-	expression tag	UNP A4AFX2
C	391	HIS	-	expression tag	UNP A4AFX2
C	392	HIS	-	expression tag	UNP A4AFX2
C	393	HIS	-	expression tag	UNP A4AFX2
C	394	HIS	-	expression tag	UNP A4AFX2
C	395	HIS	-	expression tag	UNP A4AFX2
C	396	HIS	-	expression tag	UNP A4AFX2
D	18	MSE	-	expression tag	UNP A4AFX2
D	19	SER	-	expression tag	UNP A4AFX2
D	389	GLU	-	expression tag	UNP A4AFX2
D	390	GLY	-	expression tag	UNP A4AFX2
D	391	HIS	-	expression tag	UNP A4AFX2
D	392	HIS	-	expression tag	UNP A4AFX2
D	393	HIS	-	expression tag	UNP A4AFX2
D	394	HIS	-	expression tag	UNP A4AFX2
D	395	HIS	-	expression tag	UNP A4AFX2
D	396	HIS	-	expression tag	UNP A4AFX2
E	18	MSE	-	expression tag	UNP A4AFX2
E	19	SER	-	expression tag	UNP A4AFX2
E	389	GLU	-	expression tag	UNP A4AFX2
E	390	GLY	-	expression tag	UNP A4AFX2
E	391	HIS	-	expression tag	UNP A4AFX2
E	392	HIS	-	expression tag	UNP A4AFX2
E	393	HIS	-	expression tag	UNP A4AFX2
E	394	HIS	-	expression tag	UNP A4AFX2
E	395	HIS	-	expression tag	UNP A4AFX2
E	396	HIS	-	expression tag	UNP A4AFX2
F	18	MSE	-	expression tag	UNP A4AFX2
F	19	SER	-	expression tag	UNP A4AFX2
F	389	GLU	-	expression tag	UNP A4AFX2
F	390	GLY	-	expression tag	UNP A4AFX2
F	391	HIS	-	expression tag	UNP A4AFX2
F	392	HIS	-	expression tag	UNP A4AFX2
F	393	HIS	-	expression tag	UNP A4AFX2
F	394	HIS	-	expression tag	UNP A4AFX2
F	395	HIS	-	expression tag	UNP A4AFX2

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	396	HIS	-	expression tag	UNP A4AFX2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	65	Total O 65 65	0	0
2	B	62	Total O 62 62	0	0
2	C	63	Total O 63 63	0	0
2	D	72	Total O 72 72	0	0
2	E	55	Total O 55 55	0	0
2	F	57	Total O 57 57	0	0







- Molecule 1: Mandelate racemase/muconate lactonizing enzyme




- Molecule 1: Mandelate racemase/muconate lactonizing enzyme





- Molecule 1: Mandelate racemase/muconate lactonizing enzyme





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.00Å 103.39Å 234.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.31 – 2.50 47.31 – 2.22	Depositor EDS
% Data completeness (in resolution range)	94.6 (47.31-2.50) 96.0 (47.31-2.22)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.76	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.58 (at 2.22Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.238 , 0.268 0.225 , 0.269	Depositor DCC
$R_{free}$ test set	3050 reflections (3.99%)	DCC
Wilson B-factor (Å <sup>2</sup> )	21.6	Xtriage
Anisotropy	0.508	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 41.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	16869	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/2790	0.67	0/3779
1	B	0.38	0/2807	0.67	1/3802 (0.0%)
1	C	0.38	0/2847	0.67	1/3856 (0.0%)
1	D	0.38	0/2807	0.68	0/3802
1	E	0.37	0/2796	0.69	0/3787
1	F	0.36	0/2807	0.65	1/3802 (0.0%)
All	All	0.37	0/16854	0.67	3/22828 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	68	GLY	N-CA-C	6.79	130.08	113.10
1	B	30	MSE	CB-CG-SE	-5.69	95.63	112.70
1	C	363	LEU	CA-CB-CG	5.17	127.19	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2731	0	2624	170	0
1	B	2747	0	2636	183	0
1	C	2786	0	2675	200	0
1	D	2747	0	2636	148	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2737	0	2629	208	0
1	F	2747	0	2636	151	0
2	A	65	0	0	19	0
2	B	62	0	0	14	0
2	C	63	0	0	16	0
2	D	72	0	0	6	0
2	E	55	0	0	30	0
2	F	57	0	0	9	0
All	All	16869	0	15836	1015	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

The worst 5 of 1015 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:376:ASN:HD21	1:E:378:ASP:HB2	1.16	1.06
1:D:67:THR:HG23	1:D:293:SER:HB2	1.36	1.05
1:E:30:MSE:HE2	1:E:89:LEU:HB3	1.35	1.03
1:C:253:MSE:HE3	1:C:264:VAL:HG12	1.42	1.00
1:C:37:GLU:H	1:C:37:GLU:CD	1.66	0.99

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/379 (91%)	326 (94%)	16 (5%)	3 (1%)	20	36
1	B	347/379 (92%)	323 (93%)	22 (6%)	2 (1%)	28	48
1	C	351/379 (93%)	332 (95%)	17 (5%)	2 (1%)	28	48

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	347/379 (92%)	325 (94%)	18 (5%)	4 (1%)	15	27
1	E	346/379 (91%)	315 (91%)	26 (8%)	5 (1%)	13	23
1	F	347/379 (92%)	327 (94%)	18 (5%)	2 (1%)	28	48
All	All	2083/2274 (92%)	1948 (94%)	117 (6%)	18 (1%)	20	36

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	33	PRO
1	B	69	ASP
1	D	61	ILE
1	E	69	ASP
1	C	33	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/297 (94%)	255 (91%)	25 (9%)	11	22
1	B	282/297 (95%)	259 (92%)	23 (8%)	13	25
1	C	286/297 (96%)	263 (92%)	23 (8%)	14	27
1	D	282/297 (95%)	265 (94%)	17 (6%)	22	41
1	E	281/297 (95%)	265 (94%)	16 (6%)	24	44
1	F	282/297 (95%)	262 (93%)	20 (7%)	17	32
All	All	1693/1782 (95%)	1569 (93%)	124 (7%)	16	31

5 of 124 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	141	LEU
1	C	381	ASP
1	F	240	PHE
1	C	184	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	279	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 53 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	318	HIS
1	D	126	ASN
1	F	247	HIS
1	C	359	ASN
1	C	376	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	339/379 (89%)	-0.27	0 100 100	10, 22, 37, 49	0
1	B	341/379 (89%)	-0.25	4 (1%) 79 80	10, 22, 37, 52	0
1	C	345/379 (91%)	-0.23	3 (0%) 84 85	10, 23, 38, 52	0
1	D	341/379 (89%)	-0.21	4 (1%) 79 80	10, 21, 36, 57	0
1	E	340/379 (89%)	0.04	3 (0%) 84 85	12, 27, 44, 61	0
1	F	341/379 (89%)	-0.06	4 (1%) 79 80	12, 26, 43, 57	0
All	All	2047/2274 (90%)	-0.16	18 (0%) 84 85	10, 23, 40, 61	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	68	GLY	3.3
1	F	42	HIS	2.9
1	B	46	THR	2.9
1	B	42	HIS	2.8
1	B	378	ASP	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.