



Full wwPDB NMR Structure Validation Report i

Feb 12, 2017 – 06:53 pm GMT

PDB ID : 1MUZ
Title : NMR STRUCTURE OF THE TUMOR SUPPRESSOR BIN1: ALTERNATIVE SPLICING IN MELANOMA AND INTERACTION WITH C-MYC
Authors : Pineda-Lucena, A.; Arrowsmith, C.H.
Deposited on : 2002-09-24

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) i) were used in the production of this report:


Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	trunk28760
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain				
1	A	81	27%	41%	12%	•	19%

2 Ensemble composition and analysis i

This entry contains 20 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:407-A:460, A:470-A:481 (66)	0.13	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 6, 7, 8, 12, 13, 15, 19, 20
2	5, 10, 17
3	9, 14, 16
Single-model clusters	11; 18

3 Entry composition [\(i\)](#)

There is only 1 type of molecule in this entry. The entry contains 1287 atoms, of which 629 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Myc box dependent interacting protein 1.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	81	1287	418	629	111	126	3	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	465	LYS	GLU	SEE REMARK 999	UNP O00499

4 Residue-property plots [\(i\)](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Myc box dependent interacting protein 1




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Myc box dependent interacting protein 1



4.2.2 Score per residue for model 2


- Molecule 1: Myc box dependent interacting protein 1






4.2.3 Score per residue for model 3

- Molecule 1: Myc box dependent interacting protein 1




4.2.4 Score per residue for model 4

- Molecule 1: Myc box dependent interacting protein 1




4.2.5 Score per residue for model 5 (medoid)

- Molecule 1: Myc box dependent interacting protein 1




4.2.6 Score per residue for model 6

- Molecule 1: Myc box dependent interacting protein 1




4.2.7 Score per residue for model 7

- Molecule 1: Myc box dependent interacting protein 1




4.2.8 Score per residue for model 8

- Molecule 1: Myc box dependent interacting protein 1




4.2.9 Score per residue for model 9

- Molecule 1: Myc box dependent interacting protein 1




4.2.10 Score per residue for model 10

- Molecule 1: Myc box dependent interacting protein 1




4.2.11 Score per residue for model 11

- Molecule 1: Myc box dependent interacting protein 1




4.2.12 Score per residue for model 12

- Molecule 1: Myc box dependent interacting protein 1




4.2.13 Score per residue for model 13

- Molecule 1: Myc box dependent interacting protein 1




4.2.14 Score per residue for model 14

- Molecule 1: Myc box dependent interacting protein 1




4.2.15 Score per residue for model 15

- Molecule 1: Myc box dependent interacting protein 1




4.2.16 Score per residue for model 16

- Molecule 1: Myc box dependent interacting protein 1




4.2.17 Score per residue for model 17

- Molecule 1: Myc box dependent interacting protein 1




4.2.18 Score per residue for model 18

- Molecule 1: Myc box dependent interacting protein 1




4.2.19 Score per residue for model 19

- Molecule 1: Myc box dependent interacting protein 1



4.2.20 Score per residue for model 20

- Molecule 1: Myc box dependent interacting protein 1



5 Refinement protocol and experimental data overview i

The models were refined using the following method: *TORSION ANGLE DYNAMICS*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	1.5
DYANA	refinement	1.5

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	535	498	498	48±5
All	All	10700	9960	9960	970

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 47.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:430:LEU:HD11	1:A:436:VAL:HG21	1.01	1.31	5	17
1:A:430:LEU:HD11	1:A:436:VAL:CG2	0.95	1.90	15	17
1:A:416:ALA:HB2	1:A:430:LEU:HD12	0.94	1.39	18	8
1:A:477:PHE:CD1	1:A:478:THR:HG23	0.92	1.99	16	20
1:A:420:TYR:CE2	1:A:422:ALA:HB2	0.84	2.07	14	20
1:A:414:VAL:HG22	1:A:436:VAL:O	0.81	1.73	12	9
1:A:420:TYR:CZ	1:A:422:ALA:HB2	0.81	2.11	6	20
1:A:455:VAL:HG22	1:A:459:ASP:HB2	0.80	1.50	11	16
1:A:451:TRP:C	1:A:452:LEU:HD23	0.80	1.97	1	4
1:A:452:LEU:HD11	1:A:475:GLU:HB2	0.79	1.53	16	10
1:A:416:ALA:HB2	1:A:430:LEU:HD23	0.77	1.56	20	3
1:A:455:VAL:HG22	1:A:459:ASP:CB	0.77	2.08	9	9
1:A:416:ALA:HA	1:A:478:THR:HG22	0.76	1.56	16	19
1:A:437:LEU:HD11	1:A:460:TRP:HB2	0.76	1.58	14	5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	437	LEU	20
1	A	451	TRP	19
1	A	458	SER	19
1	A	453	MET	16
1	A	443	ASN	16
1	A	442	GLN	11
1	A	447	GLN	11
1	A	475	GLU	11
1	A	428	LEU	10
1	A	476	ASN	9
1	A	445	GLU	9
1	A	427	GLU	8
1	A	449	GLU	7
1	A	456	LYS	7
1	A	419	ASP	7
1	A	479	GLU	6
1	A	415	GLN	5
1	A	480	ARG	5
1	A	478	THR	5
1	A	430	LEU	4
1	A	470	ARG	4
1	A	446	GLU	4
1	A	435	VAL	3
1	A	472	VAL	3
1	A	424	ASP	3
1	A	426	ASP	3
1	A	414	VAL	2
1	A	459	ASP	1
1	A	420	TYR	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

6.7 Other polymers [\(i\)](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

7 Chemical shift validation [\(i\)](#)

No chemical shift data were provided